



HOKKAIDO UNIVERSITY

Title	The Helmholtz decomposition of a space of vector fields with bounded mean oscillation in a bounded domain
Author(s)	Giga, Yoshikazu; Gu, Zhongyang
Citation	Hokkaido University Preprint Series in Mathematics, 1139, 1-42
Issue Date	2021-10-14
DOI	https://doi.org/10.14943/100096
Doc URL	https://hdl.handle.net/2115/82938
Type	departmental bulletin paper
File Information	TheHelDec.pdf



The Helmholtz decomposition of a space of vector fields with bounded mean oscillation in a bounded domain

Yoshikazu Giga
The University of Tokyo
labgiga@ms.u-tokyo.ac.jp

Zhongyang Gu
The University of Tokyo
zgu@ms.u-tokyo.ac.jp

Abstract

We introduce a space of vector fields with bounded mean oscillation whose “tangential” and “normal” components to the boundary behave differently. We establish its Helmholtz decomposition when the domain is bounded. This substantially extends the authors’ earlier result for a half space.

Keywords: Helmholtz decomposition, *BMO* space, Neumann problem, normal component.

1 Introduction

The Helmholtz decomposition of a vector field is a fundamental tool to analyze the Stokes and the Navier-Stokes equations. It is formally a decomposition of a vector field $v = (v^1, \dots, v^n)$ in a domain Ω of $\mathbf{R}^n$ into

$$v = v_0 + \nabla q; \tag{1}$$

here v_0 is a divergence free vector field satisfying supplemental conditions like boundary condition and ∇q denotes the gradient of a function (scalar field) q . If v is in L^p ($1 < p < \infty$) in Ω , such a decomposition is well-studied. For example, a topological direct sum decomposition

$$(L^p(\Omega))^n = L_\sigma^p(\Omega) \oplus G^p(\Omega)$$

holds for various domains including $\Omega = \mathbf{R}^n$, a half space $\mathbf{R}_+^n$, a bounded smooth domain [8]; see e.g. G. P. Galdi [9]. Here, $L_\sigma^p(\Omega)$ denotes the L^p -closure of the space of all div-free vector fields compactly supported in Ω and $G^p(\Omega)$ denotes the totality of L^p gradient fields. It is impossible

to extend this Helmholtz decomposition to L^∞ even if $\Omega = \mathbf{R}^n$ since the projection $v \mapsto \nabla q$ is a composite of the Riesz operators which is not bounded in L^∞ . We have to replace L^∞ with a class of functions of bounded mean oscillation. However, if the vector field is of bounded mean oscillation (BMO for short), such a problem is only studied when Ω is a half space $\mathbf{R}_+^n$ [10], where the boundary is flat.

Our goal is to establish the Helmholtz decomposition of BMO vector fields in a smooth bounded domain in $\mathbf{R}^n$, which is a typical example of a domain with curved boundary. Although the space of BMO functions in $\mathbf{R}^n$ is well studied, the situation is less clear when one considers such a space in a domain, because there are several possible definitions. One should be careful about the behavior of a function near the boundary $\Gamma = \partial\Omega$. In this paper we study a space of BMO vector fields introduced in [11] and establish its Helmholtz decomposition when Ω is a bounded C^3 domain.

Let us recall the space $vBMO(\Omega)$ introduced in [11]. We first recall the BMO seminorm for $\mu \in (0, \infty]$. For a locally integrable function f , i.e., $f \in L_{\text{loc}}^1(\Omega)$ we define

$$[f]_{BMO^\mu(\Omega)} := \sup \left\{ \frac{1}{|B_r(x)|} \int_{B_r(x)} |f(y) - f_{B_r(x)}| dy \mid B_r(x) \subset \Omega, r < \mu \right\},$$

where f_B denotes the average over B , i.e.,

$$f_B := \frac{1}{|B|} \int_B f(y) dy$$

and $B_r(x)$ denotes the closed ball of radius r centered at x and $|B|$ denotes the Lebesgue measure of B . The space $BMO^\mu(\Omega)$ is defined as

$$BMO^\mu(\Omega) := \{f \in L_{\text{loc}}^1(\Omega) \mid [f]_{BMO^\mu} < \infty\}.$$

This space may not agree with the space of restrictions $r_\Omega f$ of $f \in BMO^\mu(\mathbf{R}^n)$. As in [1], [2], [3], [4] we introduce a seminorm controlling the boundary behavior. For $\nu \in (0, \infty]$, we set

$$[f]_{b^\nu} := \sup \left\{ r^{-n} \int_{\Omega \cap B_r(x)} |f(y)| dy \mid x \in \Gamma, 0 < r < \nu \right\}.$$

In these papers, the space

$$BMO_b^{\mu, \nu}(\Omega) := \{f \in BMO^\mu(\Omega) \mid [f]_{b^\nu} < \infty\}$$

is considered. Note that this space $BMO_b^{\infty, \infty}(\Omega)$ is identified with Miyachi's BMO introduced by [21] if Ω is a bounded Lipschitz domain or a Lipschitz half space as proved in [4]. However,

unfortunately, it turns out such a boundary control for whole components of vector fields is too strict to have the Helmholtz decomposition. We separate tangential and normal components. Let $d_\Gamma(x)$ denote the distance from the boundary Γ , i.e.,

$$d_\Gamma(x) := \inf \{|x - y|, y \in \Gamma\}.$$

For vector fields, we consider

$$vBMO^{\mu,\nu}(\Omega) := \{v \in (BMO^\mu(\Omega))^n \mid [\nabla d_\Gamma \cdot v]_{b^\nu} < \infty\},$$

where $\cdot$ denotes the standard inner product in $\mathbf{R}^n$. The quantity $(\nabla d_\Gamma \cdot v)\nabla d_\Gamma$ on Γ is the component of v normal to the boundary Γ . We set

$$[v]_{vBMO^{\mu,\nu}(\Omega)} := [v]_{BMO^\mu(\Omega)} + [\nabla d_\Gamma \cdot v]_{b^\nu}.$$

If Ω is the half space, this is not a norm but a seminorm. However, if it has a fully curved part in the sense of [11, Definition 7], then this becomes a norm [11, Lemma 8]. In particular, when Ω is a bounded C^2 domain, this is a norm. Roughly speaking, the boundary behavior of a vector field v is controlled for only normal part of v if $v \in vBMO^{\mu,\nu}(\Omega)$. For a bounded domain, this norm is equivalent no matter how μ and ν are taken; in other words, $vBMO^{\mu,\nu}(\Omega) = vBMO^{\infty,\infty}(\Omega)$. This is because $vBMO^{\mu,\nu}(\Omega) \subset L^1(\Omega)$ when Ω is bounded, which follows from the characterization of $vBMO^{\mu,\nu}(\Omega)$ in [11, Theorem 9]. We shall simply write $vBMO^{\mu,\nu}(\Omega)$ as $vBMO(\Omega)$. We are now in a position to state our main result.

Theorem 1. *Let Ω be a bounded C^3 domain in $\mathbf{R}^n$. Then the topological direct sum decomposition*

$$vBMO(\Omega) = vBMO_\sigma(\Omega) \oplus GvBMO(\Omega) \tag{2}$$

holds with

$$\begin{aligned} vBMO_\sigma(\Omega) &:= \{v \in vBMO(\Omega) \mid \operatorname{div} v = 0 \text{ in } \Omega, v \cdot \mathbf{n} = 0 \text{ on } \Gamma\}, \\ GvBMO(\Omega) &:= \{\nabla q \in vBMO(\Omega) \mid q \in L^1_{\text{loc}}(\Omega)\}, \end{aligned}$$

where $\mathbf{n}$ denotes the exterior unit normal vector field. In other words, for $v \in vBMO(\Omega)$, there is unique $v_0 \in vBMO_\sigma(\Omega)$ and $\nabla q \in GvBMO(\Omega)$ satisfying $v = v_0 + \nabla q$. Moreover, the mapping $v \mapsto v_0, v \mapsto \nabla q$ is bounded in $vBMO(\Omega)$.

As shown in [11], the norm trace $v \cdot \mathbf{n}$ is well defined as an element of $L^\infty(\Gamma)$ for $v \in vBMO(\Omega)$ with $\operatorname{div} v = 0$. So far, the Helmholtz decomposition BMO type space in a domain is only known

for $vBMO^{\infty,\infty}$ when Ω is the half space

$$\mathbf{R}_+^n = \{x = (x_1, \dots, x_n) \in \mathbf{R}^n \mid x_n > 0\}$$

as shown in [10], where the normal trace is taken in locally $H^{-1/2}$ sense.

Here is our strategy to show Theorem 1. For a vector field v , we construct a linear map $v \mapsto q_1$ such that q_1 satisfies

$$-\Delta q_1 = \operatorname{div} v \quad \text{in } \Omega,$$

where the divergence is taken in the sense of distribution. There are many ways to construct such a map because there is no boundary condition. A naive way is to extend v in a suitable way to a function $\bar{v}$ on $\mathbf{R}^n$ so that $v \mapsto \bar{v}$ is linear. We next consider the volume potential of $\operatorname{div} \bar{v}$, i.e.,

$$q_0(x) := \int_{\mathbf{R}^n} E(x-y) \operatorname{div} \bar{v}(y) dy = E * \operatorname{div} \bar{v},$$

where E is the fundamental solution of $-\Delta$ in $\mathbf{R}^n$, i.e.,

$$E(x) := \begin{cases} -\log |x|/2\pi & (n=2) \\ |x|^{2-n}/(n(n-2)\alpha(n)) & (n \geq 3), \end{cases}$$

where $\alpha(n)$ denotes the volume of the unit ball $B_1(0)$ of $\mathbf{R}^n$. By the famous *BMO-BMO* estimate due to Fefferman and Stein [7], we have

$$[\nabla q_0]_{BMO^\infty(\mathbf{R}^n)} \leq C_0[\bar{v}]_{BMO^\infty(\mathbf{R}^n)}$$

with $C_0 > 0$ independent of $\bar{v}$. However, it is difficult to control $[\nabla d_\Gamma \cdot \nabla q_0]_{b^\nu}$ so we construct another function q_1 instead of q_0 .

Although *BMO* space does not allow the standard cut-off procedure, our space is in L^1 , so we are able to decompose v into two parts $v = v_1 + v_2$ such that the support of v_2 is close to Γ while the support of v_1 is away from Γ ; see Proposition 5. For v_1 we just set

$$q_1^1 = E * \operatorname{div} v_1$$

by extending v_1 as zero outside its support. Then, the L^∞ bound for ∇q_1^1 is well controlled near Γ , which yields a bound for b^ν semi-norm. To estimate v_2 , we use a normal coordinate system near Γ and reduce the problem to the half space. Let d denotes the signed distance function where $d = d_\Gamma$ in Ω and $d = -d_\Gamma$ outside Ω . We extend v_2 to $\mathbf{R}^n$ so that the normal part $(\nabla d \cdot \bar{v}_2)\nabla d$ is odd and the tangential part $\bar{v}_2 - (\nabla d \cdot \bar{v}_2)\nabla d$ is even in the direction of ∇d with respect to Γ . In such type of coordinate system, the minus Laplacian can be transformed

as

$$L = A - B + \text{lower order terms}, \quad A = -\Delta_\eta, \quad B = \sum_{1 \leq i, j \leq n-1} \partial_{\eta_i} b_{ij} \partial_{\eta_j},$$

where η_n is the normal direction to the boundary so that $\{\eta_n > 0\}$ is the half space. By choosing a suitable coordinate system to represent Γ locally, we are able to arrange $b_{ij} = 0$ at one point of the boundary of the local coordinate system. We use a freezing coefficient method to construct volume potential q_1^2 and q_1^3 , which corresponds to the contribution from the tangential part $\bar{v}_2^{\text{tan}}$ and the normal part $\bar{v}_2^{\text{nor}}$ respectively. Since the leading term of $\text{div } \bar{v}_2^{\text{nor}}$ in normal coordinate consists of the differential of η_n only, if we extend the coefficient b_{ij} even in η_n , q_1^3 is constructed so that the leading term of $\nabla d \cdot \nabla q_1^3$ is odd in the direction of ∇d . On the other hand, as the leading term of $\text{div } \bar{v}_2^{\text{tan}}$ in normal coordinate consists of the differential of $\eta' = (\eta_1, \dots, \eta_{n-1})$ only, the even extension of b_{ij} in η_n gives rise to q_1^2 so that the leading term of $\nabla d \cdot \nabla q_1^2$ is also odd in the direction of ∇d . Disregarding lower order terms and localization procedure, we set q_1^2 and q_1^3 of the form

$$\begin{aligned} q_1^2 &= -L^{-1} \text{div } \bar{v}_2^{\text{tan}} = -A^{-1}(I - BA^{-1})^{-1} \text{div } \bar{v}_2^{\text{tan}}, \\ q_1^3 &= -L^{-1} \text{div } \bar{v}_2^{\text{nor}} = -A^{-1}(I - BA^{-1})^{-1} \text{div } \bar{v}_2^{\text{nor}}. \end{aligned}$$

One is able to arrange BA^{-1} small by taking a small neighborhood of a boundary point. Then $(I - BA^{-1})^{-1}$ is given as the Neumann series $\sum_{m=0}^{\infty} (BA^{-1})^m$. We are able to establish BMO - BMO estimate for ∇q_1^2 and ∇q_1^3 , i.e.

$$[\nabla q_1^2]_{BMO(\mathbf{R}^n)} \leq C'_0 [\bar{v}_2^{\text{tan}}]_{BMO(\mathbf{R}^n)}, \quad [\nabla q_1^3]_{BMO(\mathbf{R}^n)} \leq C'_0 [\bar{v}_2^{\text{nor}}]_{BMO(\mathbf{R}^n)}$$

with some constant C'_0 independent of $\bar{v}_2$. Since the leading term of $\nabla d \cdot (\nabla q_1^2 + \nabla q_1^3)$ is odd in the direction of ∇d with respect to Γ , the BMO bound implies b^ν bound. Note that $[\bar{v}_2^{\text{nor}}]_{BMO(\mathbf{R}^n)}$ is controlled by $[v_2]_{b^\nu}$ and $[v_2]_{BMO(\Omega)}$ since $\bar{v}_2^{\text{nor}}$ is odd in the direction of ∇d with respect to Γ . By the procedure sketched above, we are able to construct a suitable operator by setting $q_1 = q_1^1 + q_1^2 + q_1^3$.

Theorem 2 (Construction of a suitable volume potential). *Let Ω be a bounded C^3 domain in $\mathbf{R}^n$. Then, there exists a linear operator $v \mapsto q_1$ from $vBMO(\Omega)$ to $L^\infty(\Omega)$ such that*

$$-\Delta q_1 = \text{div } v \quad \text{in } \Omega$$

and that there exists a constant $C_1 = C_1(\Omega)$ satisfying

$$\|\nabla q_1\|_{vBMO(\Omega)} \leq C_1 \|v\|_{vBMO(\Omega)}.$$

In particular, the operator $v \mapsto \nabla q_1$ is a bounded linear operator in $vBMO(\Omega)$.

By this operator, we observe that $w = v - \nabla q_1$ is divergence free in Ω . Unfortunately, this w may not fulfill the trace condition $w \cdot \mathbf{n} = 0$ on the boundary Γ . We construct another potential q_2 by solving the Neumann problem

$$\begin{aligned} \Delta q_2 &= 0 \quad \text{in } \Omega \\ \frac{\partial q_2}{\partial n} &= w \cdot \mathbf{n} \quad \text{on } \Gamma. \end{aligned}$$

We then set $q = q_1 + q_2$. Since $\partial q_2 / \partial \mathbf{n} = \nabla q_2 \cdot \mathbf{n}$, $v_0 = v - \nabla q$ gives the Helmholtz decomposition (1). To complete the proof of Theorem 1, it suffices to prove that $\|\nabla q_2\|_{vBMO(\Omega)}$ is bounded by a constant multiply of $\|v\|_{vBMO(\Omega)}$.

Lemma 3 (Estimate of the normal trace). *Let Ω be a bounded $C^{2+\kappa}$ domain in $\mathbf{R}^n$ with $\kappa \in (0, 1)$. Then there is a constant $C_2 = C_2(\Omega)$ such that*

$$\|w \cdot \mathbf{n}\|_{L^\infty(\Gamma)} \leq C_2 \|w\|_{vBMO(\Omega)}$$

for all $w \in vBMO(\Omega)$ with $\operatorname{div} w = 0$.

This is a special case of the trace theorem established in [11]. We finally need the estimate for the Neumann problem.

Lemma 4 (Estimate for the Neumann problem). *Let Ω be a bounded C^2 domain. For $g \in L^\infty(\Gamma)$ satisfying $\int_\Gamma g d\mathcal{H}^{n-1} = 0$, there exists a unique (up to constant) solution u to the Neumann problem*

$$\begin{aligned} \Delta u &= 0 \quad \text{in } \Omega \\ \frac{\partial u}{\partial \mathbf{n}} &= g \quad \text{on } \Gamma \end{aligned} \tag{3}$$

such that the operator $g \mapsto u$ is linear and that there exists a constant $C_3 = C_3(\Omega)$ such that

$$\|\nabla u\|_{vBMO(\Omega)} \leq C_3 \|g\|_{L^\infty(\Gamma)}.$$

Combining these two lemmas, Theorem 2 yields

$$\begin{aligned} \|\nabla q_2\|_{vBMO(\Omega)} &\leq C_3 C_2 \|v - \nabla q_1\|_{vBMO(\Omega)} \\ &\leq C_3 C_2 (1 + C_1) \|v\|_{vBMO(\Omega)}. \end{aligned}$$

Setting $q = q_1 + q_2$ and $v_0 = v - \nabla q$, we now observe that the projections $v \mapsto v_0$, $v \mapsto \nabla q$

are bounded in $vBMO(\Omega)$, which yields (2) in Theorem 1.

To show Lemma 4 let $N(x, y)$ be the Neumann Green function. Then a solution of (3) is given by $\int_{\Gamma} N(x, y)g(y) d\mathcal{H}^{n-1}$. It is well-known (see e.g. [12, Appendix]) that leading part of N is $E(x - y)$. We have to estimate

$$\|\nabla E * (\delta_{\Gamma} \otimes g)\|_{vBMO^{\infty, \nu}(\Omega)}.$$

Here δ_{Γ} denotes the delta function supported on Γ , i.e.,

$$\delta_{\Gamma} : \psi \mapsto \int_{\Gamma} \psi d\mathcal{H}^{n-1}$$

for $\psi \in C_c^{\infty}(\mathbb{R}^n)$. We take a C^2 cutoff function $\theta \geq 0$ such that $\theta(\sigma) = 1$ for $\sigma \leq 1$, $\theta(\sigma) = 0$ for $\sigma \geq 2$. We take δ small so that 2δ is smaller than the reach of Γ . By this choice, $\theta_d = \theta(d/\delta)$ is C^2 in $\mathbf{R}^n$, where d denotes the signed distance function from Γ so that $\nabla d = -\mathbf{n}$ on Γ . For $g \in L^{\infty}(\Gamma)$, we extend g so that $\nabla d \cdot g = 0$ near the 2δ -neighborhood of Γ . Let g_e denotes this extension and set $g_{e,c} = \theta_d g_e$. A key observation is that

$$\begin{aligned} \delta_{\Gamma} \otimes g &= (\nabla 1_{\Omega} \cdot \nabla d)g_{e,c} = \operatorname{div}(g_{e,c} 1_{\Omega} \nabla d) - 1_{\Omega} \operatorname{div}(g_{e,c} \nabla d) \\ \operatorname{div}(g_{e,c} \nabla d) &= g_{e,c} \Delta d + \nabla d \cdot \nabla g_{e,c} = g_{e,c} \Delta d + \frac{\theta'(d/\delta)}{\delta} g_e, \end{aligned}$$

where 1_{Ω} is the characteristic function of Ω . The leading (singular) part of $\nabla E * (\delta_{\Gamma} \otimes g)$ is the term involving $\operatorname{div}(g_{e,c} 1_{\Omega} \nabla d)$. The famous L^{∞} - BMO estimate for the singular integral operator $\nabla E * \operatorname{div}$ yields

$$\|\nabla E * \operatorname{div}(g_{e,c} 1_{\Omega} \nabla d)\|_{BMO(\mathbf{R}^n)} \leq C \|g_{e,c} \nabla d\|_{L^{\infty}(\Omega)} \leq C' \|g\|_{L^{\infty}(\Gamma)}$$

with C and C' independent of g . All other terms can be estimated easily since the integral kernel is integrable. A direct calculation gives an L^{∞} estimate near Γ for $\nabla d \cdot \nabla E * (\delta_{\Gamma} \otimes g)$ which yields

$$[\nabla d \cdot \nabla E * (\delta_{\Gamma} \otimes g)]_{b^{\nu}} \leq C_4 \|g\|_{L^{\infty}(\Gamma)}$$

with C_4 independent of g , but it is impossible to estimate b^{ν} -seminorm of the tangential part. This is the main reason why we use $vBMO$ instead of BMO_b -type space where b^{ν} -boundedness of ALL components of vector fields is imposed; see the end of Section 4.2.

To extend our results to a more general domain it seems to be reasonable to consider $vBMO \cap L^2$. This is because $L^p \cap L^2$ ($p > 2$) admits the Helmholtz decomposition for arbitrary uniformly C^2 domains as proved in [5], [6].

Our approach in this paper is to derive the boundedness of the operator $v \mapsto \nabla q$ by a

potential-theoretic approach. In L^p setting there is a variational approach based on duality introduced by [23]; see also [5]. The key estimate is

$$\|\nabla q\|_{L^p(\Omega)} \leq C_5 \sup \left\{ \int_{\Omega} \nabla q \cdot \nabla \varphi \, dx \mid \|\nabla \varphi\|_{L^{p'}(\Omega)} \leq 1 \right\}$$

with C_5 independent of q , where $1/p + 1/p' = 1$, $1 < p < \infty$. Formally, this estimate yields the desired bound $\|\nabla q\|_{L^p(\Omega)} \leq C_5 \|v\|_{L^p(\Omega)}$ since $\int_{\Omega} \nabla q \cdot \nabla \varphi \, dx = \int_{\Omega} v \cdot \nabla \varphi \, dx$. At this moment, it is not clear that similar estimate holds if one replaces $L^p(\Omega)$ by $vBMO$ since the predual space of $vBMO$ is not clear.

For BMO_b type solution, it is known that the Stokes semigroup is analytic [1], [3]. However, it is nontrivial to extend to the space $vBMO$ since in the half space the Stokes operator with Dirichlet boundary condition does not generate a semigroup because $[u(t)]_{vBMO}$ for the solution $u(t)$ may be non-zero for $t > 0$ for initial data u_0 with $[u_0]_{vBMO} = 0$ so that $u_0^{\tan}$ may be a non-zero constant [1, Example 6.5].

This paper is organized as follows. In Section 2, to construct a volume potential of $\operatorname{div} v$, we localize the problem and reduce the problem to small neighborhoods of points on the boundary. In Section 3, we construct a leading part of the volume potential by a perturbation method called the freezing coefficient method. In these two sections, we complete the proof of Theorem 2. In Section 4, we prove Lemma 4 by estimating the single layer potential.

2 Construction of volume potentials

For $v \in vBMO(\Omega)$, we shall construct a suitable potential q_1 so that $v \mapsto \nabla q_1$ is a bounded linear operator in $vBMO$ as stated in Theorem 2. In this section, as a preliminary, we reduce the problem to the case that the support of v is contained in a small neighborhood of a point of the boundary and it consists of only normal part.

2.1 Localization procedure

Let Ω be a uniformly C^k domain in $\mathbf{R}^n$ ($k \geq 1$). In other words, there exists $r_*, \delta_* > 0$ such that for each $z_0 \in \Gamma$, up to translation and rotation, there exists a function h_{z_0} which is C^k in a closed ball $B_{r_*}(0')$ of radius r_* centered at the origin $0'$ of $\mathbf{R}^{n-1}$ satisfying following properties:

- (i) $K_{\Gamma} := \sup_{B_{r_*}(0')} |(\nabla')^s h_{z_0}| < \infty$ for $s = 0, 1, 2, \dots, k$, where ∇' denotes the gradient in $x' \in \mathbf{R}^{n-1}$; $\nabla' h(0') = 0$, $h(0') = 0$,

(ii) $\Omega \cap U_{r_*, \delta_*, h_{z_0}}(z_0) = \{(x', x_n) \in \mathbf{R}^n \mid h_{z_0}(x') < x_n < h_{z_0}(x') + \delta_*, |x'| < r_*\}$ for

$$U_{r_*, \delta_*, h_{z_0}}(z_0) := \{(x', x_n) \in \mathbf{R}^n \mid h_{z_0}(x') - \delta_* < x_n < h_{z_0}(x') + \delta_*, |x'| < r_*\},$$

(iii) $\Gamma \cap U_{r_*, \delta_*, h_{z_0}}(z_0) = \{(x', x_n) \in \mathbf{R}^n \mid x_n = h_{z_0}(x'), |x'| < r_*\}$.

A bounded C^k domain is, of course, a uniformly C^k domain.

Let d denote the signed distance function from Γ which is defined by

$$d(x) = \begin{cases} \inf_{y \in \Gamma} |x - y| & \text{for } x \in \Omega, \\ -\inf_{y \in \Gamma} |x - y| & \text{for } x \notin \Omega \end{cases} \quad (4)$$

so that $d(x) = d_\Gamma(x)$ for $x \in \Omega$. If Ω is a bounded C^2 domain, then there is $R_* > 0$ such that if $|d(x)| < R_*$, there is unique point πx such that $|x - \pi x| = |d(x)|$. The supremum of such R_* is called the reach of Ω and Ω^c . Moreover, d is C^2 in the R_* -neighborhood of Γ , i.e., $d \in C^2(\Gamma_{R_*}^{\mathbf{R}^n})$ with

$$\Gamma_{R_*}^{\mathbf{R}^n} := \{x \in \mathbf{R}^n \mid |d(x)| < R_*\};$$

see [13, Chap. 14, Appendix], [19, §4.4]. Note that R_* satisfies

$$R_* = \min(R_*^\Omega, R_*^{\Omega^c}),$$

where R_*^Ω is the reach of Γ in Ω while $R_*^{\Omega^c}$ is the reach of Γ in the complement Ω^c of Ω . Let $K_\Gamma^* := \max\{K_\Gamma, 1\}$. There exists $0 < \rho_0 < \min(r_*, \delta_*, \frac{R_*}{2}, \frac{1}{2nK_\Gamma^*})$ such that

$$U_\rho(z_0) := \{x \in \mathbf{R}^n \mid (\pi x)' \in \text{int } B_\rho(0'), |d(x)| < \rho\}$$

is contained in the coordinate chart $U_{r_*, \delta_*, h_{z_0}}(z_0)$ for any $\rho \leq \rho_0$.

We always take $\rho < \rho_0$. Since Ω is bounded and

$$\bigcup_{z \in \Gamma} U_\rho(z)$$

covers the compact set $K = \text{cl}\left(\Gamma_{\rho/2}^{\mathbf{R}^n}\right)$, there exists a finite subcover $\{U_\rho(z_j)\}_{j=1}^m$ of K , where the number m depends on ρ . For $\sigma > 0$, we denote that

$$\Omega^\sigma = \Omega \setminus \Gamma_\sigma^{\mathbf{R}^n}, \quad U_{\sigma,j} := U_\sigma(z_j).$$

Observe that

$$\overline{\Omega} \subset \bigcup_{j=1}^m U_{\rho,j} \cup \Omega^{\rho/2}.$$

Let $\{\varphi_j\}_{j=0}^m$ be a partition of the unity associated with $\{U_{\rho,j}\} \cup \{\Omega^{\rho/2}\}$ in the sense that

$$\begin{aligned} \varphi_j &\in C_c^\infty(U_{\rho,j} \cap \overline{\Omega}), \quad 0 \leq \varphi_j \leq 1 \quad \text{for } j = 1, \dots, m, \\ \varphi_0 &\in C_c^\infty(\Omega^{\rho/2}), \quad 0 \leq \varphi_0 \leq 1, \quad \varphi_0 = 1 \quad \text{in } \Omega^\rho \end{aligned}$$

and

$$\sum_{j=0}^m \varphi_j = 1 \quad \text{in } \overline{\Omega}.$$

Here $C_c^\infty(W)$ denotes the space of all smooth function in W whose support is compact in W .

Throughout this paper, unless otherwise specified, the symbol C in an inequality represents a positive constant independent of quantities that appeared in the inequality. For a fixed $\rho > 0$, C_ρ represents a constant depending only on ρ . C_n represents a constant depending only on n and $C_{\Omega,n}$ represents a constant depending only on Ω and n .

2.2 Cut-off and extension

In general, multiplication by a smooth function to BMO is not bounded in BMO . Fortunately, our space is closed by multiplication.

Proposition 5 (Multiplication). *Let Ω be a bounded C^2 domain in $\mathbf{R}^n$. Let $\varphi \in C^\gamma(\Omega)$, $\gamma \in (0, 1)$. For each $v \in vBMO(\Omega)$, the function $\varphi v \in vBMO(\Omega)$ satisfies*

$$\|\varphi v\|_{vBMO(\Omega)} \leq C \|\varphi\|_{C^\gamma(\Omega)} \|v\|_{vBMO(\Omega)}$$

with C independent of φ and v .

Proof. Since

$$[\nabla d \cdot \varphi v]_{b^\nu} \leq \|\varphi\|_{L^\infty(\Omega)} [\nabla d \cdot v]_{b^\nu},$$

it suffices to establish the estimate

$$[\varphi v]_{BMO(\Omega)} \leq c_0 \|\varphi\|_{C^\gamma(\Omega)} \|v\|_{vBMO(\Omega)} \tag{5}$$

with c_0 independent of φ and v . Since a bounded Lipschitz domain is a uniform domain, we are able to apply [11, Theorem 13] to get

$$[\varphi v]_{BMO(\Omega)} \leq c_1 \|\varphi\|_{C^\gamma(\Omega)} ([v]_{BMO(\Omega)} + \|v\|_{L^1(\Omega)}).$$

This is based on the product estimate of a Hölder function and a function in $bmo(\mathbf{R}^n) := BMO(\mathbf{R}^n) \cap L_{ul}^1(\mathbf{R}^n)$ where

$$L_{ul}^1(\mathbf{R}^n) := \left\{ f \in L_{loc}^1(\mathbf{R}^n) \mid \|f\|_{L_{ul}^1(\mathbf{R}^n)} := \sup_{x \in \mathbf{R}^n} \int_{B_1(x)} |f(y)| dy < \infty \right\}.$$

The space $bmo(\mathbf{R}^n)$ is equipped with the norm

$$\|f\|_{bmo(\mathbf{R}^n)} := [f]_{BMO(\mathbf{R}^n)} + \|f\|_{L_{ul}^1(\mathbf{R}^n)}$$

for $f \in bmo(\mathbf{R}^n)$. The product estimate for bmo follows from a similar result for a local Hardy space $h^1 = F_{1,2}^0$ [22, Remark 4.4] and duality $bmo = (h^1)'$ [22, Theorem 3.26]. To handle a function in Ω , we need an extension to conclude [11, Theorem 13]. Fortunately, by the characterization of $vBMO$ for a bounded C^2 domain [11, Theorem 9],

$$\|v\|_{L^1(\Omega)} \leq c_2 \|v\|_{vBMO(\Omega)}.$$

Here c_j denotes a constant independent of v and φ for $j = 1, 2$. Combining these two estimates, we obtain (5) with $c_0 = c_1(1 + c_2)$. This yields Proposition 5. $\square$

For a bounded C^3 domain, we next consider an extension based on the normal coordinate in $U_\rho(z_0)$ for $\rho \leq \rho_0$ of the form

$$\begin{cases} x' &= \eta' + \eta_n \nabla' d(\eta', h_{z_0}(\eta')); \\ x_n &= h_{z_0}(\eta') + \eta_n \partial_{x_n} d(\eta', h_{z_0}(\eta')). \end{cases} \quad (6)$$

Let $V_\sigma := B_\sigma(0') \times (-\sigma, \sigma)$ for $\sigma \in (0, \rho_0)$. We shall write this coordinate change by $x = \psi(\eta)$ with $\psi \in C^2(V_{\rho_0})$ and

$$x = \pi x - d(x) \mathbf{n}(\pi x), \quad \mathbf{n}(\pi x) = -\nabla d(\pi x).$$

We consider the projection to the direction to ∇d . For $x \in \Gamma_{\rho_0}^{\mathbf{R}^n}$, we set

$$P(x) = \nabla d(\pi x) \otimes \nabla d(\pi x) = \mathbf{n}(\pi x) \otimes \mathbf{n}(\pi x).$$

For later convenience, we set $Q(x) = I - P(x)$ which is the tangential projection for $x \in \Gamma_{\rho_0}^{\mathbf{R}^n}$. For a function f in $\Gamma_\rho^{\mathbf{R}^n} \cap \overline{\Omega}$, let f_{even} (resp. f_{odd}) denote its even (odd) extension to $\Gamma_\rho^{\mathbf{R}^n}$ defined

by

$$\begin{aligned} f_{\text{even}}(\pi x + d(x)\mathbf{n}(\pi x)) &= f(\pi x - d(x)\mathbf{n}(\pi x)) & \text{for } x \in \Gamma_\rho^{\mathbf{R}^n} \setminus \overline{\Omega}, \\ f_{\text{odd}}(\pi x + d(x)\mathbf{n}(\pi x)) &= -f(\pi x - d(x)\mathbf{n}(\pi x)) & \text{for } x \in \Gamma_\rho^{\mathbf{R}^n} \setminus \overline{\Omega}. \end{aligned}$$

We denote r_W to be the restriction in W for any subset $W \subset \mathbf{R}^n$. Let f be a function (or a vector field) defined in V_σ for some $\sigma \in (0, \infty]$. We set $E_{\text{even}}f$ to be the even extension of f in $V_\sigma \cap \mathbf{R}_+^n$ to V_σ with respect to the n -th variable, i.e.,

$$E_{\text{even}}f(\eta', -\eta_n) = f(\eta', \eta_n)$$

for any $(\eta', \eta_n) \in V_\sigma \cap \mathbf{R}_+^n$.

For $v \in vBMO(\Omega)$ with $\text{supp } v \subset U_\rho(z_0) \cap \overline{\Omega}$, let $\bar{v}$ be its extension of the form

$$\bar{v}(x) := (Pv_{\text{odd}})(x) + (Qv_{\text{even}})(x) \quad (7)$$

for $x \in U_\rho(z_0)$. Notice that $\text{supp } \bar{v} \subset U_\rho(z_0)$, $\bar{v}$ is indeed defined in $\mathbf{R}^n$ with $\bar{v}(x) = 0$ for any $x \in U_\rho(z_0)^c$. Define

$$L_* := \sup_{z_0 \in \Gamma, \rho \leq \rho_0} \max \{ \|\nabla \psi\|_{L^\infty(V_\rho)} + \|\nabla \psi^{-1}\|_{L^\infty(U_\rho(z_0))}, 1 \}.$$

Since the boundary Γ is uniformly C^3 , L_* is finite that depends on Ω only. We set $\rho_{0,*} = \rho_0/12L_*$.

Proposition 6. *Let $\Omega \subset \mathbf{R}^n$ be a bounded C^2 domain, $z_0 \in \Gamma$ and $\rho \in (0, \rho_{0,*})$. There exists a constant C_ρ , which depends on ρ only, such that*

$$\begin{aligned} [\bar{v}]_{BMO(\mathbf{R}^n)} &\leq C_\rho \|v\|_{vBMO(\Omega)}, \\ [\nabla d \cdot \bar{v}]_{b^\nu(\Gamma)} &\leq C_\rho \|v\|_{vBMO(\Omega)} \end{aligned}$$

for all $v \in vBMO(\Omega)$ with $\text{supp } v \subset U_\rho(z_0) \cap \overline{\Omega}$ and $\nu > 0$.

In the normal coordinate, $P\bar{v} = Pv_{\text{odd}}$ is odd in η_n and $Q\bar{v} = Qv_{\text{even}}$ is even in η_n . The key idea of proving this proposition is to reduce the problem to the case where the boundary is locally flat by invoking the normal coordinate.

Proof. Since $vBMO(\Omega) \subset L^1(\Omega)$, see e.g. [11, Theorem 9], by considering the normal coordinate change $y = \psi(\eta)$ in $U_\rho(z_0)$ we can deduce that $v_{\text{even}}, v_{\text{odd}} \in L^1(\mathbf{R}^n)$ satisfying

$$\|v_{\text{even}}\|_{L^1(\mathbf{R}^n)} = \|v_{\text{odd}}\|_{L^1(\mathbf{R}^n)} \leq 2L_*^2 \|v\|_{L^1(\Omega)}.$$

Hence $\bar{v} \in L^1(\mathbf{R}^n)$ satisfies the estimate $\|\bar{v}\|_{L^1(\mathbf{R}^n)} \leq C_{\Omega,n}\|v\|_{L^1(\Omega)}$. Since Ω is a uniform domain, by [17, Theorem 1] there exists $v_J \in BMO(\mathbf{R}^n)$ with $r_\Omega v_J = v$ and

$$[v_J]_{BMO(\mathbf{R}^n)} \leq C_{\Omega,n}[v]_{BMO^\infty(\Omega)}.$$

Suppose that $B_r(\zeta) \subset V_{4\rho L_*}^+ := V_{4\rho L_*} \cap \mathbf{R}_+^n$. The mean value theorem implies that $\psi(B_r(\zeta)) \subset B_{L_*r}(x)$ with $x = \psi(\zeta)$. By change of variables $y = \psi(\eta)$ in $U_{4\rho L_*}(z_0)$, we see that

$$\begin{aligned} \frac{1}{|B_r(\zeta)|} \int_{B_r(\zeta)} |v \circ \psi(\eta) - c| d\eta &\leq L_* \cdot \frac{1}{|B_r(\zeta)|} \int_{\psi(B_r(\zeta))} |v(y) - c| dy \\ &\leq C_n L_*^{n+1} \cdot \frac{1}{|B_{L_*r}(x)|} \int_{B_{L_*r}(x)} |v_J(y) - c| dy \end{aligned}$$

for any constant vector $c \in \mathbf{R}^n$. By considering an equivalent definition of the BMO -seminorm, see e.g. [14, Proposition 3.1.2], we deduce that

$$[v \circ \psi]_{BMO^\infty(V_{4\rho L_*}^+)} \leq C_{\Omega,n}[v]_{BMO^\infty(\Omega)}.$$

By recalling the results concerning the even extension of BMO functions in the half space, see [10, Lemma 3.2] and [10, Lemma 3.4], we can deduce that

$$[v_{\text{even}} \circ \psi]_{BMO^\infty(V_{4\rho L_*})} \leq C_{\Omega,n}[v]_{BMO^\infty(\Omega)}. \quad (8)$$

Next, we shall estimate the BMO -seminorm of v_{even} . Let $B_r(x)$ be a ball with radius $r \leq \frac{\rho}{2}$. If either $B_r(x) \cap U_\rho(z_0) = \emptyset$ or $B_r(x) \subset \Omega$, there is nothing to prove. It is sufficient to consider $B_r(x)$ that intersects both $U_\rho(z_0)$ and Ω^c . In this case we can find $x_0 \in B_r(x) \cap U_\rho(z_0)$. Since $B_r(x) \subset B_{2r}(x_0) \subset B_{4\rho}(z_0) \subset U_{8\rho}(z_0)$, by considering change of variables $y = \psi(\eta)$ in $U_{8\rho}(x_0)$, we have that

$$\frac{1}{|B_r(x)|} \int_{B_r(x)} |v_{\text{even}}(y) - c| dy \leq \frac{L_*}{|B_r(x)|} \int_{\psi^{-1}(B_r(x))} |v_{\text{even}} \circ \psi(\eta) - c| d\eta.$$

For any $y \in B_r(x)$, we have that $|y - z_0| < 4\rho$. Hence $\psi^{-1}(B_r(x)) \subset B_{L_*r}(\zeta) \subset B_{4\rho L_*}(0) \subset V_{4\rho L_*}$. By (8), we deduce that

$$\frac{1}{|B_r(x)|} \int_{B_r(x)} |v_{\text{even}}(y) - (v_{\text{even}})_{B_r(x)}| dy \leq C_{\Omega,n}[v]_{BMO^\infty(\Omega)}.$$

Thus, we obtain that

$$[v_{\text{even}}]_{BMO^{\frac{\rho}{2}}(\mathbf{R}^n)} \leq C_{\Omega,n}[v]_{BMO^\infty(\Omega)}.$$

For a ball B with radius $r(B) > \frac{\rho}{2}$, a simple triangle inequality implies that

$$\frac{1}{|B|} \int_B |v_{\text{even}}(y) - (v_{\text{even}})_B| dy \leq \frac{2}{|B|} \int_B |v_{\text{even}}(y)| dy \leq \frac{C_n}{\rho^n} \|v_{\text{even}}\|_{L^1(\mathbf{R}^n)}.$$

Therefore, we obtain the BMO estimate for v_{even} , i.e.,

$$[v_{\text{even}}]_{BMO(\mathbf{R}^n)} \leq \frac{C_{\Omega,n}}{\rho^n} \|v\|_{vBMO(\Omega)}.$$

We shall then give the BMO estimate for Pv_{odd} . Since $\nabla d \in C^1(\Gamma_{\rho_0}^{\mathbf{R}^n})$, there exists $D_e \in C^1(\mathbf{R}^n)$ such that $\|D_e\|_{C^1(\mathbf{R}^n)} \leq \|\nabla d\|_{C^1(\Gamma_{\rho_0}^{\mathbf{R}^n})}$ and $r_{\Gamma_{\rho_0}^{\mathbf{R}^n}} D_e = \nabla d$, see the proof of [11, Theorem 13]. By the multiplication rule for bmo functions, we have that $(Pv)_E := (D_e \cdot v_{\text{even}})D_e \in bmo(\mathbf{R}^n)$, see also [11, Theorem 13]. Consider the normal coordinate change in $U_{4\rho L_*}(z_0)$. Since $(Pv)_E = Pv$ in $U_{4\rho L_*}(z_0) \cap \Omega$, same argument in the second paragraph implies that

$$[Pv \circ \psi]_{BMO^\infty(V_{4\rho L_*}^+)} \leq C_{\Omega,n} \|(Pv)_E\|_{bmo(\mathbf{R}^n)} \leq \frac{C_{\Omega,n}}{\rho^n} \|v\|_{vBMO(\Omega)}.$$

Let $\zeta \in V_{12\rho L_*} = \psi^{-1}(U_{12\rho L_*}(z_0))$ with $\zeta_n = 0$. Let $B_r(\zeta) \subset V_{12\rho L_*}$ and $x = \psi(\zeta)$. Since $F(B_r(\zeta) \cap V_{12\rho L_*}^+) \subset B_{L_*r}(x) \cap \Omega$, by considering change of variables $y = \psi(\eta)$ in $U_{12\rho L_*}(z_0)$, we can deduce that

$$\frac{1}{|B_r(\zeta)|} \int_{B_r(\zeta) \cap V_{12\rho L_*}^+} |Pv_{\text{odd}} \circ \psi(\eta)| d\eta \leq L_*^{n+1} [\nabla d \cdot v]_{b^\nu}. \quad (9)$$

Recall the results concerning the odd extension of BMO functions in the half space, see [10, Lemma 3.1], we have the estimate

$$[Pv_{\text{odd}} \circ \psi]_{BMO^\infty(V_{4\rho L_*})} \leq \frac{C_{\Omega,n}}{\rho^n} \|v\|_{vBMO(\Omega)}. \quad (10)$$

By considering (10) and the fact that $Pv_{\text{odd}} = (Pv)_E$ in Ω , same argument in the third paragraph implies the BMO estimate for Pv_{odd} , i.e.,

$$[Pv_{\text{odd}}]_{BMO(\mathbf{R}^n)} \leq \frac{C_{\Omega,n}}{\rho^n} \|v\|_{vBMO(\Omega)}.$$

Combining the BMO estimates for v_{even} and Pv_{odd} , we have that

$$[\bar{v}]_{BMO(\mathbf{R}^n)} \leq \frac{C_{\Omega,n}}{\rho^n} \|v\|_{vBMO(\Omega)}.$$

Notice that $\nabla d \cdot \bar{v} = v_{\text{odd}} \cdot \nabla d$ in $\mathbf{R}^n$. Let $x \in \Gamma$ and $r \leq \frac{\rho}{L_*}$. If $B_r(x) \cap U_\rho(z_0) = \emptyset$, then $v_{\text{odd}} = 0$ in $B_r(x)$. Suppose that $B_r(x) \cap U_\rho(z_0) \neq \emptyset$. Then we can find $x_0 \in B_r(x) \cap U_\rho(z_0) \cap \Gamma$. Let

$\zeta_0 = \psi^{-1}(x_0)$, we have that $\psi^{-1}(B_r(x)) \subset B_{2L_*r}(\zeta_0) \subset V_{12\rho L_*}$. Hence,

$$\begin{aligned} r^{-n} \int_{B_r(x)} |v_{\text{odd}} \cdot \nabla d| dy &\leq \frac{2L_*}{r^n} \int_{B_{2L_*r}(\zeta_0) \cap V_{12\rho L_*}^+} |(v \cdot \nabla d) \circ \psi| d\eta \\ &\leq \frac{2L_*^2}{r^n} \int_{B_{2L_*^2r}(x_0) \cap \Omega} |\nabla d \cdot v| dy \leq C_{\Omega,n} [\nabla d \cdot v]_{b^\nu}. \end{aligned}$$

For $r > \frac{\rho}{L_*}$, we simply have that

$$r^{-n} \int_{B_r(x)} |v_{\text{odd}} \cdot \nabla d| dy \leq \frac{C_{\Omega,n}}{\rho^n} \|v_{\text{odd}}\|_{L^1(\mathbf{R}^n)} \leq \frac{C_{\Omega,n}}{\rho^n} \|v\|_{vBMO(\Omega)}.$$

□

2.3 Volume potentials

To construct mapping $v \mapsto q_1$ in Theorem 2, for some ρ_* to be determined later in the next section, we localize v by using the partition of the unity $\{\varphi_j\}_{j=0}^m$ associated with the covering

$$\{U_{\rho,j}\}_{j=1}^m \cup \Omega^{\rho/2}$$

as in Section 2.1, where ρ is always assumed to satisfy $\rho \leq \rho_*/2$. Here and hereafter we always assumed that Ω is a bounded C^3 domain in $\mathbf{R}^n$.

Proposition 7. *There exists a constant C_ρ , which depends on ρ only, such that*

$$\begin{aligned} [\nabla q_1^1]_{BMO^\infty(\mathbf{R}^n)} &\leq C_\rho \|v\|_{vBMO(\Omega)}, \\ \|\nabla q_1^1(x)\|_{L^\infty(\Gamma_{\rho/4}^{\mathbf{R}^n})} &\leq C_\rho \|v\|_{vBMO(\Omega)} \end{aligned}$$

for $q_1^1 = E * \text{div}(\varphi_0 v)$ and $v \in vBMO(\Omega)$. In particular,

$$[\nabla q_1^1]_{b^\nu(\Gamma)} \leq C_\rho \|v\|_{vBMO(\Omega)}$$

for $\nu < \rho/4$.

Proof. By the BMO - BMO estimate [7], we have the estimate

$$[\nabla q_1^1]_{BMO(\mathbf{R}^n)} \leq C[\varphi_0 v]_{BMO(\mathbf{R}^n)}.$$

Consider $x \in \Gamma_{\rho/4}^{\mathbf{R}^n}$. Since ∇q_1^1 is harmonic in $\Gamma_{\rho/2}^{\mathbf{R}^n}$ and $B_{\frac{\rho}{4}}(x) \subset \Gamma_{\rho/2}^{\mathbf{R}^n}$, the mean value property

for harmonic functions implies that

$$\nabla q_1^1(x) = \frac{C_n}{\rho^n} \int_{B_{\rho/4}(x)} \nabla q_1^1(y) dy.$$

By Hölder's inequality, we can estimate $|\nabla q_1^1(x)|$ by $\frac{C_n}{\rho^{n/2}} \|\nabla q_1^1\|_{L^2(\mathbf{R}^n)}$. Since the convolution with $\nabla^2 E$ is bounded in L^p for any $1 < p < \infty$, see e.g. [14, Theorem 5.2.7 and Theorem 5.2.10], an interpolation inequality (cf. [4, Lemma 5]) implies that

$$\|\nabla q_1^1\|_{L^2(\mathbf{R}^n)} \leq C \|\varphi_0 v\|_{L^2(\mathbf{R}^n)} \leq C \|\varphi_0 v\|_{L^1(\mathbf{R}^n)}^{\frac{1}{2}} [\varphi_0 v]_{BMO(\mathbf{R}^n)}^{\frac{1}{2}}.$$

View $\varphi_0 v$ as the extension of $\varphi_0 v$ from Ω to $\mathbf{R}^n$. By the extension theorem for *bmo* functions [11, Theorem 12], we estimate $[\varphi_0 v]_{BMO(\mathbf{R}^n)}$ by $C_\rho [\varphi_0 v]_{BMO^\infty(\Omega)}$. Since $vBMO(\Omega) \subset L^1(\Omega)$, see [11, Theorem 9], Proposition 5 implies that

$$|\nabla q_1^1(x)| \leq C_\rho \|v\|_{vBMO(\Omega)}$$

for any $x \in \Gamma_{\rho/4}^{\mathbf{R}^n}$. □

We next set $v_1 := \varphi_0 v$ and $v_2 := 1 - v_1$. For each $\varphi_j v_2$ ($j = 1, \dots, m$), we extend as in Proposition 6 to get $\overline{\varphi_j v_2}$ and set

$$\overline{v_2} := \sum_{j=1}^m \overline{\varphi_j v_2}.$$

Indeed, this extension is independent of the choice φ_j 's but we do not use this fact. We next set

$$\overline{v_2}^{\text{tan}} := Q \overline{v_2} = \sum_{j=1}^m Q (\varphi_j v_2)_{\text{even}}.$$

For $1 \leq j \leq m$, $\varphi_j \in C_c^\infty(U_{\rho,j} \cap \overline{\Omega})$ implies that the even extension of φ_j in $U_{\rho,j}$ with respect to Γ is Hölder continuous in the sense that $(\varphi_j)_{\text{even}} \in C^{0,1}(U_{\rho,j})$. Moreover, we have that $(\varphi_j)_{\text{even}} \in C^{0,1}(\mathbf{R}^n)$ satisfies

$$\|(\varphi_j)_{\text{even}}\|_{C^{0,1}(\mathbf{R}^n)} \leq C_\rho \|(\varphi_j)_{\text{even}}\|_{C^{0,1}(U_{\rho,j})}.$$

For simplicity of notations, we denote $Q (\varphi_j v_2)_{\text{even}}$ by w_j^{tan} for every $1 \leq j \leq m$. Now, we are ready to construct the suitable potential corresponding to $\overline{v_2}^{\text{tan}}$.

Proposition 8. *There exists $\rho_* > 0$ such that if $\rho < \rho_*/2$, then for every $1 \leq j \leq m$, there*

exists a linear operator $v \mapsto p_j^{\text{tan}}$ from $vBMO(\Omega)$ to $L^\infty(\mathbf{R}^n)$ such that

$$-\Delta p_j^{\text{tan}} = \text{div } w_j^{\text{tan}} \text{ in } U_{2\rho,j} \cap \Omega$$

and that there exists a constant C_ρ , independent of v , such that

$$\begin{aligned} [\nabla p_j^{\text{tan}}]_{BMO(\mathbf{R}^n)} &\leq C_\rho \|v\|_{vBMO(\Omega)}, \\ \sup_{x \in \Gamma, r < \rho} \frac{1}{r^n} \int_{B_r(x)} |\nabla d \cdot \nabla p_j^{\text{tan}}| dy &\leq C_\rho \|v\|_{vBMO(\Omega)}. \end{aligned}$$

Having the estimate for the volume potential near the boundary regarding its tangential component, we are left to handle the contribution from $\bar{v}_2^{\text{nor}} := \bar{v}_2 - \bar{v}_2^{\text{tan}}$. We recall its decomposition

$$\bar{v}_2^{\text{nor}} = \sum_{j=1}^m P(\varphi_j v_2)_{\text{odd}}.$$

For simplicity of notations, we denote $P(\varphi_j v_2)_{\text{odd}}$ by w_j^{nor} for every $1 \leq j \leq m$. With a similar idea of proof, we can establish the suitable potential corresponding to $\bar{v}_2^{\text{nor}}$.

Proposition 9. *There exists $\rho_* > 0$ such that if $\rho < \rho_*/2$, then for every $1 \leq j \leq m$, there exists a linear operator $v \mapsto p_j^{\text{nor}}$ from $vBMO(\Omega)$ to $L^\infty(\mathbf{R}^n)$ such that*

$$-\Delta p_j^{\text{nor}} = \text{div } w_j^{\text{nor}} \text{ in } U_{2\rho,j} \cap \Omega$$

and that there exists a constant C_ρ , independent of v , such that

$$\begin{aligned} [\nabla p_j^{\text{nor}}]_{BMO(\mathbf{R}^n)} &\leq C_\rho \|v\|_{vBMO(\Omega)}, \\ \sup_{x \in \Gamma, r < \rho} \frac{1}{r^n} \int_{B_r(x)} |\nabla d \cdot \nabla p_j^{\text{nor}}| dy &\leq C_\rho \|v\|_{vBMO(\Omega)}. \end{aligned}$$

Once these two propositions are proved, we are able to prove Theorem 2.

Theorem 2 admitting Proposition 8 and 9. Fix $1 \leq j \leq m$. Let us first consider the contribution from the tangential part. We take a cut-off function $\theta_j \in C_c^\infty(U_{2\rho,j})$ such that $\theta_j = 1$ on $U_{\rho,j}$ and $0 \leq \theta_j \leq 1$. We next set

$$q_{1,j}^{\text{tan}} := \theta_j p_j^{\text{tan}} + E * (p_j^{\text{tan}} \Delta \theta_j + 2 \nabla \theta_j \cdot \nabla p_j^{\text{tan}}).$$

By definition, Proposition 8 says that

$$\begin{aligned} -\Delta q_{1,j}^{\tan} &= -\Delta(\theta_j p_j^{\tan}) + p_j^{\tan} \Delta \theta_j + 2\nabla \theta_j \cdot \nabla p_j^{\tan} \\ &= \theta_j \operatorname{div} w_j^{\tan} = \operatorname{div} w_j^{\tan} \end{aligned}$$

in Ω as $\operatorname{supp} w_j^{\tan} \subset U_{\rho,j}$. By interpolation as in the proof of Proposition 8, we observe that $\|p_j^{\tan}\|_{L^\infty(\mathbf{R}^n)}$, $\|\nabla p_j^{\tan}\|_{L^p(\mathbf{R}^n)}$ are controlled by $\|v\|_{BMO(\Omega)}$. Since ∇E is in $L^{p'}(B_R)$ for $p' < n/(n-1)$ where $R = \operatorname{diam} \Omega + 4\rho$, it follows that

$$\sup_{\mathbf{R}^n} |\nabla E * (p_j^{\tan} \Delta \theta_j + 2\nabla \theta_j \cdot \nabla p_j^{\tan})| \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

Thus, by Proposition 8, we conclude that the restriction of $q_{1,j}^{\tan}$ on Ω , which is still denoted by $q_{1,j}^{\tan}$, fulfills

$$\|\nabla q_{1,j}^{\tan}\|_{vBMO(\Omega)} \leq C_\rho \|v\|_{vBMO(\Omega)}. \quad (11)$$

By Proposition 9, a similar argument yields an estimate of type (11) for

$$q_{1,j}^{\text{nor}} := \theta_j p_j^{\text{nor}} + E * (p_j^{\text{nor}} \Delta \theta_j + 2\nabla \theta_j \cdot \nabla p_j^{\text{nor}}).$$

Set

$$q_1^2 = \sum_{j=1}^m q_{1,j}^{\tan}, \quad q_1^3 = \sum_{j=1}^m q_{1,j}^{\text{nor}}, \quad q_1 = q_1^1 + q_1^2 + q_1^3.$$

Observe that q_1^2 and q_1^3 satisfy the desired estimates in Theorem 2. Moreover, by construction we have that

$$\begin{aligned} -\Delta q_1 &= -\Delta q_1^1 - \Delta q_1^2 - \Delta q_1^3 \\ &= \operatorname{div} v_1 + \sum_{j=1}^m \operatorname{div} w_j^{\tan} + \sum_{j=1}^m \operatorname{div} w_j^{\text{nor}} \\ &= \operatorname{div}(v_1 + v_2) = \operatorname{div} v \end{aligned}$$

in Ω . □

3 Volume potentials based on normal coordinates

Our goal in this section is to prove Proposition 8 and Proposition 9. We write the Laplace operator by a normal coordinate system and construct a volume potential keeping the parity of functions with respect to the boundary. For this purpose, we adjust a perturbation method called a freezing coefficient method which is often used to construct a fundamental solution to

an operator with variable coefficients.

3.1 A perturbation method keeping parity

We consider an elliptic operator of the form

$$L_0 = A - B, \quad A = -\Delta_\eta, \quad B = \sum_{1 \leq i, j \leq n-1} \partial_{\eta_i} b_{ij}(\eta) \partial_{\eta_j}$$

in a cylinder $V_{4\rho}$. We assume that

(B1) (Regularity) $b_{ij} \in \text{Lip}(V_{4\rho})$ ($1 \leq i, j \leq n-1$),

(B2) (Parity) b_{ij} is even in η_n , i.e., $b_{ij}(\eta', \eta_n) = b_{ij}(\eta', -\eta_n)$ for $\eta \in V_{4\rho}$,

(B3) (Smallness) $b_{ij}(0) = 0$ ($1 \leq i, j \leq n-1$).

For $\rho > 0$, let Y_ρ denotes the space

$$\{g \in BMO(\mathbf{R}^n) \cap L^2(\mathbf{R}^n) \mid \text{supp } g \subset V_\rho, \ g(\eta', \eta_n) = g(\eta', -\eta_n) \text{ for } \eta \in V_\rho\},$$

whereas Z_ρ denotes the space

$$\{f \in BMO(\mathbf{R}^n) \mid \text{supp } f \subset V_\rho, \ f(\eta', \eta_n) = -f(\eta', -\eta_n) \text{ for } \eta \in V_\rho\}.$$

The oddness condition in Z_ρ guarantees that

$$\frac{1}{r^n} \int_{B_r(\eta', 0)} f \, d\eta = 0$$

for any $r > 0$ and $\eta' \in \mathbf{R}^{n-1}$, which implies that

$$\frac{1}{r^n} \int_{B_r(\eta', 0)} |f| \, d\eta \leq [f]_{BMO(\mathbf{R}^n)}$$

for any $r > 0$ and $\eta' \in \mathbf{R}^{n-1}$. Hence f is L^1 in $\mathbf{R}^n$.

Lemma 10. *Assume that (B1) – (B3). Then, there exists $\rho_* > 0$ depending only on n and b such that the following property holds provided that $\rho \in (0, \rho_*)$. There exists a bounded linear operator $f \mapsto q_o$ from Z_ρ to $L^\infty(\mathbf{R}^n)$ such that*

(i)

$$[\nabla_\eta q_o]_{BMO(\mathbf{R}^n)} \leq C[f]_{BMO(\mathbf{R}^n)} \quad \text{for all } f \in Z_\rho$$

with some C independent of f ;

(ii)

$$L_0 q_o = \partial_{\eta_n} f \quad \text{in} \quad V_{2\rho};$$

(iii) q_o is even in $\mathbf{R}^n$ with respect to η_n , i.e. $q_o(\eta', \eta_n) = q_o(\eta', -\eta_n) \forall \eta \in \mathbf{R}^n$;

(iv)

$$\sup \left\{ \frac{1}{r^n} \int_{B_r(\eta', 0)} |\partial_{\eta_n} q_o| \, d\eta \mid 0 < r < \infty, \eta' \in \mathbf{R}^{n-1} \right\} \leq C[f]_{BMO(\mathbf{R}^n)}.$$

Proof. By (B3) and (B1), we observe that

$$\overline{\lim}_{\rho \downarrow 0} \|b_{ij}\|_{C^\gamma(V_{4\rho})} / \rho^{1-\gamma} < \infty$$

for any $\gamma \in (0, 1)$ and $1 \leq i, j \leq n-1$. Indeed, for $1 \leq i, j \leq n-1$, (B1) and (B3) imply that

$$\begin{aligned} \|b_{ij}\|_{L^\infty(V_{4\rho})} &\leq 8L\rho, \\ [b_{ij}]_{C^\gamma(V_{4\rho})} &:= \sup \left\{ |b_{ij}(\eta) - b_{ij}(\zeta)| / |\eta - \zeta|^\gamma \mid \eta, \zeta \in V_{4\rho} \right\} \\ &\leq L(16\rho)^{1-\gamma}, \end{aligned}$$

where L is the maximum of Lipschitz bound for b_{ij} for all $1 \leq i, j \leq n-1$. We next take a cut-off function. We take $\theta \in C_c^\infty(V_4)$ such that $\theta = 1$ on V_2 and $0 \leq \theta \leq 1$ in V_4 , we may assume θ is radial so that θ is even in η_n . We rescale θ by setting

$$\theta_\rho(\eta) = \theta(\eta/\rho)$$

so that $\theta_\rho = 1$ on $V_{2\rho}$. Since $\|\nabla \theta_\rho\|_\infty \rho$ is bounded as $\rho \rightarrow 0$, we see that

$$\overline{\lim}_{\rho \downarrow 0} [\theta_\rho]_{C^\gamma(V_{4\rho})} \rho^\gamma < \infty.$$

Hence, the estimate

$$[\theta_\rho b_{ij}]_{C^\gamma(V_{4\rho})} \leq [\theta_\rho]_{C^\gamma(V_{4\rho})} \|b_{ij}\|_{L^\infty(V_{4\rho})} + [b_{ij}]_{C^\gamma(V_{4\rho})} \|\theta_\rho\|_{L^\infty(V_{4\rho})}$$

implies that

$$\overline{\lim}_{\rho \downarrow 0} \|\theta_\rho b_{ij}\|_{C^\gamma(V_{4\rho})} / \rho^{1-\gamma} < \infty.$$

We then set

$$L_1 = A - B_1, \quad B_1 = \sum_{1 \leq i, j \leq n-1} \partial_{\eta_i} b_{ij}^\rho \partial_{\eta_j}, \quad b_{ij}^\rho = b_{ij} \theta_\rho.$$

For $1 \leq i, j \leq n-1$, notice that b_{ij}^ρ satisfies the same property of b_{ij} in (B1) – (B3). Moreover,

$$\text{supp } b_{ij}^\rho \subset V_{4\rho} \quad \text{and} \quad \|b_{ij}^\rho\|_{C^\gamma(V_{4\rho})} \leq c_b \rho^{1-\gamma}, \quad \rho > 0$$

with some c_b independent of ρ . Since $\text{supp } b_{ij}^\rho \subset V_{4\rho}$, we actually have that $b_{ij}^\rho \in C^\gamma(\mathbf{R}^n)$ together with the estimate

$$\|b_{ij}^\rho\|_{C^\gamma(\mathbf{R}^n)} \leq \|b_{ij}^\rho\|_{C^\gamma(V_{4\rho})}.$$

For a given $f \in Z_\rho$, we define q_o by

$$q_o := \sum_{k=0}^{\infty} A^{-1} (B_1 A^{-1})^k \partial_{\eta_n} f,$$

where formally for a function h we mean $A^{-1}h$ by $E * h$. The parity condition (iii) is clear once q_o is well defined as a function. Since

$$L_1 q_o = \sum_{k=0}^{\infty} (B_1 A^{-1})^k \partial_{\eta_n} f - \sum_{k=1}^{\infty} (B_1 A^{-1})^k \partial_{\eta_n} f = \partial_{\eta_n} f$$

in $\mathbf{R}^n$, the property (ii) then follows since $L_1 = L_0$ in $V_{2\rho}$.

It remains to prove the convergence of q_o as well as (i). For this purpose, we reinterpret q_o in a different way. We rewrite

$$B_1 = \text{div}' \cdot \nabla'_B \quad \text{with} \quad \nabla'_B = \left(\sum_{j=1}^{n-1} b_{ij}^\rho \partial_{\eta_j} \right)_{1 \leq i \leq n-1}$$

and observe that

$$\begin{aligned} q_o &= \sum_{k=0}^{\infty} A^{-1} \text{div}' \cdot G^k \cdot \nabla'_B A^{-1} \partial_{\eta_n} f + A^{-1} \partial_{\eta_n} f, \\ G &:= \nabla'_B A^{-1} \text{div}' . \end{aligned}$$

Denote

$$b^\rho := \left(b_{ij}^\rho \right)_{1 \leq i, j \leq n-1}.$$

Since $\partial_{\eta_\alpha} A^{-1} \partial_{\eta_\beta}$ is bounded in BMO [7] and also in L^p ($1 < p < \infty$) for all $\alpha, \beta = 1, \dots, n$, see e.g. [14, Theorem 5.2.7 and Theorem 5.2.10], by a multiplication theorem we can deduce the

estimates

$$\|Gh\|_{L^p(\mathbf{R}^n)} \leq C_p \|b^\rho\|_{L^\infty(\mathbf{R}^n)} \|h\|_{L^p(\mathbf{R}^n)}, \quad (12)$$

$$[Gh]_{BMO(\mathbf{R}^n)} \leq C'_\infty \|b^\rho\|_{C^\gamma(\mathbf{R}^n)} ([h]_{BMO(\mathbf{R}^n)} + \|h\|_{L^1(\mathbf{R}^n)}) \quad (13)$$

provided that $\text{supp } h \subset V_{4\rho}$ and $\rho < 1$. Here C_p and C'_∞ are independent of ρ and h . Similar estimate holds for $\nabla'_B A^{-1} \partial_{\eta_n}$. Since $\|f\|_{L^1(\mathbf{R}^n)} \leq C_\rho [f]_{BMO(\mathbf{R}^n)}$ for $f \in Z_\rho$, by an interpolation (cf. [4, Lemma 5]) we see that the L^p norm of f is also controlled, i.e., $\|f\|_{L^p(\mathbf{R}^n)} \leq C_\rho [f]_{BMO(\mathbf{R}^n)}$ for any $1 \leq p < \infty$. By the support condition, $A^{-1} \text{div}'$ and $A^{-1} \partial_{\eta_n}$ is bounded from $L^p \rightarrow L^\infty$ for $p > n$ with bound K , we see that

$$\begin{aligned} \|q_o\|_{L^\infty(\mathbf{R}^n)} &\leq K \left(\left\| \sum_{k=0}^{\infty} G^k \nabla'_B A^{-1} \partial_{\eta_n} f \right\|_{L^p(\mathbf{R}^n)} + \|f\|_{L^p(\mathbf{R}^n)} \right) \\ &\leq K \left(\sum_{k=0}^{\infty} C_p^{k+1} \|b^\rho\|_{L^\infty(\mathbf{R}^n)}^{k+1} \|f\|_{L^p(\mathbf{R}^n)} + \|f\|_{L^p(\mathbf{R}^n)} \right), \quad p > n. \end{aligned}$$

If ρ is taken small so that

$$\sum_{k=0}^{\infty} (C_p \cdot 8L\rho)^{k+1} < \infty,$$

then q_o converges uniformly in $\mathbf{R}^n$ and $\|q_o\|_{L^\infty(\mathbf{R}^n)} \leq C_\rho [f]_{BMO(\mathbf{R}^n)}$ with some C_ρ independent of f .

Set

$$\|h\|_{BMOL^p(\mathbf{R}^n)} := [h]_{BMO(\mathbf{R}^n)} + \|h\|_{L^p(\mathbf{R}^n)}.$$

By estimates (12) and (13), we observe that

$$\|Gh\|_{BMOL^p(\mathbf{R}^n)} \leq C_* \|b^\rho\|_{C^\gamma(\mathbf{R}^n)} \|h\|_{BMOL^p(\mathbf{R}^n)}, \quad 1 < p < \infty,$$

where $C_* = C_p + C'_\infty \cdot C_n$ with C_n independent of ρ and h . We next estimate ∇q_o . By the similar estimate for $\nabla'_B A^{-1} \text{div}'$ and $\nabla'_B A^{-1} \partial_{\eta_n}$, we have that

$$\|\nabla q_o\|_{BMOL^p(\mathbf{R}^n)} \leq \left(\sum_{k=0}^{\infty} C_*^{k+1} \|b^\rho\|_{C^\gamma(\mathbf{R}^n)}^{k+1} + C_* \|b^\rho\|_{C^\gamma(\mathbf{R}^n)} \right) \|f\|_{BMOL^p(\mathbf{R}^n)}.$$

We fix $p > n$ and take $\rho < \frac{1}{8LC_p}$ sufficiently small so that

$$\sum_{k=0}^{\infty} (C_* \cdot c_b \rho^{1-\gamma})^{k+1} < \infty.$$

Then we get our desired estimate

$$\|\nabla q_o\|_{BMOL^p(\mathbf{R}^n)} \leq C_\rho \|f\|_{BMOL^p(\mathbf{R}^n)} \leq C_\rho [f]_{BMO(\mathbf{R}^n)}$$

for $f \in Z_\rho$. This completes the proof of (i).

Since $\partial_{\eta_n} q_o$ is odd in η_n so that

$$\frac{1}{r^n} \int_{B_r(\eta', 0)} \partial_{\eta_n} q_o \, d\eta = 0$$

for any $\eta' \in \mathbf{R}^{n-1}$, the left-hand side of (iv) is estimated by a constant multiple of $[\partial_{\eta_n} q_o]_{BMO(\mathbf{R}^n)}$, which is estimated by a constant multiple of $[f]_{BMO(\mathbf{R}^n)}$. The proof of (iv) is now complete. $\square$

Similarly, we are able to establish the following which corresponds to a version of Lemma 10 for the space Y_ρ .

Lemma 11. *Assume that (B1) – (B3). Then, there exists $\rho_* > 0$ depending only on n and b such that the following property holds provided that $\rho \in (0, \rho_*)$. For each $1 \leq i \leq n-1$, there exists a bounded linear operator $g \mapsto q_{e,i}$ from Y_ρ to $L^\infty(\mathbf{R}^n)$ such that*

(i)

$$[\nabla q_{e,i}]_{BMO(\mathbf{R}^n)} \leq C \|g\|_{BMOL^2(\mathbf{R}^n)} \quad \text{for all } g \in Y_\rho$$

with some C independent of f ;

(ii)

$$L_0 q_{e,i} = \partial_{\eta_i} g \quad \text{in } V_{2\rho};$$

(iii) $q_{e,i}$ is even in $\mathbf{R}^n$ with respect to η_n , i.e. $q_{e,i}(\eta', \eta_n) = q_{e,i}(\eta', -\eta_n) \, \forall \eta \in \mathbf{R}^n$;

(iv)

$$\sup \left\{ \frac{1}{r^n} \int_{B_r(\eta', 0)} |\partial_{\eta_n} q_{e,i}| \, d\eta \mid 0 < r < \infty, \eta' \in \mathbf{R}^{n-1} \right\} \leq C \|g\|_{BMOL^2(\mathbf{R}^n)}.$$

Proof. Fix $1 \leq i \leq n-1$. Since g is even in $\mathbf{R}^n$ with respect to η_n , $\partial_{\eta_i} g$ is also even in $\mathbf{R}^n$ with respect to η_n . This means that $\partial_{\eta_i} g$ has the same parity with $\partial_{\eta_n} f$ in Lemma 10. By considering

$$q_{e,i} := \sum_{k=0}^{\infty} A^{-1} (B_1 A^{-1})^k \partial_{\eta_i} g,$$

exactly the same arguments of the proof of Lemma 10 finish the rest of the work. $\square$

We take ρ_* in Lemma 10 and Lemma 11 to be

$$\rho_* := \min \left\{ \rho_{0,*}, \frac{1}{8LC_p}, \left(\frac{1}{C_* \cdot c_b} \right)^{\frac{1}{1-\gamma}} \right\}.$$

3.2 Laplacian in a normal coordinate system

Take $z_0 \in \Gamma$. Let us recall the normal coordinate system (6) introduced in Section 2.1, i.e.,

$$\begin{cases} x' &= \eta' + \eta_n \nabla' d(\eta', h_{z_0}(\eta')); \\ x_n &= h_{z_0}(\eta') + \eta_n \partial_{\eta_n} d(\eta', h_{z_0}(\eta')) \end{cases}$$

in $U_{\rho_0}(z_0)$ with $\nabla' h_{z_0}(0') = 0$, $h_{z_0}(0') = 0$ up to translation and rotation such that $z_0 = 0$ and

$$-\mathbf{n}(\eta', h_{z_0}(\eta')) = (-\nabla' h_{z_0}(\eta'), 1) \Big/ \left(1 + |\nabla'_{z_0} h(\eta')|^2 \right)^{1/2}, \quad \eta' \in B_{\rho_0}.$$

Since Γ is C^3 , the mapping $x = \psi(\eta) \in C^2(V_{\rho_0})$ in $U_{\rho_0}(z_0)$, it is a local C^2 -diffeomorphism. Moreover, its Jacobi matrix $D\psi$ is the identity at 0, i.e.,

$$\nabla \psi(0) = I = \nabla \psi^{-1}(0).$$

A direct calculation shows that in $U_{\rho_0}(z_0) \cap \Omega$,

$$\begin{aligned} -\Delta_x &= -\Delta_\eta - \left\{ \sum_{\substack{1 \leq i, j \leq n-1 \\ i \neq j}} \gamma_{ij} \partial_{\eta_i} \partial_{\eta_j} + \sum_{j=1}^{n-1} (\gamma_{jj} - 1) \partial_{\eta_j}^2 \right\} \\ &\quad - \sum_{1 \leq i, j \leq n} \frac{\partial^2 \eta_j}{\partial x_i^2} \partial_{\eta_j}, \quad \gamma_{ij} = \sum_{k=1}^n \frac{\partial \eta_j}{\partial x_k} \frac{\partial \eta_i}{\partial x_k}. \end{aligned}$$

Note that $\gamma_{jj}(0) = 1$ while $\gamma_{ij}(0) = 0$ if $i \neq j$. Changing order of multiplication and differentiation, we conclude that

$$\begin{aligned} -\Delta_x &= \tilde{L}_0 + \tilde{M}, \\ \tilde{L}_0 &:= A - \tilde{B}, \quad A := -\Delta_\eta, \quad \tilde{B} := \sum_{1 \leq i, j \leq n-1} \partial_{\eta_i} \tilde{b}_{ij}(\eta) \partial_{\eta_j}, \\ \tilde{M} &:= \sum_{j=1}^n \tilde{c}_j(\eta) \partial_{\eta_j} \end{aligned}$$

with $\tilde{b}_{ij} = \gamma_{ij} - \delta_{ij}$, $\tilde{c}_j = -\sum_{i=1}^n \frac{\partial^2 \eta_j}{\partial x_i^2} + \sum_{i=1}^n \partial_{\eta_i} \gamma_{ij}$. Note that if $\Gamma = \partial\Omega$ is C^3 , $\tilde{b}_{ij} \in C^1(V_{\rho_0})$ and $\tilde{c}_j \in C(V_{\rho_0})$. We restrict $\tilde{b}_{ij}$, $\tilde{c}_j$ in $V_{\rho_0} \cap \mathbf{R}_+^n$ and extend to V_{ρ_0} so that the extended

function b_{ij} , c_j 's are even in V_{ρ_0} with respect to η_n , i.e., we set $b_{ij} = E_{\text{even}} r_{V_{\rho_0} \cap \mathbf{R}_+^n} \tilde{b}_{ij}$ and $c_j = E_{\text{even}} r_{V_{\rho_0} \cap \mathbf{R}_+^n} \tilde{c}_j$. By this extension, b_{ij} may not be in C^1 but still Lipschitz and $c_j \in C(V_{\rho_0})$. We set

$$B := \sum_{1 \leq i, j \leq n-1} \partial_{\eta_i} b_{ij}(\eta) \partial_{\eta_j},$$

$$M := \sum_{j=1}^n c_j(\eta) \partial_{\eta_j}$$

and

$$L := L_0 + M, \quad L_0 = A - B.$$

The operator L may not agree with $-\Delta_x$ outside $U_{\rho_0}(z_0) \cap \Omega$. We summarize what we observe so far.

Proposition 12. *Let $\Gamma = \partial\Omega$ be C^3 and ρ_0 be chosen as in Section 2.1. For $z_0 \in \Gamma$, L_0 satisfies (B1) – (B3). Moreover, $-\Delta_x = L$ in $U_{\rho_0}(z_0) \cap \Omega$ and the coefficient of M is in $C(V_{\rho_0})$.*

Although we do not use the explicit formula of Δ in normal coordinates, we give it for $n = 2$ when we take the arc length parameter s to represent Γ . The coordinate transform is of the form

$$x_1 = \phi_1(x) + r\phi_2'(s)$$

$$x_2 = \phi_2(x) - r\phi_1'(s)$$

with $\phi_1'^2 + \phi_2'^2 = 1$ and $r = d(x)$. A direct calculation yields

$$-\Delta_x = -\Delta_{s,r} - \partial_s \left(\frac{1}{J^2} - 1 \right) \partial_s - \frac{\partial_s J}{J^3} \partial_s - \frac{1}{r} \left(1 - \frac{1}{J} \right) \partial_r,$$

where $J = 1 + r\kappa$ and κ is the curvature. We see that that the even extension of coefficient does not agree with $-\Delta_x$ outside Ω .

3.3 bmo invariant under local C^1 -diffeomorphism

Before we give the proofs to Proposition 8 and 9, we shall first establish the fact that the bmo estimate of a compactly supported function is preserved under a local C^1 -diffeomorphism. Let $V, U \subset \mathbf{R}^n$ be two domains, we consider a local C^1 -diffeomorphism $\psi : V \mapsto U$. Suppose that

$$\|\nabla_\eta \psi\|_{L^\infty(V)} + \|\nabla_x \psi^{-1}\|_{L^\infty(U)} < \infty.$$

Let $\rho > 0$. Assume that there exist two bounded subdomains $V_\rho \subset V, U_\rho \subset U$ such that $\psi : V_\rho \mapsto U_\rho$ is also a local C^1 -diffeomorphism. Set

$$K_* := \max \{1, \|\nabla_\eta \psi\|_{L^\infty(V)} + \|\nabla_x \psi^{-1}\|_{L^\infty(U)}\}.$$

We assume further that there exists a constant c_0 such that for some $\eta_0 \in V_\rho$,

$$V_\rho \subset B_{c_0\rho}(\eta_0) \subset B_{K_*(c_0+3)\rho}(\eta_0) \subset V, \quad U_\rho \subset B_{c_0\rho}(x_0) \subset B_{K_*(c_0+3)\rho}(x_0) \subset U$$

where $x_0 = \psi(\eta_0)$.

Proposition 13. *Let $f \in bmo(\mathbf{R}^n)$ with $\text{supp } f \subset V_\rho$, then $f \circ \psi^{-1} \in bmo(\mathbf{R}^n)$ satisfies*

$$\|f \circ \psi^{-1}\|_{bmo(\mathbf{R}^n)} \leq C_\rho \|f\|_{bmo(\mathbf{R}^n)}.$$

Proof. Since $\text{supp } f \circ \psi^{-1} \subset U_\rho$, we can treat $f \circ \psi^{-1}$ as a function in $\mathbf{R}^n$ with value zero outside U_ρ . The compactness of V_ρ in $\mathbf{R}^n$ implies that $\|f\|_{bmo(\mathbf{R}^n)} = \|f\|_{BMO L^1(\mathbf{R}^n)}$. Thus, the L^1 estimate

$$\|f \circ \psi^{-1}\|_{L^1(\mathbf{R}^n)} \leq C \|f\|_{L^1(\mathbf{R}^n)}$$

is obvious. Since $\psi \in C^1(V_\rho)$, an equivalent definition of the BMO -seminorm (cf. [14, Proposition 3.1.2]) implies that

$$[f \circ \psi^{-1}]_{BMO^\infty(B_{(c_0+1)\rho}(x_0))} \leq \|\nabla_x \psi^{-1}\|_{L^\infty(U)}^n \cdot \|\nabla_\eta \psi\|_{L^\infty(V)} \cdot [f]_{BMO(\mathbf{R}^n)}.$$

As $U_\rho \subset B_{c_0\rho}(x_0)$, by the extension theorem of bmo functions [11, Theorem 12], we obtain that

$$\|f \circ \psi^{-1}\|_{bmo(\mathbf{R}^n)} \leq C_\rho \|f \circ \psi^{-1}\|_{bmo_\infty(B_{(c_0+1)\rho}(x_0))} \leq C_\rho \|f\|_{bmo(\mathbf{R}^n)}.$$

□

Similarly, if $g \in bmo(\mathbf{R}^n)$ with $\text{supp } g \subset U_\rho$, then we have that $g \circ \psi \in bmo(\mathbf{R}^n)$ satisfying

$$\|g \circ \psi\|_{bmo(\mathbf{R}^n)} \leq C_\rho \|g\|_{bmo(\mathbf{R}^n)}.$$

Even if we are considering vector fields instead of scalar functions, similar results hold.

Proposition 14. *Let $\nabla_\eta f \in bmo(\mathbf{R}^n)$ with $\text{supp } \nabla_\eta f \subset V_\rho$, then $\nabla_x(f \circ \psi^{-1}) \in bmo(\mathbf{R}^n)$ satisfying*

$$\|\nabla_x(f \circ \psi^{-1})\|_{bmo(\mathbf{R}^n)} \leq C_\rho \|\nabla_\eta f\|_{bmo(\mathbf{R}^n)}.$$

Proof. Since $\nabla_\eta f$ is compactly supported, the L^1 estimate

$$\|\nabla_x(f \circ \psi^{-1})\|_{L^1(\mathbf{R}^n)} \leq C \|\nabla_\eta f\|_{L^1(\mathbf{R}^n)}$$

is obvious. Pick a cut-off function $\theta_{*,\rho} \in C_c^\infty(B_{K_*(c_0+3)\rho}(\eta_0))$ such that $\theta_{*,\rho} = 1$ in $B_{K_*(c_0+2)\rho}(\eta_0)$. Consider $B_r(x) \subset B_{(c_0+1)\rho}(x_0)$ with $r < \rho$. Let $\eta = \psi^{-1}(x)$. Since $\psi^{-1}(B_r(x)) \subset B_{K_*(c_0+2)\rho}(\eta_0)$, we have that

$$\frac{1}{r^n} \int_{B_r(x)} |\partial_{x_i}(f \circ \psi^{-1}) - c| dy \leq \frac{K_*}{r^n} \int_{\psi^{-1}(B_r(x))} \left| \sum_{1 \leq l \leq n} \theta_{*,\rho} \left(\frac{\partial \eta_l}{\partial x_i} \right)_\psi \partial_{\eta_l} f - c \right| d\eta$$

for any $c \in \mathbf{R}^n$, $1 \leq i \leq n$. By considering an equivalent definition of the BMO -seminorm, see e.g. [14, Proposition 3.1.2], we deduce that

$$\begin{aligned} [\nabla_x(f \circ \psi^{-1})]_{BMO^\infty(B_{(c_0+1)\rho}(x_0))} &\leq K_*^{n+1} \left[\sum_{1 \leq i, l \leq n} \theta_{*,\rho} \left(\frac{\partial \eta_l}{\partial x_i} \right)_\psi \partial_{\eta_l} f \right]_{BMO(\mathbf{R}^n)} \\ &\leq C_\rho \|\nabla_\eta f\|_{bmo(\mathbf{R}^n)}. \end{aligned}$$

As $U_\rho \subset B_{c_0\rho}(x_0)$, by the extension theorem of bmo functions [11, Theorem 12], we obtain that

$$\|\nabla_x(f \circ \psi^{-1})\|_{bmo(\mathbf{R}^n)} \leq C_\rho \|\nabla_x(f \circ \psi^{-1})\|_{bmo_\infty(B_{(c_0+1)\rho}(x_0))} \leq C_\rho \|\nabla_\eta f\|_{bmo(\mathbf{R}^n)}.$$

□

If $\nabla_x g \in bmo(\mathbf{R}^n)$ with $\text{supp } \nabla_x g \subset U_\rho$, same proof of Proposition 14 shows that $\nabla_\eta(g \circ \psi) \in bmo(\mathbf{R}^n)$ satisfying

$$\|\nabla_\eta(g \circ \psi)\|_{bmo(\mathbf{R}^n)} \leq C_\rho \|\nabla_x g\|_{bmo(\mathbf{R}^n)}.$$

Let h be either a scalar function or a vector field which is compactly supported in U_ρ , for simplicity of notations we denote $h_\psi := h \circ \psi$. If h is a vector field, we denote $h_{\psi,i} := h_i \circ \psi$ for $1 \leq i \leq n$.

3.4 Volume potential for tangential component

Let $\rho \in (0, \rho_*/2)$ and fix $1 \leq j \leq m$. Since $\varphi_j v_2 \in vBMO(\Omega)$ with $\text{supp } \varphi_j v_2 \subset U_{\rho,j} \cap \overline{\Omega}$, Proposition 6 implies that $(\varphi_j v_2)_{\text{even}} \in BMOL^1(\mathbf{R}^n)$. By the product estimate for bmo functions [11, Theorem 13], we see that $w_j^{\text{tan}} = Q(\varphi_j v_2) \in BMOL^1(\mathbf{R}^n)$ with $\text{supp } w_j^{\text{tan}} \subset U_{\rho,j}$. For simplicity of notations, we set $v_{2,j} := (\varphi_j v_2)_{\text{even}}$.

Let $\psi : V_{4\rho} \mapsto U_{4\rho,j}$ be the normal coordinate change defined by (6) in Section 2.1. Since

$\rho < \rho_*/2$, we have that

$$V_{4\rho} \subset B_{12\rho}(0) \subset B_{24L^*\rho}(0) \subset V_{\rho_0}, \quad U_{4\rho,j} \subset B_{12\rho}(z_j) \subset B_{24L^*\rho}(z_j) \subset U_{\rho_0,j}.$$

By Proposition 13 and 14, we see that ψ , in this case, is a local C^2 -diffeomorphism that preserves bmo estimates for functions or vector fields compactly supported in $V_{4\rho}$. As a result, $(v_{2,j})_\psi \in BMOL^1(\mathbf{R}^n)$ satisfies the estimate

$$\|(v_{2,j})_\psi\|_{BMOL^1(\mathbf{R}^n)} \leq C_\rho \|v_{2,j}\|_{BMOL^1(\mathbf{R}^n)}.$$

Note that similar conclusions hold if we consider $\psi^{-1} : U_{4\rho,j} \mapsto V_{4\rho}$ instead.

Proposition 8. For $1 \leq i \leq n$ and $1 \leq k \leq n-1$, we define

$$\left(\frac{\partial \eta_k}{\partial x_i}\right)_* := E_{\text{even}} r_{V_{4\rho} \cap \mathbf{R}_+^n} \left(\frac{\partial \eta_k}{\partial x_i}\right)_\psi \quad \text{and} \quad g_{i,k} := \left(\frac{\partial \eta_k}{\partial x_i}\right)_* \cdot (v_{2,j})_{\psi,i}.$$

We consider

$$\begin{aligned} (\operatorname{div}_x w_j^{\tan})_{\psi,*} &:= \sum_{\substack{1 \leq i \leq n, \\ 1 \leq k \leq n-1}} \left\{ \partial_{\eta_k} g_{i,k} - \partial_{\eta_k} \left(\frac{\partial \eta_k}{\partial x_i}\right)_\psi \cdot (v_{2,j})_{\psi,i} \right\} \\ &\quad - \sum_{\substack{1 \leq i \leq n, \\ 1 \leq k \leq n-1}} \left(\frac{\partial \eta_k}{\partial x_i}\right)_\psi \cdot \left(\sum_{1 \leq l \leq n} (v_{2,j})_{\psi,l} \cdot \left(\frac{\partial \eta_l}{\partial x_i}\right)_\psi \right) \cdot \frac{\partial^2 x_i}{\partial \eta_k \partial \eta_n} \end{aligned}$$

in $V_{4\rho} = \psi^{-1}(U_{4\rho,j})$. Let $L = L_0 + M$ be the operator in Proposition 12 and L_0^{-1} be the operator in Lemma 11. Let $1 \leq i \leq n$ and $1 \leq k \leq n-1$. We set

$$q_{j,1,\psi}^{i,k} := -\theta_\rho L_0^{-1} \partial_{\eta_k} g_{i,k}$$

where θ_ρ is the cut-off function defined in the proof of Lemma 10. There exists $\overline{(\frac{\partial \eta_k}{\partial x_i})_*} \in C^{0,1}(\mathbf{R}^n)$, see e.g. [11, Theorem 13], such that the restriction of $\overline{(\frac{\partial \eta_k}{\partial x_i})_*}$ in $V_{4\rho}$ equals $(\frac{\partial \eta_k}{\partial x_i})_*$ and $\|\overline{(\frac{\partial \eta_k}{\partial x_i})_*}\|_{C^{0,1}(\mathbf{R}^n)} \leq \|(\frac{\partial \eta_k}{\partial x_i})_*\|_{C^{0,1}(V_{4\rho})}$. By viewing $g_{i,k}$ as $\overline{(\frac{\partial \eta_k}{\partial x_i})_*} \cdot (v_{2,j})_{\psi,i}$, we see that $g_{i,k} \in BMOL^1(\mathbf{R}^n)$. Hence, $q_{j,1,\psi}^{i,k} \in L^\infty(\mathbf{R}^n)$ is well-defined which satisfies all conditions in Lemma 11. Let $f_{j,1,\psi}^{i,k} := M\theta_\rho L_0^{-1} \partial_{\eta_k} g_{i,k}$. We can define

$$q_{j,1}^{i,k} := q_{j,1,\psi}^{i,k} \circ \psi^{-1}, \quad f_{j,1}^{i,k} := f_{j,1,\psi}^{i,k} \circ \psi^{-1}$$

in $U_{\rho_0,j}$. Notice that $\operatorname{supp} q_{j,1}^{i,k}, \operatorname{supp} f_{j,1}^{i,k} \subset U_{4\rho,j}$, we can indeed treat $q_{j,1}^{i,k}, f_{j,1}^{i,k}$ as functions defined in $\mathbf{R}^n$ where their values outside $U_{4\rho,j}$ equal zero. Proposition 14 shows that $\nabla_x q_{j,1}^{i,k} \in BMO(\mathbf{R}^n)$

satisfies the estimate

$$[\nabla_x q_{j,1}^{i,k}]_{BMO(\mathbf{R}^n)} \leq C_\rho \|\nabla_\eta q_{j,1,\psi}^{i,k}\|_{BMO L^2(\mathbf{R}^n)} \leq C_\rho \|g_{i,k}\|_{BMO L^2(\mathbf{R}^n)}.$$

Let $p_{j,1}^{i,k} := E * f_{j,1}^{i,k}$. By Lemma 11 again, we can prove that

$$\|p_{j,1}^{i,k}\|_{L^\infty(\mathbf{R}^n)} + \|\nabla_x p_{j,1}^{i,k}\|_{L^\infty(\mathbf{R}^n)} \leq C_\rho \|f_{j,1,\psi}^{i,k}\|_{L^p(V_{2\rho})} \leq C_\rho \|g_{i,k}\|_{L^p(\mathbf{R}^n)}$$

with some $p > n$. Thus, $p_{j,1}^{i,k}$ is well-defined. By Proposition 6, we have that

$$\|g_{i,k}\|_{BMO L^1(\mathbf{R}^n)} \leq C_\rho \|v_{2,j}\|_{BMO L^1(\mathbf{R}^n)} \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

Hence, by an interpolation (cf. [4, Lemma 5]),

$$\|g_{i,k}\|_{L^p(\mathbf{R}^n)} \leq C_\rho \|v\|_{vBMO(\Omega)}$$

for any $1 < p < \infty$.

For lower order term $q_{j,2,\psi}^{i,k} := \partial_{\eta_k}(\frac{\partial \eta_k}{\partial x_i})_\psi \cdot (v_{2,j})_{\psi,i}$, we set $q_{j,2}^{i,k} := q_{j,2,\psi}^{i,k} \circ \psi^{-1}$ in $U_{\rho_0,j}$. Similar as $q_{j,1}^{i,k}$, we can treat $q_{j,2}^{i,k}$ as a function in $\mathbf{R}^n$ with value zero outside $U_{\rho,j}$ since $\text{supp } q_{j,2}^{i,k} \subset U_{\rho,j}$. Define $p_{j,2}^{i,k} := E * q_{j,2}^{i,k}$. Since E and $\nabla_x E$ are locally integrable, we have that

$$\|p_{j,2}^{i,k}\|_{L^\infty(\mathbf{R}^n)} + \|\nabla_x p_{j,2}^{i,k}\|_{L^\infty(\mathbf{R}^n)} \leq C_\rho \|q_{j,2,\psi}^{i,k}\|_{L^p(V_\rho)} \leq C_\rho \|v_{2,j}\|_{L^p(U_{\rho,j})}$$

for some $p > n$. By an interpolation (cf. [4, Lemma 5]) again, we deduce that

$$\|p_{j,2}^{i,k}\|_{L^\infty(\mathbf{R}^n)} + \|\nabla_x p_{j,2}^{i,k}\|_{L^\infty(\mathbf{R}^n)} \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

This argument also holds for lower order term

$$q_{j,3,\psi}^{i,k} := \left(\frac{\partial \eta_k}{\partial x_i}\right)_\psi \cdot \left(\sum_{1 \leq l \leq n} (v_{2,j})_{\psi,l} \cdot \left(\frac{\partial \eta_n}{\partial x_l}\right)_\psi\right) \cdot \frac{\partial^2 x_i}{\partial \eta_k \partial \eta_n}.$$

By letting $q_{j,3}^{i,k} := q_{j,3,\psi}^{i,k} \circ \psi^{-1}$ in $U_{\rho_0,j}$ and $p_{j,3}^{i,k} := E * q_{j,3}^{i,k}$, we can show that

$$\|p_{j,3}^{i,k}\|_{L^\infty(\mathbf{R}^n)} + \|\nabla_x p_{j,3}^{i,k}\|_{L^\infty(\mathbf{R}^n)} \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

Set

$$p_j^{\text{tan}} := \sum_{\substack{1 \leq i \leq n, \\ 1 \leq k \leq n-1}} (q_{j,1}^{i,k} + p_{j,1}^{i,k} + p_{j,2}^{i,k} + p_{j,3}^{i,k}).$$

Since a direct calculation implies that

$$\begin{aligned} (\operatorname{div}_x w_j^{\tan})_\psi &= \sum_{\substack{1 \leq i \leq n, \\ 1 \leq k \leq n-1}} \left(\frac{\partial \eta_k}{\partial x_i} \right)_\psi \cdot \partial_{\eta_k} (v_{2,j})_{\psi,i} \\ &\quad - \sum_{\substack{1 \leq i \leq n, \\ 1 \leq k \leq n-1}} \left(\frac{\partial \eta_k}{\partial x_i} \right)_\psi \cdot \left(\sum_{1 \leq l \leq n} (v_{2,j})_{\psi,l} \cdot \left(\frac{\partial \eta_n}{\partial x_l} \right)_\psi \right) \cdot \frac{\partial^2 x_i}{\partial \eta_k \partial \eta_n} \end{aligned}$$

in normal coordinate in $V_{4\rho} = \psi^{-1}(U_{4\rho,j})$, it is easy to see that

$$-\Delta_x p_j^{\tan} = \operatorname{div} w_j^{\tan}$$

in $U_{2\rho,j} \cap \Omega$. Calculations above ensures that

$$[\nabla_x p_j^{\tan}]_{BMO(\mathbf{R}^n)} \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

Since $\operatorname{supp} q_{j,1}^{i,k} \subset U_{4\rho,j}$, we consider $x \in \Gamma$ and $r < \rho$ such that $B_r(x) \cap U_{4\rho,j} \neq \emptyset$. By change of variables $y = \psi(\eta)$ in $U_{4\rho,j}$, we deduce that

$$\int_{B_r(x) \cap U_{4\rho,j}} |\nabla_y q_{j,1}^{i,k} \cdot \nabla_y d| dy \leq C \int_{B_{L_*r}(\zeta)} |\partial_{\eta_n} q_{j,1,\psi}^{i,k}| d\eta$$

where $\zeta = \psi^{-1}(x)$ and $\zeta_n = 0$. By Lemma 11, we see that

$$\int_{B_{L_*r}(\zeta)} |\partial_{\eta_n} q_{j,1,\psi}^{i,k}| d\eta \leq r^n C_\rho \|v\|_{vBMO(\Omega)}.$$

Since $\nabla_x p_{j,l}^{i,k} \in L^\infty(\mathbf{R}^n)$ for $l = 1, 2, 3$, we finally obtain that

$$\frac{1}{r^n} \int_{B_r(x)} |\nabla_y p_j^{\tan} \cdot \nabla_y d| dy \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

□

3.5 Volume potential for normal component

Consider $\rho \in (0, \rho_*/2)$ and $1 \leq j \leq m$. We let $g_j := \nabla d \cdot (\varphi_j v_2)_{\text{odd}}$. Since $\varphi_j v_2 \in vBMO(\Omega)$ with $\operatorname{supp} \varphi_j v_2 \subset U_{\rho,j} \cap \overline{\Omega}$, by Proposition 6 we see that $g_j \in BMO(\mathbf{R}^n) \cap b^\nu(\Gamma)$. In particular, we have the estimate

$$[g_j]_{BMO(\mathbf{R}^n)} + [g_j]_{b^\nu(\Gamma)} \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

Considering the normal coordinate in $U_{4\rho,j}$, g_j is odd in η_n . Note that $w_j^{\text{nor}} = g_j \nabla d$.

Proposition 9. Since $\nabla d \in C^1(U_{\rho_0,j})$, by Proposition 6 we have that

$$[w_j^{\text{nor}}]_{BMO(\mathbf{R}^n)} \leq C \|\nabla d\|_{C^\gamma(U_{\rho_0,j})} \|g_j\|_{BMOL^1(\mathbf{R}^n)} \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

We note that

$$\operatorname{div}_x w_j^{\text{nor}} = \nabla_x g_j \cdot \nabla_x d + g_j \Delta_x d.$$

Let $g_{j,\psi} := g_j \circ \psi$ in $U_{\rho_0,j}$. We may treat $g_{j,\psi}$ as a function in $\mathbf{R}^n$ with value zero outside V_ρ . By Proposition 13, we have that

$$[g_{j,\psi}]_{BMO(\mathbf{R}^n)} \leq C_\rho \|g_j\|_{BMOL^1(\mathbf{R}^n)}.$$

In normal coordinate, $\nabla_x g_j \cdot \nabla_x d = \partial_{\eta_n} g_{j,\psi}$. We introduce the operator $L = L_0 + M$ in Proposition 12. Since $g_{j,\psi} \in Z_\rho$, we set

$$p_{1,j,\psi} := \theta_\rho L_0^{-1} \partial_{\eta_n} g_{j,\psi}$$

where θ_ρ is the cut-off function of $V_{2\rho}$ in the proof of Lemma 10. $p_{1,j,\psi}$ satisfies all conditions in Lemma 10. Set $f_{j,\psi} := -M \theta_\rho L_0^{-1} \partial_{\eta_n} g_{j,\psi}$. We define

$$p_{1,j} := p_{1,j,\psi} \circ \psi^{-1}, f_j := f_{j,\psi} \circ \psi^{-1}$$

in $U_{\rho_0,j}$. Notice that $p_{1,j} \in L^\infty(\mathbf{R}^n)$ and $f_j \in L^p(\mathbf{R}^n)$ with some $p > n$. By Proposition 14,

$$[\nabla_x p_{1,j}]_{BMO(\mathbf{R}^n)} \leq C_\rho [\nabla_\eta p_{1,j,\psi}]_{BMO(\mathbf{R}^n)} \leq C_\rho [g_{j,\psi}]_{BMO(\mathbf{R}^n)}.$$

Set

$$p_j^{\text{nor}} = p_{1,j} + p_{2,j} + p_{3,j}$$

with $p_{2,j} = E * f_j$ and $p_{3,j} = E * (g_j \Delta_x d)$. This p_j^{nor} satisfies all desired properties required. For lower order terms $p_{2,j}$ and $p_{3,j}$, we have that

$$\|p_2\|_{L^\infty(\mathbf{R}^n)} + \|\nabla p_2\|_{L^\infty(\mathbf{R}^n)} + \|\nabla p_3\|_{L^\infty(\mathbf{R}^n)} + \|p_3\|_{L^\infty(\mathbf{R}^n)} \leq C_\rho \|g_j\|_{L^p(\mathbf{R}^n)}$$

as E and $\nabla_x E$ are both locally integrable. By an interpolation (cf. [4, Lemma 5]), we obtain that

$$[\nabla_x p_j^{\text{nor}}]_{BMO(\mathbf{R}^n)} \leq C_\rho \|g_j\|_{BMOL^1(\mathbf{R}^n)} \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

Since $\operatorname{supp} p_{1,j} \subset U_{\rho,j}$, we consider $x \in \Gamma$ and $r < \rho$ such that $B_r(x) \cap U_{\rho,j} \neq \emptyset$. Set $\zeta = \psi^{-1}(x)$

with $\zeta_n = 0$. Consider change of variable $y = \psi(\eta)$ in $U_{4\rho,j}$, by Lemma 10 we see that

$$\int_{B_r(x) \cap U_{\rho,j}} |\nabla_y d \cdot \nabla_y p_{1,j}| dy \leq C \int_{B_{L_*r}(\zeta)} |\partial_{\eta_n} p_{1,j,\psi}| d\eta \leq C_\rho [g_{j,\psi}]_{BMO(\mathbf{R}^n)}.$$

By the L^∞ -estimates of $\nabla_y p_2$ and $\nabla_y p_3$, we get that

$$\frac{1}{r^n} \int_{B_r(x)} |\nabla_y d \cdot \nabla_y p_j^{\text{nor}}| dy \leq C_\rho \|v\|_{vBMO(\Omega)}.$$

Finally, a simple substitution shows that

$$-\Delta_x p_j^{\text{nor}} = \nabla_x d \cdot \nabla_x g_j - f_j + f_j + g_j \Delta_x d = \text{div}_x w_j^{\text{nor}}$$

in $U_{2\rho}(z_0) \cap \Omega$. □

4 Neumann problem with bounded data

We consider the Neumann problem for the Laplace equation problem (3) for the Laplace equation. If Ω is a smooth bounded domain, as well-known, for $g \in H^{-1/2}(\Gamma)$, there is a unique (up to constant) weak solution $u \in H^1(\Omega)$ provided that g fulfills the compatibility condition

$$\int_{\Gamma} g d\mathcal{H}^{n-1} = 0; \tag{14}$$

see e.g. [20]. The main goal of this section is to prove that ∇u belongs to $vBMO^{\infty,\infty}(\Omega)$ provided that $g \in L^\infty(\Gamma)$. In other words, we prove Lemma 4.

To prove Lemma 4, we represent the solution by using the Neumann-Green function. Let $N(x, y)$ be the Green function, i.e., a solution v of

$$\begin{aligned} -\Delta_x v &= \delta(x - y) - |\Omega|^{-1} && \text{in } \Omega \\ \frac{\partial v}{\partial \mathbf{n}_x} &= 0 && \text{on } \partial\Omega \end{aligned}$$

for $y \in \Omega$. It is easy to see that the solution u of (3) satisfying $\int_{\Omega} u dx = 0$ is given as

$$u(x) = \int_{\Gamma} N(x, y) g(y) d\mathcal{H}^{n-1}(y).$$

The function N is decomposed as

$$N(x, y) = E(x - y) + h(x, y),$$

where $h \in C^\infty(\Omega \times \Omega)$ is a milder part. We recall $h(x, y) = h(y, x)$ and

$$\sup_{x \in \Omega} \int_{\Omega} \left| \nabla_y^k h(x, y) \right|^{1+\delta} dy < \infty$$

for $k = 0, 1, 2$ with some $\delta > 0$; see [12, Lemma 3.1]. In particular, by applying the standard L^p estimate for the Neumann problem in the proof of [12, Lemma 3.1] to $\nabla_y h(\cdot, y)$, we can deduce that

$$\sup_{x \in \Omega} \int_{\Omega} |\nabla_x \nabla_y h(x, y)|^{1+\delta} dy < \infty.$$

Hence, we see that $\nabla_x h(x, \cdot) \in W^{1,1+\delta}(\Omega_y)$. By the trace theorem for Sobolev space $W^{1,1+\delta}(\Omega_y)$, this yields

$$M_0 := \sup_{x \in \Omega} \int_{\Gamma} |\nabla_x h(x, y)|^{1+\delta} d\mathcal{H}^{n-1}(y) < \infty. \quad (15)$$

We decompose u as

$$u(x) = E * (\delta_{\Gamma} \otimes g) + \int_{\Gamma} h(x, y) g(y) d\mathcal{H}^{n-1}(y) = I + II.$$

The estimate (15) yields

$$\|\nabla II\|_{L^\infty(\Omega)} \leq M_0 \|g\|_{L^\infty(\Gamma)},$$

so to prove Lemma 4 it suffices to estimate ∇I . In other words, Lemma 4 follows from the next lemma.

Lemma 15. *Let Ω be a bounded domain in $\mathbf{R}^n$ with C^2 boundary $\Gamma = \partial\Omega$.*

(i) *(BMO estimate) There exists a constant C_1 such that*

$$[\nabla (E * (\delta_{\Gamma} \otimes g))]_{BMO(\mathbf{R}^n)} \leq C_1 \|g\|_{L^\infty(\Gamma)} \quad (16)$$

for all $g \in L^\infty(\Gamma)$.

(ii) *(L^∞ estimate for normal component) There exists a constant C_2 such that*

$$\|\nabla d \cdot \nabla (E * (\delta_{\Gamma} \otimes g))\|_{L^\infty(\Gamma_{\rho_0}^{\mathbf{R}^n} \cap \Omega)} \leq C_2 \|g\|_{L^\infty(\Gamma)} \quad (17)$$

for all $g \in L^\infty(\Gamma)$.

Here $E * (\delta_{\Gamma} \otimes g)$ is defined as $E * (\delta_{\Gamma} \otimes g)(x) := \int_{\Gamma} E(x - y) g(y) d\mathcal{H}^{n-1}(y)$ for a function g on Γ . We shall prove Lemma 15 in following subsections.

4.1 BMO estimate

To see the idea, we shall prove (16) when Γ is flat. Let $\Gamma = \partial\mathbf{R}_+^n$ and $\mathbf{R}_+^n = \{(x_1, \dots, x_n) \mid x_n > 0\}$. In this case,

$$\nabla(E * (\delta_\Gamma \otimes g)) = \nabla \partial_{x_n} E * 1_{\mathbf{R}_+^n} \tilde{g}$$

where $\tilde{g} \in L^\infty(\mathbf{R}^n)$ is defined by $\tilde{g}(x', x_n) := g(x', 0)$ for any $x \in \mathbf{R}^n$. By the L^∞ -BMO estimate for the singular integral operator [15, Theorem 4.2.7], we obtain (16) when $\Gamma = \partial\mathbf{R}_+^n$.

Lemma 15 (i). Note that the signed distance function d is C^2 in $\Gamma_{\rho_0}^{\mathbf{R}^n}$, see [13, Section 14.6]. Let $\delta \in \rho_0/2$. We take a C^2 cut-off function $\theta \geq 0$ such that $\theta(\sigma) = 1$ for $\sigma \leq 1$ and $\theta(\sigma) = 0$ for $\sigma \geq 2$. By the choice of δ , we see that $\theta_d = \theta(d/\delta)$ is C^2 in $\mathbf{R}^n$. We extend $g \in L^\infty(\Gamma)$ to $g_e \in L^\infty(\Gamma_{2\delta}^{\mathbf{R}^n})$ by setting

$$g_e(x) := g(\pi x)$$

for any $x \in \Gamma_{2\delta}^{\mathbf{R}^n}$ with πx denoting the projection of x on Γ . For $x \in \Gamma_{2\delta}^{\mathbf{R}^n}$, by considering the normal coordinate $x = \psi(\eta)$ in $U_{2\delta}(\pi x)$, we have that

$$(\nabla_x d)_\psi \cdot (\nabla_x g_e)_\psi = \partial_{\eta_n}(g_e)_\psi = 0$$

as $(g_e)_\psi(\eta', \alpha) = (g_e)_\psi(\eta', \beta)$ for any $|\eta'| < 2\delta$ and $\alpha, \beta \in (-2\delta, 2\delta)$. Hence, we see that $\nabla d \cdot \nabla g_e = 0$ in $\Gamma_{2\delta}^{\mathbf{R}^n}$.

Let us consider $g_{e,c} := \theta_d g_e$. A key observation is that

$$\begin{aligned} \delta_\Gamma \otimes g &= (\nabla 1_\Omega \cdot \nabla d) g_{e,c} \\ &= \operatorname{div}(g_{e,c} 1_\Omega \nabla d) - 1_\Omega \operatorname{div}(g_{e,c} \nabla d), \\ \operatorname{div}(g_{e,c} \nabla d) &= g_{e,c} \Delta d + \nabla d \cdot \nabla g_{e,c} = g_{e,c} \Delta d + \frac{\theta'(d/\delta)}{\delta} g_e. \end{aligned}$$

Thus

$$\nabla E * (\delta_\Gamma \otimes g) = \nabla \operatorname{div}(E * (g_{e,c} 1_\Omega \nabla d)) - \nabla E * (1_\Omega g_e f_{\theta,\delta}) = I_1 + I_2$$

where $f_{\theta,\delta} := \theta_d \Delta d + \frac{\theta'(d/\delta)}{\delta}$. By the L^∞ -BMO estimate for the singular integral operator [15, Theorem 4.2.7], the first term is estimated as

$$[I_1]_{BMO(\mathbf{R}^n)} \leq C \|g_{e,c} \nabla d\|_{L^\infty(\Omega)} \leq C \|g\|_{L^\infty(\Gamma)}.$$

Since

$$A = \sup_{x \in \mathbf{R}^n \setminus \{0\}} |x|^{n-1} |\nabla E(x)| < \infty,$$

for $x \in \mathbf{R}^n$ with $d(x, \Omega) = \inf_{y \in \Omega} |x - y| < 1$ we have that

$$|I_2(x)| \leq A \int_{\Omega} \frac{1}{|x - y|^{n-1}} dy \|f_{\theta, \delta}\|_{L^\infty(\Gamma_{2\delta}^{\mathbf{R}^n})} \|g_{e, c}\|_{L^\infty(\Gamma_{2\delta}^{\mathbf{R}^n})} \leq C_{\Omega, \delta} \|g\|_{L^\infty(\Gamma)}$$

with $C_{\Omega, \delta}$ depending only on Ω and δ . For $x \in \mathbf{R}^n$ with $d(x, \Omega) = \inf_{y \in \Omega} |x - y| \geq 1$, the above estimate is trivial as $|x - y|^{-(n-1)} \leq 1$ for any $y \in \Omega$. The proof of (i) is now complete. $\square$

4.2 Estimate for normal derivative

We shall estimate normal derivative of E .

Lemma 16. *Let Ω be a bounded domain in $\mathbf{R}^n$ with C^2 boundary Γ . Then*

(i)

$$\int_{\Gamma} \frac{\partial E}{\partial \mathbf{n}_y}(x - y) d\mathcal{H}^{n-1}(y) = -1 \quad \text{for } x \in \Omega,$$

(ii)

$$\sup_{x \in \Omega} \int_{\Gamma} \left| \frac{\partial E}{\partial \mathbf{n}_y}(x - y) \right| d\mathcal{H}^{n-1}(y) < \infty.$$

Proof. (i) This follows from the Gauss divergence theorem. We observe that

$$\int_{\Gamma} \frac{\partial E}{\partial \mathbf{n}_y}(x - y) d\mathcal{H}^{n-1}(y) = \int_{\Omega} \Delta_y E(x - y) dy.$$

Since $\Delta_y E(x - y) = -\delta(x - y)$, we obtain

$$\int_{\Gamma} \frac{\partial E}{\partial \mathbf{n}_y}(x - y) d\mathcal{H}^{n-1}(y) = -1$$

for $x \in \Omega$.

(ii) We recall our local coordinate patches $\{U_i\}_{i=1}^m$ with $U_i = U_{\rho, i}$ as in Section 2.1. For $x \in \Omega^\rho$ and $y \in \Gamma$, obviously $|\nabla E(x - y)| \leq C\rho^{-(n-1)}$. Let $x \in \Gamma_\rho^{\mathbf{R}^n} \cap \Omega$. If $d(x, U_i \cap \Gamma) \geq \rho$, similarly $|\nabla E(x - y)| \leq C\rho^{-(n-1)}$ for $y \in U_i \cap \Gamma$. Hence, it is sufficient to consider U_i such that $d(x, U_i \cap \Gamma) < \rho$, i.e., it suffices to prove

$$\int_{U_i \cap \Gamma} \left| \frac{\partial E}{\partial \mathbf{n}_y}(x - y) \right| d\mathcal{H}^{n-1}(y) < \infty.$$

for U_i such that $d(x, U_i \cap \Gamma) < \rho$. Since $-\partial E / \partial \mathbf{n}_y(x - y)$ is invariant under translations and rotations, we can write $-\partial E / \partial \mathbf{n}_y(x - y)$ in the local coordinate. Let U_i be such that

$d(x, U_i \cap \Gamma) < \rho$ and denote h_{z_i} by h_i for simplicity. Let us observe that

$$-\mathbf{n}(y', h_i(y')) = (-\nabla' h_i(y'), 1) / \omega_i(y')$$

with $\omega_i(y') = (1 + |\nabla' h_i(y')|^2)^{1/2}$, where ∇' is the gradient in y' variables. This implies that

$$-n\alpha(n) \frac{\partial E}{\partial \mathbf{n}_y}(x - y) = \frac{\sigma_i(y')}{\omega_i(y') \left(|x' - y'|^2 + (x_n - h_i(y'))^2 \right)^{n/2}}$$

for $y \in \Gamma_i$ with

$$\sigma_i(y') := -\nabla' h_i(y) \cdot (x' - y') + (x_n - h_i(y')) \quad \text{where } x_n > h_i(x'), \quad x' \in B_{3\rho}(0').$$

We set

$$K_i(x', y', x_n) = \frac{\sigma_i(y')}{\left(|x' - y'|^2 + (x_n - h_i(y'))^2 \right)^{n/2}}.$$

By the Taylor expansion

$$h_i(x') = h_i(y') + \nabla' h_i(y') \cdot (x' - y') + r_i(x', y')$$

with

$$r_i(x', y') = (x' - y')^T \cdot \int_0^1 (1 - \theta) \nabla'^2 h_i(\theta x' + (1 - \theta)y') d\theta \cdot (x' - y'),$$

we obtain

$$\sigma_i(y') = x_n - h_i(x') + r_i(x', y')$$

with an estimate

$$|r_i(x', y')| \leq \|\nabla'^2 h_i\|_{L^\infty(B_{3\rho}(0'))} |x' - y'|^2. \quad (18)$$

We decompose K_i into a leading term and a remainder term

$$K_i(x', y', x_n) = K_0^i(x', y', x_n) + R_i(x', y', x_n)$$

with

$$K_0^i(x', y', x_n) := \frac{x_n - h_i(x')}{\left(|x' - y'|^2 + (x_n - h_i(y'))^2 \right)^{n/2}}$$

$$R_i(x', y', x_n) := \frac{r_i(x, y)}{\left(|x' - y'|^2 + (x_n - h_i(y'))^2 \right)^{n/2}}.$$

The term K_0^i is very singular but it is positive. The term R_i is estimated as

$$|R_i(x', y', x_n)| \leq \|\nabla'^2 h_i\|_{L^\infty(B_{3\rho}(0'))} |x' - y'|^{2-n}$$

by the estimate (18). Hence,

$$\int_{\Gamma \cap U_i} \left| \frac{R_i(x', y', x_n)}{\omega_i(y')} \right| d\mathcal{H}^{n-1}(y) \leq C \int_{B_\rho(0')} \frac{1}{|x' - y'|^{n-2}} dy' \leq C\rho$$

with C independent of ρ and i . By (i), we observe that

$$\begin{aligned} n\alpha(n) &= \sum_{i: d(x, U_i \cap \Gamma) < \rho} \int_{B_\rho(0')} \frac{K_i(x', y', x_n)}{\omega_i(y')} dy' \\ &\quad - n\alpha(n) \sum_{j: d(x, U_j \cap \Gamma) \geq \rho} \int_{U_j \cap \Gamma} \frac{\partial E}{\partial \mathbf{n}_y}(x - y) d\mathcal{H}^{n-1}(y). \end{aligned}$$

Since K_0^i is positive for any i such that $d(x, U_i \cap \Gamma) < \rho$,

$$\sum_{i: d(x, U_i \cap \Gamma) < \rho} \int_{B_\rho(0')} \frac{K_0^i(x', y', x_n)}{\omega_i(y')} dy' \leq n\alpha(n) \cdot \left(1 + \frac{m \cdot C \cdot S(\Gamma)}{\rho^{n-1}}\right) + m \cdot C \cdot \rho$$

where $S(\Gamma)$ denotes the surface area of Γ , which is bounded. Thus, the estimate

$$\int_{U_i \cap \Gamma} \left| \frac{\partial E}{\partial \mathbf{n}_y}(x - y) \right| d\mathcal{H}^{n-1}(y) \leq \frac{1}{n\alpha(n)} \int_{B_\rho(0')} \frac{K_0^i + |R_i|}{\omega_i(y')} dy' < \infty$$

holds for any U_i such that $d(x, U_i \cap \Gamma) < \rho$. The proof of (ii) is now complete. $\square$

Based on Lemma 16, we are able to prove Lemma 15 (ii).

Lemma 15 (ii). We decompose

$$\begin{aligned} \nabla d(x) \cdot \nabla (E * (\delta_\Gamma \otimes g))(x) &= \int_\Gamma (\nabla d(x) - \nabla d(y)) \cdot \nabla E(x - y) g(y) d\mathcal{H}^{n-1}(y) \\ &\quad + \int_\Gamma \frac{\partial E}{\partial \mathbf{n}_y}(x - y) g(y) d\mathcal{H}^{n-1}(y) = I_1 + I_2. \end{aligned}$$

Let $x \in \Gamma_{\rho_0}^{\mathbf{R}^n}$ and πx be the projection of x on Γ . For $y \in U_{\rho_0}(\pi x)$, there exists a constant L' , independent of x and y , such that

$$|\nabla d(x) - \nabla d(y)| \leq L'|x - y|.$$

For $y \in \Gamma_{\rho_0}^{\mathbf{R}^n} \setminus U_{\rho_0}(\pi x)$, we have that $|x - y| \geq \frac{\rho_0}{2}$. Since $\overline{\Gamma_{\rho_0/2}^{\mathbf{R}^n}}$ is compact in $\mathbf{R}^n$, by considering a finite subcover of $\cup_{z \in \Gamma} U_{\rho_0}(z)$ we are able to show that there exists $M > 0$ such that the estimate

$$|\nabla d(x) - \nabla d(y)| \leq M|x - y|$$

holds for any $x, y \in \Gamma_{\rho_0}^{\mathbf{R}^n}$. Thus,

$$H(x, y) = (\nabla d(x) - \nabla d(y)) \cdot \nabla E(x - y)$$

is estimated as

$$|H(x, y)| \leq \frac{M}{|x - y|^{n-2}}$$

in $\Gamma_{\rho_0}^{\mathbf{R}^n} \times \Gamma_{\rho_0}^{\mathbf{R}^n}$. We observe that

$$\begin{aligned} \sup_{x \in \Gamma_{\rho_0}^{\mathbf{R}^n} \cap \Omega} |I_1(x)| &\leq \sup_{x \in \Gamma_{\rho_0}^{\mathbf{R}^n} \cap \Omega} \int_{\Gamma} H(x, y) d\mathcal{H}^{n-1}(y) \|g\|_{L^\infty(\Gamma)} \\ &\leq M \sup_{x \in \Gamma_{\rho_0}^{\mathbf{R}^n} \cap \Omega} \int_{\Gamma} \frac{d\mathcal{H}^{n-1}(y)}{|x - y|^{n-2}} \|g\|_{L^\infty(\Gamma)}. \end{aligned}$$

Since

$$\sup_{x \in \Gamma_{\rho_0}^{\mathbf{R}^n} \cap \Omega} |I_2(x)| \leq \sup_{x \in \Gamma_{\rho_0}^{\mathbf{R}^n} \cap \Omega} \int_{\Gamma} \left| \frac{\partial E}{\partial \mathbf{n}_y}(x - y) \right| d\mathcal{H}^{n-1}(y) \|g\|_{L^\infty(\Gamma)},$$

Lemma 16 (ii) now yields (17). The proof is now complete. $\square$

We wonder whether the tangential component of $\nabla E * (\delta_\Gamma \otimes g)$ satisfies the same estimate. Unfortunately, the estimate

$$\|\nabla (E * (\delta_\Gamma \otimes g))\|_{L^\infty(\Gamma_{\rho_0}^{\mathbf{R}^n} \cap \Omega)} \leq C\|g\|_{L^\infty(\Gamma)}$$

should not hold even if Γ is flat. Even weaker estimate

$$[\nabla (E * (\delta_\Gamma \otimes g))]_{b^\nu(\Gamma)} \leq C\|g\|_{L^\infty(\Gamma)}$$

should not hold in general.

To illustrate the problem, we consider the case that Γ is flat. We may assume $\Gamma = \partial \mathbf{R}_+^n$, $\mathbf{R}_+^n = \{x_n > 0\}$.

Lemma 17. *The estimate*

$$\|\partial_{x_n} (E * (\delta_\Gamma \otimes g))\|_{L^\infty(\mathbf{R}_+^n)} \leq \frac{1}{2}\|g\|_{L^\infty(\mathbf{R}^{n-1})}$$

holds for $g \in L^\infty(\mathbf{R}^{n-1})$.

Proof. This is because $-\partial_{x_n}(E * (\delta_\Gamma \otimes g))$ is the half of the Poisson integral, i.e.,

$$-\partial_{x_n}(E * (\delta_\Gamma \otimes g))(x) = \frac{1}{2} \int_{\mathbf{R}^{n-1}} P_{x_n}(x' - y') g(y') dy',$$

where P_{x_n} denotes the Poisson kernel. Thus the desired L^∞ estimate follows from the maximum principle of the Dirichlet problem for the Laplacian or from the property that $\int_{\mathbf{R}^{n-1}} P_{x_n}(x') dx' = 1$ and $P_{x_n} \geq 0$. $\square$

Theorem 18. *There is a bounded sequence of smooth functions $\{g_\ell\}_{\ell \in \mathbf{N}} \subset L^\infty(\mathbf{R}^{n-1})$ such that*

$$\lim_{\ell \rightarrow \infty} [\partial_{x'}(E * (\delta_\Gamma \otimes g_\ell))]_{b^\nu} = \infty$$

for any $\nu > 0$.

Proof. If g is smooth, $E * (\delta_\Gamma \otimes g)$ is smooth up to the boundary. In this case, if $[\partial_{x'}(E * (\delta_\Gamma \otimes g))]_{b^\nu}$ is bounded by $C\|g\|_{L^\infty(\mathbf{R}^{n-1})}$, $\|\partial_{x'}(E * (\delta_\Gamma \otimes g))\|_{L^\infty(\Gamma)}$ is also bounded by $c_0 C\|g\|_{L^\infty(\mathbf{R}^{n-1})}$ with a constant c_0 depending only on n since the mean value over r -ball around x converges to its value at x as $r \rightarrow 0$.

We consider the Neumann problem

$$\begin{aligned} \Delta u &= 0 \quad \text{in } \mathbf{R}_+^n, \\ \frac{\partial u}{\partial \mathbf{n}} &= g \quad \text{on } \Gamma = \partial \mathbf{R}_+^n. \end{aligned}$$

By using the tangential Fourier transform, we see that

$$u(x, t) = \Lambda^{-1} \exp(-x_n \Lambda) g$$

where $\Lambda = (-\Delta')^{1/2}$. If $\|\nabla' u\|_{L^\infty(\Gamma)} \leq C\|g\|_{L^\infty(\mathbf{R}^{n-1})}$ were true, sending $x_n > 0$ to zero would imply L^∞ boundedness of the Riesz operator $\nabla' \Lambda^{-1}$, which is absurd.

The operator $E * (\delta_\Gamma \otimes g)$ is the half of the solution operator of the Neumann problem, so L^∞ bound for $\nabla' E * (\delta_\Gamma \otimes g)$ should not hold even if it is restricted to smooth functions. $\square$

Corollary 19. *Assume that $\Omega = \mathbf{R}_+^n$. Let $v \mapsto \nabla q$ be the Helmholtz projection to a gradient field. Then, this projection is unbounded from $(L^\infty(\Omega))^n$ to $(BMO_b^{\mu, \nu}(\Omega))^n$ for any $\mu, \nu > 0$.*

Proof. We consider

$$v = (0, \dots, 0, v_n(x'))$$

with $v_n \in L^\infty(\mathbf{R}^{n-1})$. This evidently solves $\operatorname{div} v = 0$. The normal trace equals $-v_n(x')$. If

$$[\nabla q]_{b^\nu} \leq C \|v_n\|_{L^\infty(\mathbf{R}^{n-1})}$$

for all $v_n \in L^\infty(\mathbf{R}^{n-1})$ with C independent of v , then this would contradict Theorem 18. $\square$

Acknowledgement

The first author was partly supported by the Japan Society for the Promotion of Science through grants No. 19H00639 (Kiban A), No. 18H05323 (Kaitaku), No. 17H01091 (Kiban A). The second author is partly supported by the Mizuho international foundation through foreign students scholarship program.

References

- [1] M. Bolkart and Y. Giga, On L^∞ - BMO estimates for derivatives of the Stokes semigroup, *Math. Z.*, 284, 1163–1183 (2016)
- [2] M. Bolkart, Y. Giga, T.-H. Miura, T. Suzuki, Y. Tsutsui, On analyticity of the L^p -Stokes semigroup for some non-Helmholtz domains, *Math. Nachr.*, 290, 2524–2546 (2017)
- [3] M. Bolkart, Y. Giga and T. Suzuki, Analyticity of the Stokes semigroup in BMO -type spaces, *J. Math. Soc. Japan*, 70, 153–177 (2018)
- [4] M. Bolkart, Y. Giga, T. Suzuki, Y. Tsutsui, Equivalence of BMO -type norms with applications to the heat and Stokes semigroup, *Potential Anal.*, 49, 105–130 (2018)
- [5] R. Farwig, H. Kozono and H. Sohr, An L^q -approach to Stokes and Navier-Stokes equations in general domains, *Acta Math.*, 195, 21–53 (2005)
- [6] R. Farwig, H. Kozono and H. Sohr, On the Helmholtz decomposition in general unbounded domains, *Arch. Math. (Basel)*, 88, 239–248 (2007)
- [7] C. Fefferman and E. M. Stein, H^p spaces of several variables, *Acta Math.*, 129, 137–193 (1972)
- [8] D. Fujiwara and H. Morimoto, An L_r -theorem of the Helmholtz decomposition of vector fields, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 24, 685–700 (1977)
- [9] G. P. Galdi, An introduction to the mathematical theory of the Navier-Stokes equations. Steady-state problems. Second edition, Springer Monographs in Mathematics, xiv+1018 pp. Springer, New York (2011)

- [10] Y. Giga and Z. Gu, On the Helmholtz decompositions of vector fields of bounded mean oscillation and in real Hardy spaces over the half space, *Adv. Math. Sci. Appl.*, 29, 87–128 (2020)
- [11] Y. Giga and Z. Gu, Normal trace for vector fields of bounded mean oscillation, *arXiv:2011.12029* (2020)
- [12] Y. Giga, Z. Gu and P.-Y. Hsu, Continuous alignment of vorticity direction prevents the blow-up of the Navier-Stokes flow under the no-slip boundary condition, *Nonlinear Anal.*, 189, 111579 (2019)
- [13] D. Gilbarg, N. S. Trudinger, Elliptic partial differential equations of second order, Second edition, *Grundlehren der Mathematischen Wissenschaften*, 224. Springer-Verlag, Berlin (1983)
- [14] L. Grafakos, Classical Fourier Analysis, Third edition, *Graduate Texts in Mathematics*, 249. Springer-Verlag, New York (2014)
- [15] L. Grafakos, Modern Fourier Analysis, Third edition, *Graduate Texts in Mathematics*, 250. Springer-Verlag, New York (2014)
- [16] R. Hanks, Interpolation by the real method between BMO , $L^\alpha(0 < \alpha < \infty)$ and $H^\alpha(0 < \alpha < \infty)$, *Indiana Univ. Math. J.*, 26, 679–689 (1977)
- [17] P. W. Jones, Extension theorems for BMO , *Indiana Univ. Math. J.*, 29, 41–66 (1980)
- [18] H. Kozono and H. Wadade, Remarks on Gagliardo-Nirenberg type inequality with critical Sobolev space and BMO , *Math. Z.*, 259, 935–950 (2008)
- [19] S. G. Krantz, H. R. Parks, The implicit function theorem. History, theory, and applications, Reprint of the 2003 edition. *Modern Birkhäuser Classics*, Birkhäuser/Springer, New York (2013)
- [20] J.-L. Lions and E. Magenes, Non-homogeneous boundary value problems and applications, Vol. I, xvi+357 pp. Springer-Verlag, New York-Heidelberg (1972)
- [21] A. Miyachi, H^p spaces over open subsets of R^n , *Studia Math.*, 95, 205–228 (1990)
- [22] Y. Sawano, Theory of Besov spaces. *Developments in Mathematics*, 56, Springer, Singapore (2018)
- [23] C. G. Simader and H. Sohr, A new approach to the Helmholtz decomposition and the Neumann problem in L^q -spaces for bounded and exterior domains, *Mathematical problems*

relating to the Navier-Stokes equation, 1–35, Ser. Adv. Math. Appl. Sci., 11, World Sci. Publ., River Edge, NJ, 1992