



Title	Enzootic nasal tumor virus type 2 envelope of goats acts as a retroviral oncogene in cell transformation
Author(s)	Maeda, Naoyoshi; Inoshima, Yasuo; De las Heras, Marcelo et al.
Citation	Virus Genes, 57(1), 50-59 https://doi.org/10.1007/s11262-020-01808-7
Issue Date	2020-11-05
Doc URL	https://hdl.handle.net/2115/83203
Rights	This is a post-peer-review, pre-copyedit version of an article published in Virus Genes. The final authenticated version is available online at: http://dx.doi.org/10.1007/s11262-020-01808-7 .
Type	journal article
File Information	Maeda et al., Virus Genes 2020.pdf



Virus Genes

Original article

Title: Enzootic nasal tumor virus type 2 envelope of goats acts as a retroviral oncogene in cell transformation

Authors: Naoyoshi Maeda^{a,*}, Yasuo Inoshima^c, Marcelo De las Heras^d, and Katsumi Maenaka^{a,b}

Affiliations: ^aCenter for Research and Education on Drug Discovery, and ^bLaboratory of Biomolecular Science, Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, JAPAN, ^cLaboratory of Food and Environmental Hygiene, Cooperative Department of Veterinary Medicine, Gifu University, 1-1 Yanagido, Gifu 501-1193, JAPAN, and ^dDepartment of Animal Pathology, Zaragoza University, Zaragoza, SPAIN

*Corresponding author: Naoyoshi Maeda, Ph. D., Center for Research and Education on Drug Discovery, Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, JAPAN, Tel: +81-11-706-3752/Fax: +81-11-706-4983, E-mail: nmaeda@pharm.hokudai.ac.jp

Keywords: Envelope, Enzootic nasal adenocarcinoma, Enzootic nasal tumor virus, Jaagsiekte sheep retrovirus, Oncogenic transformation, Signal transduction

Text: 5,457 words (including figure legends)

Abstract: 213 words

25 70 references

26 6 figures

Abstract

Enzootic nasal tumor virus type 1 (ENTV-1) (ovine nasal tumor virus) and ENTV-2 (caprine nasal tumor virus) are known to be causative agents of enzootic nasal adenocarcinoma (ENA) in sheep and goats, respectively. Although the nucleotide and amino acid sequences of ENTV-1 and ENTV-2 are quite similar, they are recognized as phylogenetically distinct viruses. The envelope protein of ENTV-1 functions as an oncoprotein in the *in vitro* transformation of epithelial cells and fibroblasts. Thus, it is the primary determinant of *in vivo* tumorigenesis in ENA. As per our knowledge, no previous studies have reported in detail the role of ENTV-2 in ENA tumorigenesis. Here, in order to investigate the molecular mechanism of caprine ENA oncogenesis by ENTV-2, we have attempted to identify the transforming potential of ENTV-2 envelope, and investigated the activation of cell signaling pathways in oncogenic transformation. Our findings confirmed that ENTV-2 envelope was capable of inducing oncogenic transformation of rat cell lines *in vitro*. Further, we found that MAPK, Akt, and p38 were constitutively activated in ENTV-2 envelope-transformed clone cells. In addition, inhibitor experiments revealed that MEK-MAPK and PI3K-Akt signaling pathways are involved in the ENTV-2 envelope-induced cell transformation. These data indicate that ENTV-2 envelope could induce oncogenic transformation by signaling pathways that are also utilized by ENTV-1 envelope.

Introduction

Enzootic nasal adenocarcinoma (ENA), previously known as enzootic intranasal tumor, is a contagious neoplasia in sheep and goats [1]. It has been reported worldwide [2-9], with several cases being reported in China in recent times [10-13]. The etiological agent of ENA has been identified as retroviruses [14-17]. Experimental infection was successful in inducing ENA by cell-free tumor filtrates in sheep [18-20], and by concentrated ENA nasal exudate in goats [21]. The sequences of ovine nasal tumor virus (ENTV-1 in sheep) and caprine nasal tumor virus (ENTV-2 in goats) were identified as similar but phylogenetically distinct viruses [22-26]. The target tissues in ENA were the secretory epithelial cells of the ethmoid turbinate of the nasal cavity; however, the tissue distribution of the two viruses was different; while ENTV-1 was detected in tumors, lymph nodes, kidneys, and lungs, ENTV-2 was additionally detected in peripheral blood mononuclear cells (PBMCs) and bone marrow [26].

Both ENTV-1 and ENTV-2 are classified as simple ovine β -retroviruses. This category also includes jaagsiekte sheep retrovirus (JSRV), which is an etiological agent for naturally occurring and experimentally induced ovine pulmonary adenocarcinoma (OPA) in sheep [27]. These three retroviruses consist of 5'-long terminal repeat (LTR)-*gag-pro-pol-env*-3'-LTR, without any viral oncogenes [23, 26, 27]. However, they encode another open reading frame x, designated as *orf-x*, which was initially in focus for its role in oncogenesis by JSRV [28]; but it was found that it was not required for the *in vitro* transformation and induction of OPA *in vivo* [29, 30]. In the classical transformation assay, the full-length envelope glycoprotein of ENTV-1, as well as that of JSRV, was identified as an oncoprotein contributing to the transformation of fibroblasts and epithelial cells *in vitro* [29, 31-33]. The JSRV envelope itself induced lung tumors not only in sheep [34] but also in mice *in vivo* [35-38]. Interestingly, ENTV-1 envelope also induced lung tumors in mice [39]. In addition to the envelopes, the LTR region is considered to be an important factor in determining their tissue tropism [37, 40]. A recent study has further revealed that the expression of a glycosylphosphatidylinositol-anchored cell surface protein hyaluronidase-2 (Hyal-2), an entry receptor

for JSRV and ENTV [31], influences the entry of ENTV into nasal target cells, but not JSRV [41].

The JSRV envelope transmembrane (TM) domain is required for the transformation of fibroblasts [42]. Further mutational analysis revealed that the cytoplasmic tail (CT) region of JSRV envelope TM, in particular, is required for transformation *in vitro*. Certain tyrosine residues of the CT region, including Tyr590 of JSRV and Tyr592 and Tyr596 of ENTV-1, were initially regarded as critical for transformation in the context of signal transduction [43-47]; phosphatidylinositol 3-kinase (PI3K) to Akt and mitogen extracellular regulated kinase (MEK) to mitogen-activated protein kinase (MAPK) pathways are involved in transformation *in vitro* [43-49]. Besides the molecules involved in the above signaling pathways, other molecules including Ras, Rac1, p38, Hsp90, and Sprouty2 have also been reported to play a role in cell transformation [48, 50, 51]. Signaling pathways utilized by ENTV-1 resemble those used by JSRV *in vitro* and *in vivo* [52, 53].

While a lot of research has been conducted on cell transformation induced by ENTV-1 and JSRV, no previous studies have reported on the molecular mechanism of caprine ENA oncogenesis induced by ENTV-2. The present study aims to address this by focusing on the identification of the transforming potential of ENTV-2 envelope *in vitro*, and further investigation on the activation of signaling pathways associated with oncogenic transformation.

Materials and methods

Constructs

ENA tumors isolated from goats were collected and embedded in paraffin. Integrated genomic ENTV-2 genes were recovered from a paraffin block using TaKaRa DEXPAT (TaKaRa Bio, Shiga, Japan). PCR was performed using the recovered genomes as templates, forward and reverse primers designed according to the ENTV-2 sequence (GenBank accession number: AY197548) [26], dNTP mixture, *TaKaRa Ex Taq* DNA polymerase, and *TaKaRa Ex Taq* Buffer (TaKaRa Bio). PCR conditions were as follows: 95°C for 2 min,

30 cycles of 95°C for 30 sec, 45°C for 30 sec, and 68°C for 1 min. The PCR products were purified with a Gel Extraction Kit and PCR Purification Kit (QIAGEN, CA, USA), cloned into T-Vector pMD20 (TaKaRa Bio), and sequenced. Each PCR product was ligated using a DNA Ligation Kit (Mighty Mix) (TaKaRa Bio), and inserted into the HA-tagged JSRV envelope expression plasmid Δ GP(JSRV-HA) [48], designated as Δ GP(ENTV-2). New ENTV-2 sequence has been deposited as NAOM-HU3118124 strain in DNA Data Bank of Japan with accession number LC570918. The JSRV envelope expression plasmid was provided by Dr. Hung Fan (University of California, Irvine, CA, USA). The ENTV-2 envelope expression construct with 6 histidines (His)-tagged version was also developed, designated as Δ GP(ENTV-2-His).

Cell culture

Rat embryonic fibroblasts 208F cells and rat kidney epithelial RK3E cells were purchased from the European Collection of Authenticated Cell Cultures and the American Type Culture Collection, respectively. The two cell lines, along with human kidney epithelial 293T cells, were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS), penicillin (100 units/mL), and streptomycin (100 μ g/mL).

Transformation assay and establishment of transformed clone cells

208F and RK3E cells were seeded at 4×10^5 cells/plate in 6 cm dishes, 24 h prior to transfection. Five micrograms of Δ GP(ENTV-2), Δ GP(ENTV-2-His), or Δ GP(JSRV) were transfected into the cells using XtremeGENE HP DNA Transfection Reagent (Roche, Mannheim, Germany). To inhibit MAPK, PI3K, and p38 signaling pathways, cells after transfection were treated with inhibitors, PD98059, LY294002, and SB203580 (Calbiochem, CA, USA), respectively, or DMSO (WAKO, Osaka, Japan) as a control vehicle. Transformed foci were counted on days 11-20 after transfection. To establish transformed clone cells, Δ GP(ENTV-2-His) was transfected into 208F and RK3E cells. Transformed foci were picked up 3-4 weeks

after transfection, and single clone cells were isolated by a limiting dilution. The cells were lysed with lysis buffer (20 mM Tris (pH 7.4), 150 mM NaCl, 10% glycerol, 1% Triton X-100) supplemented with Complete protease inhibitor cocktail (Roche).

Treatment with epidermal growth factor (EGF)

Parental and transformed 208F cells were seeded at 5×10^5 cells/plate in 6 cm dishes. They were serum-starved for 20-24 h, and the parental 208F cells were subsequently stimulated with 100 ng/mL recombinant human EGF (BioAcademia, Osaka, Japan) for 30 min. The cells were lysed with lysis buffer supplemented with Complete protease inhibitor cocktail.

Immunoblot

The cell lysate was mixed with 5X sodium dodecyl sulfate (SDS) loading buffer (50 mM Tris (pH 6.8), 10% glycerol, 1% SDS, and 0.05% bromophenol blue) with or without reducing agent 1% β -mercaptoethanol, and boiled at 95°C for 5 min. The total proteins were separated on SDS polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto a nitrocellulose membrane (GE Healthcare, USA). The membranes were incubated with Tris-buffered saline with Tween 20 (TBS-T) having 5% nonfat dry milk or bovine serum albumin at room temperature for 1 h. The membranes were incubated with specific antibodies overnight at 4°C. Monoclonal antibodies (mAbs) against p44/42 MAPK (Erk1/2), phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204), Akt (pan), phospho-Akt (Ser473), p38 MAPK, phospho-p38 MAPK (Thr180/Tyr182) (Cell Signaling, MA, USA), and His-tag (MBL, Nagoya, Japan) were used as primary antibodies. After washing the membranes three times with TBS-T, the membranes were incubated with anti-mouse or anti-rabbit IgG labeled with Alexa Fluor 680 (Thermo Fisher Scientific, MA, USA) at room temperature for 1 h. After washing the membranes three times with TBS-T, signals were detected using an Odyssey CLx Infrared Imaging System (LI-COR Biosciences, NE, USA).

Immunohistochemistry

Naturally occurring ENA tumors and normal goat ethmoid turbinate were embedded in paraffin. For antigen retrieval, the sections were microwaved at 95°C for 5 min. in 10 mM citrate buffer (pH 6.0). Endogenous peroxidase was quenched with 0.3% hydrogen peroxidase in methanol for 30 min. The sliced sections were stained with hematoxylin and eosin (H&E), anti-phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) mAb, and anti-phospho-Akt (Ser473) mAb. Immunohistochemistry was performed by using a commercially available EnVisionTM+ (Dako, CA, USA) according to instructions by the manufacturer.

Statistical analysis

Statistically significant differences were calculated using Student's *t*-test and are indicated as *p* values. Differences of *p* < 0.05 were considered to be statistically significant.

Results

Transformation of rat cells by ENTV-2 envelope

In order to make a construct for the expression of full-length ENTV-2 envelope, we performed PCR to amplify proviral ENTV-2 *env* DNA derived from the ENA tumor genome. The nucleotide sequence of the recovered ENTV-2 *env* gene had 98% identity (1,805 out of 1,848 nucleotides) with AY197548, which had been previously identified in Spain [26]. The deduced amino acid sequence of the new ENTV-2 was identical to that of the AY197548 strain, though Ser at position 111 was mutated to Arg. For efficient expression in mammalian cells, the JSRV *env* gene in the expression plasmid ΔGP(JSRV-HA) was replaced with the full-length ENTV-2 *env* gene for its expression under the cytomegalovirus (CMV) promoter, designated as ΔGP(ENTV-2) (Fig. 1). In addition, for the easy detection of ENTV-2 envelope, an expression construct was developed for ENTV-2 envelope with His-tag (ΔGP(ENTV-2-His)) (Fig. 1).

To investigate the transforming potential of the ENTV-2 envelope, we transfected rat fibroblast 208F cells with the ENTV-2 envelope expression plasmid Δ GP(ENTV-2), JSRV envelope expression plasmid Δ GP(JSRV) or pcDNA3.1(-), with the latter two plasmids as positive and negative plasmid controls, respectively. In the transformation assay, we found that the ENTV-2 and JSRV envelopes induced foci of transformed cells in 208F cells (Fig. 2a and c), as did the ENTV-2 envelope with His-tagged construct Δ GP(ENTV-2-His) (Fig. 2b). Moreover, we found that the ENTV-2 envelope induced foci of transformed cells in rat epithelia RK3E cells (Fig. 2e-g). The negative plasmid control did not induce any foci of transformed cells in either cell line (Fig. 2d and h).

Signaling pathways in ENTV-2 envelope-induced transformation *in vitro*

In order to investigate the molecular mechanisms involved in ENTV-2 envelope-induced transformation, we utilized the chemical inhibitors PD98059 for MEK, LY294002 for PI3K, and SB203580 for p38 in the transformation assay. The number of foci of transformed cells treated with MEK and PI3K inhibitors at day 20 after transfection was lower compared to that in cells treated with DMSO as a control vehicle (Fig. 3a). Moreover, the size of each focus on cells treated with MEK and PI3K inhibitors was smaller than that in cells treated with DMSO (Fig. 3b-j). These results suggested that MEK and PI3K pathways are involved in the ENTV-2 envelope-induced transformation. On the other hand, the number of foci of transformed cells treated with p38 inhibitor at day 13 after transfection showed an increase, compared to that in cells treated with DMSO (Fig. 4a). The foci on cells treated with p38 inhibitor were larger in size, compared to those in cells treated with DMSO (Fig. 4b-g). Similar results were also observed in the experiments conducted using RK3E cells (data not shown). These results suggest that the ENTV-2 envelope triggers MEK and PI3K signaling pathways to induce transformation, while the p38 signaling pathway may negatively regulate the transformation.

To further investigate the signaling pathways involved in ENTV-2 envelope-induced

transformation, we established 208F clone cells transformed with ENTV-2 envelope having a His-tag (Fig. 5a). The transformed cells (Fig. 5a) showed differences in morphology compared to the parental 208F cells (Fig. 5b). Full-length and TM domain of ENTV-2 envelope were detected by western blot analysis (Fig. 5c). In the ENTV-2 envelope-transformed clone cells, we found constitutive activation of MAPK (also called p44/42 or Erk1/2), Akt, and p38 (Fig. 5d-f).

MAPK and Akt activation in naturally occurring ENA tumors *in vivo*

Immunohistochemical staining of naturally occurring caprine ENA tumors, from which the ENTV-2 *env* gene was recovered, showed the typical histological features of adenocarcinoma (Fig. 6a and d). In serial sections of the same sample, we detected phosphorylation of p44/42 MAPK at Thr202/Tyr204 (Fig. 6b and e) and Akt at Ser 473 (Fig. 6c and f). Normal goat ethmoid turbinate did not show any phosphorylation of these molecules (Fig. 6g-l). These results, together with those of the *in vitro* experiments, strongly suggest that the MEK-MAPK and PI3K-Akt signaling pathways are involved in ENTV-2 envelope-induced ENA oncogenesis *in vivo*.

Discussion

ENA and OPA caused by ENTV-1 and JSRV, respectively, are interesting and unusual examples of neoplasms induced by virus envelopes having transforming potential [54]. While the mechanisms of ENTV-1 and JSRV transformation have been widely studied, there's been a lack of in-depth investigation into caprine ENA oncogenesis by ENTV-2. Thus, in order to understand the molecular mechanisms of caprine ENA, this study focused on identifying the transforming potential of ENTV-2 envelope *in vitro*. Our findings confirmed that ENTV-2 envelope was indeed capable of inducing oncogenic transformation of rat cell lines *in vitro*. Further, we found that MAPK, Akt, and p38 were constitutively activated in ENTV-2 envelope-transformed clone cells. In addition, inhibitor experiments revealed that MEK-MAPK and

PI3K-Akt signaling pathways are involved in the ENTV-2 envelope-induced cell transformation. Collectively, these results revealed that the transforming potential of ENTV-2 envelope seems to be quite similar to that of JSRV *in vitro*. In previous studies, activation of MAPK phosphorylated on Thr202/Tyr204 was observed not only in naturally occurring ENA, but also in naturally occurring OPA as well as experimentally induced OPA *in vivo* [48, 53]. However, while the staining of ENA tumors from sheep and goats with anti-phosphorylated Akt mAb was positive, the staining of OPA tumor section with that was not positive in all cases, although phosphorylation of Akt at Ser473 was observed in JSRV envelope-transformed cells *in vitro* [44]. This led us to postulate that although the signaling pathways leading to oncogenic transformation by the ENTV-1 and JSRV envelopes shared an overall similarity, there were slight differences in the mechanism. Additionally, we found that the caprine ENA tumor used in this study also exhibited phosphorylation of MAPK and Akt. Taken together, we concluded that ENTV-2 envelope could induce oncogenic transformation by signaling pathways that are also utilized by the ENTV-1 envelope.

The molecular mechanism through which the ENTVs and JSRV envelopes activate these signaling pathways has not yet been fully understood. In particular, the elucidation of the cellular molecules that interact with these envelopes should be a key factor towards understanding their oncogenic potential. One of the interacting partners identified as an entry receptor for JSRV as well as ENTV-1 is Hyal-2, which was initially considered to be a tumor suppressor located at human chromosome 3p21.3 [55]. However, the involvement of Hyal-2 in transformation is controversial; mouse fibroblast NIH-3T3 cells are successfully transformed, although Hyal-2 of the murine homolog does not function as a receptor for the JSRV and ENTV-1 envelopes [56]. Two studies independently reported that RalA binding protein 1 (RALBP1) and zinc finger protein 111 (Zfp111) directly interacted with the JSRV envelope CT region [57, 58]. RALBP1 was involved in JSRV transformation, via the formation of a complex with CDC42 to induce the activation of downstream effectors such as the mammalian target of rapamycin and p70S6 kinase in canine kidney epithelial MDCK cells [57]. On the other hand, Zfp111 was found to interact with a nuclear form of JSRV

envelope in rat fibroblast 208F cells, suggesting that JSRV transformation involves events in the nuclear region [58]. These studies indicated that JSRV envelope interacts with multiple cellular or nuclear proteins to induce cellular transformation *in vitro*. Nevertheless, the mechanism of activation of MAPK and other molecules, including p38, Ras, and Rac1, in the JSRV envelope-induced transformation, as well as their involvement in OPA oncogenesis, remains unclear. Further research is needed on the identification of proteins interacting with the ENTV-1 and ENTV-2 envelopes. The amino acid sequence of variable region 3 (VR3), especially in the CT region, in the ENTV and JSRV envelopes, is not conserved; thus, the proteins interacting with ENTVs may be different from those interacting with JSRV. This fact could help distinguish between the similar yet slightly different signaling pathways leading to oncogenic transformation by the ENTV and JSRV envelopes. In addition to the transforming potential of CT regions, they have also been reported to contribute to the infectivity and fusogenicity of viruses; thus, the CT regions play a critical role in exhibiting multiple functions in the viral life cycle [59-65]. It has been observed that the Moloney murine leukemia virus envelope CT region possesses an amphipathic α -helix that facilitates membrane fusion [66, 67]. Previous studies have predicted the presence of a putative amphipathic α -helical wheel structure in the CT region of JSRV envelope [47]; this possibility is backed by multiple structural analyses (Maeda *et al.*, unpublished observations). Further investigation is also needed to elucidate the structure-function and relationship of ENTV-2 envelope, especially the CT region, in ENA tumorigenesis.

So far, it is not yet unclear why ENTV-1 and ENTV-2 cause identical diseases but could be detected only in sheep and goats, respectively. JSRV could experimentally induce lung tumors in goats that differ from those induced in sheep, suggesting that there are several restricting factors which determine species specificity [68]. In addition to the main target cells, such as type II pneumocytes and club (Clara) cells, JSRV can infect other cell types, including undifferentiated cells in the respiratory tract [69], indicating that its envelope may be able to induce different neoplastic diseases in tissues other than those of the lungs. For instance, JSRV, but not ENTV-1, infection was detected in ovine ENA [70]. Thus, tissue

specificity is believed to be influenced by other viral factors apart from the envelope. The viral LTR could be a critical factor in determining retrovirus tropism. The JSRV LTR is preferentially active in lung epithelial cells [40]. Rosales *et al.* recently reported that nasal turbinate chondrocytes could be potential target cells for ENTV-1 infection *in vivo*, in cells in which ENTV-1 LTR is significantly active [41]. It is notable that the LTR sequence of ENTV-2, especially the U3 region, is highly divergent from that of ENTV-1 and JSRV [11, 26]. As mentioned above, the envelope VR3 sequence of ENTV-2 is also significantly different from that of ENTV-1 and JSRV. These facts collectively may account for the yet unexplained goat tropism of ENTV-2, which would be an interesting point of focus in future studies.

Acknowledgments

This research is partially supported by a Grant-in-Aid Scientific Research by the Japan Society for the Promotion of Science (JSPS) (to N.M. and K.M.), and by the Platform Project for Supporting in Drug Discovery and Life Science Research (Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS)) from Japan Agency for Medical Research and Development (AMED) under Grant Number JP19am0101093 (support number 1234) (to K.M.). This research was also the result of using research equipment shared in the Ministry of Education, Culture, Sports, Science and Technology (MEXT) Project for promoting public utilization of advanced research infrastructure (Program for supporting introduction of the new sharing system) under Grant Number JPMXS0420100119 (to K.M.).

Author Contributions

NM conceived and designed the experiments. MDLH and YI provided the materials. NM and MDLH carried out the experiments. NM wrote the manuscript. NM, YI, MDLH, and KM reviewed and edited the manuscript. NM, YI, MDLH, and KM read and approved the final version of the manuscript.

285 **Compliance with ethical standards**

286 **Conflict of Interest**

287 The authors declare no conflict of interest.

288

289 **Ethical approval**

290 All applicable international, national, and institutional guidelines for the care and use of animals were

291 followed.

292

293 **Informed consent**

294 Informed consent was obtained from all individual participants included in this study.

295

296

297 **References**

- 298 1. De las Heras M, Ortín A, Cousens C, Minguijón E, Sharp JM (2003) Enzootic nasal adenocarcinoma
299 of sheep and goats. *Curr Top Microbiol Immunol* 275:201-223.

- 300 2. McKinnon AO, Thorsen J, Hayes MA, Misener CR (1982) Enzootic nasal adenocarcinoma of sheep
301 in Canada. *Can Vet J* 23:88-94.

- 302 3. Vitellozzi G, Mughetti L, Palmarini M, Mandara MT, Mechelli L, Sharp JM, Manocchio I (1993)
303 Enzootic intranasal tumour of goats in Italy. *Zentralbl Veterinarmed B* 40:459-468.

- 304 4. Kane Y, Rosati S, Diop OM, Profiti M, Niang I, Kadja M, Kaboret YY, Alogninouwa T, Lena P
305 (2004) Enzootic nasal tumour virus demonstrated in sheep in Senegal by direct detection of provirus
306 from tumour DNA. *Vet Rec* 155:526-528.

- 307 5. Kawasako K, Okamoto M, Kurosawa T, Nakade T, Kirisawa R, Miyashou T, Komine M, Go T, Imazu
308 S, Takeuchi N, Tomonaga K, Ikuta K, Akihara Y, Shimoyama Y, Hirayama K, Taniyama H (2005)
309 Enzootic intranasal tumour virus infection in apparently healthy sheep in Japan. *Vet Rec* 157:118-120.

- 310 6. Svava T, Gombac M, Vrecl M, Juntos P, Kostanjsek R, Pogacnik A, Pogacnik M (2006) Enzootic
311 nasal adenocarcinoma of sheep in Slovenia. *J Vet Med A Physiol Pathol Clin Med* 53:26-29.

- 312 7. Kaufmann C, Brechbühl M, Oevermann A, Müller S, Zimmer K, Schönmann M, Leroux C, Weibel
313 D, Bertoni G (2008) Enzootic nasal adenocarcinoma in a dairy sheep flock. *Schweiz Arch Tierheilkd*
314 150:297-302.

- 315 8. Sid N, Belalmi NEH, Benhamza L, Ouhida S, Zebiri ME, Aydoğan A, Leroux C (2018) First case
316 report of enzootic nasal adenocarcinoma in "Ouled Djellal" ewe in Algeria. *Open Vet J* 8:9-12.

- 317 9. de Cecco BS, Lorenzett MP, Henker LC, Weber MN, Moséna ACS, Baumbach L, Canal CW,
318 Driemeier D, Pavarini SP, Sonne L (2019) Detection of enzootic nasal tumor virus (ENTV) in a sheep
319 flock in southern Brazil. *Trop Anim Health Prod* 51:2095-2098.
- 320 10. Yi G, Kaiyu W, Qigui Y, Zhongqiong Y, Yingdong Y, Defang C, Jinlu H (2010) Descriptive study of
321 enzootic nasal adenocarcinoma in goats in southwestern China. *Transbound Emerg Dis* 57:197-200.
- 322 11. He Y, Zhang Q, Wang J, Zhou M, Fu M, Xu X (2017) Full-length genome sequence analysis of
323 enzootic nasal tumor virus isolated from goats in China. *Virol J* 14:141.
- 324 12. Ye C, Huang Q, Chen T, Jiang J, Hou F, Xu D, Peng Y, Fang R, Chen J (2019) First detection and
325 genotypic analysis of goat enzootic nasal tumor virus 2 in Chongqing, China. *Arch Virol* 164:1647-
326 1650.
- 327 13. Zhai SL, Lv DH, Xu ZH, Yu JS, Wen XH, Zhang H, Chen QL, Jia CL, Zhou XR, Zhai Q, Li F, Woo
328 PCY, Lau SKP, Wang D, Wei WK (2019) A novel enzootic nasal tumor virus circulating in goats from
329 southern China. *Viruses* 11. pii: E956.
- 330 14. De las Heras M, García de Jalón JA, Balaguer L, Badiola JJ (1988) Retrovirus-like particles in
331 enzootic intranasal tumours in Spanish goats. *Vet Rec* 123:135.
- 332 15. De las Heras M, Sharp JM, Garcia de Jalon JA, Dewar P (1991) Enzootic nasal tumour of goats:
333 demonstration of a type D-related retrovirus in nasal fluids and tumours. *J Gen Virol* 72:2533-2535.
- 334 16. De las Heras M, Garcia de Jalon JA, Sharp JM (1991) Pathology of enzootic intranasal tumor in thirty-
335 eight goats. *Vet Pathol* 28:474-481.

- 336 17. De las Heras M, Sharp JM, Ferrer LM, García de Jalón JA, Cebrian LM (1993) Evidence for a type
337 D-like retrovirus in enzootic nasal tumour of sheep. Vet Rec 132:441.
- 338 18. Cohrs, P (1953) Infektiose Adenopapillome der Reichschleimhaut beim schaf. Berl Munch Tierarztl
339 Wochenschr 66:225-228.
- 340 19. Walsh SR, Linnerth-Petrik NM, Yu DL, Foster RA, Menzies PI, Diaz-Méndez A, Chalmers HJ,
341 Wootton SK (2013) Experimental transmission of enzootic nasal adenocarcinoma in sheep. Vet Res
342 44:66.
- 343 20. Walsh SR, Stinson KJ, Wootton SK (2016) Seroconversion of sheep experimentally infected with
344 enzootic nasal tumor virus. BMC Res Notes 9:15.
- 345 21. De las Heras M, García de Jalón JA, Minguijón E, Gray EW, Dewar P, Sharp JM (1995) Experimental
346 transmission of enzootic intranasal tumors of goats. Vet Pathol 32:19-23.
- 347 22. Cousens C, Minguijon E, Garcia M, Ferrer LM, Dalziel RG, Palmarini M, De las Heras M, Sharp JM
348 (1996) PCR-based detection and partial characterization of a retrovirus associated with contagious
349 intranasal tumors of sheep and goats. J Virol 70:7580-7583.
- 350 23. Cousens C, Minguijon E, Dalziel RG, Ortin A, Garcia M, Park J, Gonzalez L, Sharp JM, De las Heras
351 M (1999) Complete sequence of enzootic nasal tumor virus, a retrovirus associated with transmissible
352 intranasal tumors of sheep. J Virol 73:3986-3993.

- 353 24. Walsh SR, Linnerth-Petrik NM, Laporte AN, Menzies PI, Foster RA, Wootton SK (2010) Full-length
354 genome sequence analysis of enzootic nasal tumor virus reveals an unusually high degree of genetic
355 stability. *Virus Res* 151:74-87.
- 356 25. Walsh SR, Gerpe MC, Wootton SK (2016) Construction of a molecular clone of ovine enzootic nasal
357 tumor virus. *Virol J* 13:209.
- 358 26. Ortín A, Cousens C, Minguijón E, Pascual Z, Villarreal MP, Sharp JM, De las Heras M (2003)
359 Characterization of enzootic nasal tumour virus of goats: complete sequence and tissue distribution. *J*
360 *Gen Virol* 84:2245-2252.
- 361 27. Palmarini M, Sharp JM, De las Heras M, Fan H (1999) Jaagsiekte sheep retrovirus is necessary and
362 sufficient to induce a contagious lung cancer in sheep. *J Virol* 73:6964-6972.
- 363 28. Rosati S, Pittau M, Alberti A, Pozzi S, York DF, Sharp JM, Palmarini M (2000) An accessory open
364 reading frame (orf-x) of jaagsiekte sheep retrovirus is conserved between different virus isolates. *Virus*
365 *Res* 66:109-116.
- 366 29. Maeda N, Palmarini M, Murgia C, Fan H (2001) Direct transformation of rodent fibroblasts by
367 jaagsiekte sheep retrovirus DNA. *Proc Natl Acad Sci USA* 98:4449-4454.
- 368 30. Cousens C, Maeda N, Murgia C, Dagleish MP, Palmarini M, Fan H (2007) In vivo tumorigenesis by
369 Jaagsiekte sheep retrovirus (JSRV) requires Y590 in Env TM, but not full-length orfX open reading
370 frame. *Virology* 367:413-421.

- 371 31. Rai SK, Duh FM, Vigdorovich V, Danilkovitch-Miagkova A, Lerman MI, Miller AD (2001) Candidate
372 tumor suppressor HYAL2 is a glycosylphosphatidylinositol (GPI)-anchored cell-surface receptor for
373 jaagsiekte sheep retrovirus, the envelope protein of which mediates oncogenic transformation. Proc
374 Natl Acad Sci USA 98:4443-4448.
- 375 32. Dirks C, Duh FM, Rai SK, Lerman MI, Miller AD (2002) Mechanism of cell entry and transformation
376 by enzootic nasal tumor virus. J Virol 76:2141-149.
- 377 33. Alberti A, Murgia C, Liu SL, Mura M, Cousens C, Sharp M, Miller AD, Palmarini M (2002)
378 Envelope-induced cell transformation by ovine betaretroviruses. J Virol 76:5387-5394.
- 379 34. Caporale M, Cousens C, Centorame P, Pinoni C, De las Heras M, Palmarini M (2006) Expression of
380 the jaagsiekte sheep retrovirus envelope glycoprotein is sufficient to induce lung tumors in sheep. J
381 Virol 80:8030-8037.
- 382 35. Wootton SK, Halbert CL, Miller AD (2005) Sheep retrovirus structural protein induces lung tumours.
383 Nature 434:904-907.
- 384 36. Wootton SK, Metzger MJ, Hudkins KL, Alpers CE, York D, DeMartini JC, Miller AD (2006) Lung
385 cancer induced in mice by the envelope protein of jaagsiekte sheep retrovirus (JSRV) closely
386 resembles lung cancer in sheep infected with JSRV. Retrovirology 3:94.
- 387 37. Yu DL, Linnerth-Petrik NM, Halbert CL, Walsh SR, Miller AD, Wootton SK (2011) Jaagsiekte sheep
388 retrovirus and enzootic nasal tumor virus promoters drive gene expression in all airway epithelial cells
389 of mice but only induce tumors in the alveolar region of the lungs. J Virol 85:7535-7545.

38. Linnerth-Petrik NM, Santry LA, Yu DL, Wootton SK (2012) Adeno-associated virus vector mediated expression of an oncogenic retroviral envelope protein induces lung adenocarcinomas in immunocompetent mice. *PLoS One* 7:e51400.
39. Wootton SK, Halbert CL, Miller AD (2006) Envelope proteins of jaagsiekte sheep retrovirus and enzootic nasal tumor virus induce similar bronchioalveolar tumors in lungs of mice. *J Virol* 80:9322-9325.
40. McGee-Estrada K, Fan H (2007) Comparison of LTR enhancer elements in sheep beta retroviruses: insights into the basis for tissue-specific expression. *Virus Genes* 35:303-312.
41. Rosales Gerpe MC, van Lieshout LP, Domm JM, Ingrao JC, Datu J, Walsh SR, Yu DL, Jong J, Krell PJ, Wootton SK (2019) The U3 and Env proteins of jaagsiekte sheep retrovirus and enzootic nasal tumor virus both contribute to tissue tropism. *Viruses* 11. pii: E1061.
42. Chow YH, Alberti A, Mura M, Pretto C, Murcia P, Albritton LM, Palmarini M (2003) Transformation of rodent fibroblasts by the jaagsiekte sheep retrovirus envelope is receptor independent and does not require the surface domain. *J Virol* 77:6341-6350.
43. Palmarini M, Maeda N, Murgia C, De-Fraja C, Hofacre A, Fan H (2001) A phosphatidylinositol 3-kinase docking site in the cytoplasmic tail of the Jaagsiekte sheep retrovirus transmembrane protein is essential for envelope-induced transformation of NIH 3T3 cells. *J Virol* 75:11002-11009.
44. Zavala G, Pretto C, Chow YH, Jones L, Alberti A, Grego E, De las Heras M, Palmarini M (2003) Relevance of Akt phosphorylation in cell transformation induced by Jaagsiekte sheep retrovirus. *Virology* 312:95-105.

- 410 45. Liu SL, Lerman MI, Miller AD (2003) Putative phosphatidylinositol 3-kinase (PI3K) binding motifs
411 in ovine betaretrovirus Env proteins are not essential for rodent fibroblast transformation and
412 PI3K/Akt activation. *J Virol* 77:7924-7935.
- 413 46. Liu SL, Miller AD (2005) Transformation of madin-darby canine kidney epithelial cells by sheep
414 retrovirus envelope proteins. *J Virol* 79:927-933.
- 415 47. Hull S, Fan H (2006) Mutational analysis of the cytoplasmic tail of jaagsiekte sheep retrovirus
416 envelope protein. *J Virol* 80:8069-8080.
- 417 48. Maeda N, Fu W, Ortin A, De las Heras M, Fan H (2005) Roles of the Ras-MEK-mitogen-activated
418 protein kinase and phosphatidylinositol 3-kinase-Akt-mTOR pathways in Jaagsiekte sheep retrovirus-
419 induced transformation of rodent fibroblast and epithelial cell lines. *J Virol* 79:4440-4450.
- 420 49. Sun X, Du F, Liu S (2017) Modulation of autophagy in exJSRV-env-transfected cells through the
421 Akt/mTOR and MAPK signaling pathway. *Biochem Biophys Res Commun* 485:672-678.
- 422 50. Varela M, Golder M, Archer F, De las Heras M, Leroux C, Palmarini M (2008) A large animal model
423 to evaluate the effects of Hsp90 inhibitors for the treatment of lung adenocarcinoma. *Virology*
424 371:206-215.
- 425 51. Chitra E, Lin YW, Davamani F, Hsiao KN, Sia C, Hsieh SY, Wei OL, Chen JH, Chow YH (2010)
426 Functional interaction between Env oncogene from Jaagsiekte sheep retrovirus and tumor suppressor
427 Sprouty2. *Retrovirology* 7:62.

- 428 52. Maeda N, Fan H (2008) Signal transduction pathways utilized by enzootic nasal tumor virus (ENTV-
429 1) envelope protein in transformation of rat epithelial cells resemble those used by jaagsiekte sheep
430 retrovirus. *Virus Genes* 36:147-155.
- 431 53. De Las Heras M, Ortín A, Benito A, Summers C, Ferrer LM, Sharp JM (2006) In-situ demonstration
432 of mitogen-activated protein kinase Erk 1/2 signalling pathway in contagious respiratory tumours of
433 sheep and goats. *J Comp Pathol* 135:1-10.
- 434 54. Maeda N, Fan H, Yoshikai Y (2008) Oncogenesis by retroviruses: old and new paradigms. *Rev Med*
435 *Virol* 18:387-405.
- 436 55. Rai SK, DeMartini JC, Miller AD (2000) Retrovirus vectors bearing jaagsiekte sheep retrovirus Env
437 transduce human cells by using a new receptor localized to chromosome 3p21.3. *J Virol* 74:4698-
438 4704.
- 439 56. Liu SL, Duh FM, Lerman MI, Miller AD (2003) Role of virus receptor Hyal2 in oncogenic
440 transformation of rodent fibroblasts by sheep betaretrovirus env proteins. *J Virol* 77:2850-2858.
- 441 57. Monot M, Erny A, Gineys B, Desloire S, Dolmazon C, Aublin-Gex A, Lotteau V, Archer F, Leroux
442 C (2015) Early steps of jaagsiekte sheep retrovirus-mediated cell transformation involve the
443 interaction between Env and the RALBP1 cellular protein. *J Virol* 89:8462-8473.
- 444 58. Hsu T, Phung A, Choe K, Kim JW, Fan H (2015) Role for a Zinc Finger Protein (Zfp111) in
445 Transformation of 208F Rat Fibroblasts by Jaagsiekte Sheep Retrovirus Envelope Protein. *J Virol*
446 89:10453-10466.

- 447 59. Côté M, Zheng YM, Li K, Xiang SH, Albritton LM, Liu SL (2012) Critical role of leucine-valine
448 change in distinct low pH requirements for membrane fusion between two related retrovirus envelopes.
449 J Biol Chem 287:7640-7651.
- 450 60. Côté M, Zheng YM, Albritton LM, Liu SL (2011) Single residues in the surface subunits of oncogenic
451 sheep retrovirus envelopes distinguish receptor-mediated triggering for fusion at low pH and infection.
452 Virology 421:173-183.
- 453 61. Côté M, Zheng YM, Liu SL (2009) Receptor binding and low pH coactivate oncogenic retrovirus
454 envelope-mediated fusion. J Virol 83:11447-11455.
- 455 62. Côté M, Kucharski TJ, Liu SL (2008) Enzootic nasal tumor virus envelope requires a very acidic pH
456 for fusion activation and infection. J Virol 82:9023-9034.
- 457 63. Côté M, Zheng YM, Albritton LM, Liu SL (2008) Fusogenicity of Jaagsiekte sheep retrovirus
458 envelope protein is dependent on low pH and is enhanced by cytoplasmic tail truncations. J Virol
459 82:2543-2554.
- 460 64. Bertrand P, Côté M, Zheng YM, Albritton LM, Liu SL (2008) Jaagsiekte sheep retrovirus utilizes a
461 pH-dependent endocytosis pathway for entry. J Virol 82:2555-2559.
- 462 65. Walsh SR, de Jong JG, van Vloten JP, Gerpe MCR, Santry LA, Wootton SK (2017) Truncation of the
463 enzootic nasal tumor virus envelope protein cytoplasmic tail increases Env-mediated fusion and
464 infectivity. J Gen Virol 98:108-120.

- 465 66. Epand RF, Zhang YL, Mirzabekov T, Kagan B, Silberstein A, Hubbell WL, Epand RM, Chakraborti
466 S, Dimitrov DS, Anderson WF, Rozenberg-Adler Y (2008) Membrane activity of an amphiphilic
467 alpha-helical membrane-proximal cytoplasmic domain of the MoMuLV envelope glycoprotein. *Exp*
468 *Mol Pathol* 84:9-17.
- 469 67. Rozenberg-Adler Y, Conner J, Aguilar-Carreno H, Chakraborti S, Dimitrov DS, Anderson WF (2008)
470 Membrane-proximal cytoplasmic domain of Moloney murine leukemia virus envelope tail facilitates
471 fusion. *Exp Mol Pathol* 84:18-30.
- 472 68. Caporale M, Martineau H, De las Heras M, Murgia C, Huang R, Centorame P, Di Francesco G, Di
473 Gialleonardo L, Spencer TE, Griffiths DJ, Palmarini M (2013) Host species barriers to Jaagsiekte
474 sheep retrovirus replication and carcinogenesis. *J Virol* 87:10752-10762.
- 475 69. Martineau HM, Cousens C, Imlach S, Dagleish MP, Griffiths DJ (2011) Jaagsiekte sheep retrovirus
476 infects multiple cell types in the ovine lung. *J Virol* 85:3341-3355.
- 477 70. Jahns H, Cousens C (2020) Nasal adenocarcinoma associated with jaagsiekte sheep retrovirus
478 infection in a sheep. *J Vet Diagn Invest* 32:152-155.

479

Figure legends

Figure 1 Expression constructs used in this study. Δ GP(JSRV) was previously described [29]. The JSRV *env* gene in the expression plasmid (Δ GP(JSRV-HA)) [47] was replaced with the full-length ENTV-2 *env* gene, designated as Δ GP(ENTV-2). The His-tag (black box) was inserted into Δ GP(ENTV-2) at the C-terminus of the ENTV-2 *env* reading frame. All *env* genes are expressed in rodent cells driven by the CMV promoter. SD: *env* splice donor, SA: *env* splice acceptor

Figure 2 Transformation of rat 208F and RK3E cells. 208F (a-d) and RK3E (e-h) cells were transfected with 5 μ g of Δ GP(ENTV-2) (a, e), Δ GP(ENTV-2-His) (b, f), Δ GP(JSRV) (c, g), and pcDNA3.1(-) (d, h). Foci of transformed 208F cells or RK3E cells at day 11 after transfection are shown (Scale bar = 100 μ m).

Figure 3 Effects of MEK and PI3K inhibitors on the transformation of 208F cells. 208F cells were transfected with 5 μ g of Δ GP(ENTV-2) (b, e, h), Δ GP(JSRV) (c, f, i), and pcDNA3.1(-) (d, g, j), respectively. The number of foci of transformed 208F cells at day 20 after transfection was scored (Scale bar = 200 μ m). The MEK and PI3K inhibitors at a final concentration of 20 μ M were added daily to the cells. For (a), bars indicate mean values $\pm$ standard deviations of three independent experiments. Statistically significant differences are shown as *P* values (****; *P* < 0.0001).

Figure 4 Effects of p38 inhibitor on the transformation of 208F cells. 208F cells were transfected with 5 μ g of Δ GP(ENTV-2) (b, e), Δ GP(JSRV) (c, f), and pcDNA3.1(-) (d, g), respectively. The number of foci of transformed 208F cells at day 13 after transfection was scored (Scale bar = 200 μ m). The p38 inhibitor at a final concentration of 5 μ M was added daily to the cells. For (a), bars indicate mean values $\pm$ standard deviations of three independent experiments. Statistically significant differences are shown as *P* values (***; *P* < 0.001, ****; *P* < 0.0001).

504

505 Figure 5 Activation of signaling pathways in His-tagged ENTV-2 envelope-transformed clone cells.
506 Transformed clone cells (a) with a different morphology as parental 208F cells (b) were established (Scale
507 bar = 100 μ m). Expression of ENTV-2 envelope with His-tag (c), p44/42 MAPK (Erk1/2), and phospho-
508 p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (d), Akt (pan), phospho-Akt (Ser473) (e), p38 MAPK, and
509 phospho-p38 MAPK (Thr180/tyr182) (f), in the transformed clone cells was detected by western blot
510 analysis. For positive controls for detecting phosphorylated proteins, parental 208F cells were stimulated
511 with 100 ng/mL recombinant human EGF to detect phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (d).
512 T: transformed, P: parental, kDa: kilodalton

513

514 Figure 6 Immunohistochemical staining of the caprine ENA tumor and normal goat ethmoid turbinate.
515 Lower magnification micrographs of naturally occurring ENA tumor and normal goat ethmoid turbinate
516 sections immunohistochemically stained with H&E (a, g), phospho-p44/42 MAPK (Thr202/Tyr204) mAb
517 (b, h), and phospho-Akt (Ser473) mAb (c, i) are shown, respectively, (Scale bar = 100 μ m). Higher
518 magnification micrographs of each section are also shown in (d) to (f) for the caprine ENA tumor, and (j)
519 to (l) for normal goat ethmoid turbinate, respectively, (Scale bar = 50 μ m).




Figure 1




Figure 2




Figure 3

(a)





Figure 4




Figure 5




Figure 6