



# HOKKAIDO UNIVERSITY

Title	Transcriptome analysis reveals wingless regulates neural development and signaling genes in the region of wing pigmentation of a polka-dotted fruit fly
Author(s)	Fukutomi, Yuichi; Kondo, Shu; Toyoda, Atsushi et al.
Citation	FEBS Journal, 288(1), 115-126 <a href="https://doi.org/10.1111/febs.15338">https://doi.org/10.1111/febs.15338</a>
Issue Date	2021-01
Doc URL	<a href="https://hdl.handle.net/2115/83728">https://hdl.handle.net/2115/83728</a>
Rights	This is the peer reviewed version of the following article: Yuichi Fukutomi, Shu Kondo, Atsushi Toyoda, Shuji Shigenobu, Shigeyuki Koshikawa, Transcriptome analysis reveals wingless regulates neural development and signaling genes in the region of wing pigmentation of a polka-dotted fruit fly, The FEBS Journal, 288(1): 115-126, 2021, which has been published in final form at <a href="https://doi.org/10.1111/febs.15338">https://doi.org/10.1111/febs.15338</a> . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Type	journal article
File Information	FEBS Journal_fukutomi_koshikawa.pdf



## **Transcriptome analysis reveals *wingless* regulates neural development and signaling genes in the region of wing pigmentation of a polka-dotted fruit fly.**

Yuichi Fukutomi<sup>1</sup>, Shu Kondo<sup>2</sup>, Atsushi Toyoda<sup>3</sup>, Shuji Shigenobu<sup>4</sup>, Shigeyuki Koshikawa<sup>1,5</sup>

1 Graduate School of Environmental Science, Hokkaido University, N10W5, Kita-ku, Sapporo, Hokkaido 060-0810, Japan

2 Invertebrate Genetics Laboratory, National Institute of Genetics, 1111 Yata, Mishima, Shizuoka 411-8540, Japan

3 Comparative Genomics Laboratory, Department of Genomics and Evolutionary Biology, National Institute of Genetics, 1111 Yata, Mishima, Shizuoka, 411-8540, Japan

4 Functional Genomics Facility, National Institute for Basic Biology, Nishigo-naka 38, Myodaiji, Okazaki, Aichi 444-8585, Japan

5 Faculty of Environmental Earth Science, Hokkaido University, N10W5, Kita-ku, Sapporo, Hokkaido 060-0810, Japan

**iD:** S.Kondo, 0000-0002-4625-8379; S.S., 0000-0003-4640-2323; S.Koshikawa, 0000-0003-2831-555X

### **Sub-Discipline:**

Developmental Biology

### **Keywords:**

Pigmentation, fruit fly, regulatory circuit, gene regulatory network, Quartz-Seq

### **Author for correspondence:**

Shigeyuki Koshikawa

email: koshi@ees.hokudai.ac.jp

### **Data accessibility**

Sequence reads were submitted to the DDBJ/EMBL/GenBank. BioProject: PRJDB9109, BioSample: SAMD00198200-SAMD00198299. PE400 (DRX055245); MP3000

(DRX055246); MP5000 (DRX055247).

## Abstract

How evolutionary novelties have arisen is one of the central questions in evolutionary biology. Pre-existing gene regulatory networks or signaling pathways have been shown to be co-opted for building novel traits in several organisms. However, the structure of entire gene regulatory networks and evolutionary events of gene co-option for emergence of a novel trait are poorly understood. In this study, to explore the genetic and molecular bases of the novel wing pigmentation pattern of a polka-dotted fruit fly (*Drosophila guttifera*), we performed *de novo* genome sequencing and transcriptome analyses. As a result, we comprehensively identified the genes associated with the pigmentation pattern. Furthermore, we revealed that 151 of these associated genes were positively or negatively regulated by *wingless*, a master regulator of wing pigmentation. Genes for neural development, Wnt signaling, Dpp signaling, and effectors (such as enzymes) for melanin pigmentation were included among these 151 genes. None of the known regulatory genes that regulate pigmentation pattern formation in other fruit fly species were included. Our results suggest that the novel pigmentation pattern of a polka-dotted fruit fly might have emerged through multi-step co-options of multiple gene regulatory networks, signaling pathways, and effector genes, rather than recruitment of one large gene circuit.

## Introduction

How do evolutionary novelties emerge? Researchers have tried to unravel the developmental genetic program underlying traits in order to clarify the origins of evolutionary novelties [1]. Gene regulatory networks for producing novel traits have been supposed to be composed of a combination of genes forming other traits. One of the most significant current discussions regarding the production of evolutionary novelty is how pre-existing regulatory networks were utilized for this production [1,2].

So far, gene regulatory networks or signaling pathways involved in development of novel traits have been scrutinized in several animals. For example, in a horned dung beetle, limb and wing patterning genes are co-opted for horn formation [3,4]. In Nymphalid butterflies, components of the appendage-patterning gene regulatory network, such as *Distal-less*, *wingless*, and *decapentaplegic* signaling, contributed to development of eyespot, another representative novel trait [5-10]. In the fruit fly *Drosophila melanogaster*, it was shown experimentally that the gene regulatory network

for larval posterior spiracle development was re-used for the posterior lobe, a novel trait observed in male genitalia [11]. Many studies have shown or suggested which gene regulatory networks or signaling pathways are necessary for, or involved in, development of novel traits. However, the structure of entire gene regulatory networks and evolutionary events of gene co-option for emergence of a novel trait are poorly understood.

Fruit fly species have been used to study regulatory evolution of pigmentation pattern, and provided many examples of mechanisms underlying phenotypic evolution [12-15]. In a polka-dotted fruit fly (*Drosophila guttifera*), which has a novel polka-dotted pigmentation pattern on the wings, a melanin synthesis gene, *yellow*, was expressed in the polka-dotted pattern [16-18]. A Wnt signaling gene, *wingless*, was expressed in the centers of pigmentation areas, and positively regulated the expression of *yellow* through an enhancer (Fig. 1a, c, e) [16,17]. Ectopic expression of *wingless* induced ectopic wing pigmentation (Fig. 1b, d, f) [17]. The unique expression pattern of *wingless* seemed to be caused by evolutionary gain of novel enhancer activities [19,20]. In *Drosophila melanogaster*, however, there is no pigmentation around crossveins where *wingless* is expressed. If we assume the ancestral species had *wingless* expression and no pigmentation as in *D. melanogaster*, gain of novel expression pattern of *wingless* alone is not sufficient, for emergence of pigmentation pattern [17]. Also, expression of the melanin synthesis gene *yellow* is not sufficient to induce pigmentation in the *Drosophila melanogaster* wing [16,21], indicating that expression changes of multiple genes were required for the evolution of pigmentation. Therefore, comprehensive exploration of genes downstream of *wingless*, which include both regulatory and effector genes, is necessary for understanding the emergence of the novel wing pigmentation pattern.

In this study, we comprehensively identified the genes that are expressed in pigmentation areas and are also regulated by *wingless*, by *de novo* genome sequencing and two successive transcriptome analyses by Quartz-Seq, a highly sensitive method of RNA sequencing. In the first transcriptome analysis, we compared gene expression patterns between pigmentation areas and an unpigmented area and searched for differentially expressed genes (DEGs). In the second transcriptome analysis, we tested whether those DEGs were regulated by *wingless*, the master control gene for wing pigmentation.

## Results

### Genome sequencing

To obtain the genome sequence of *D. guttifera*, we constructed paired-end and mate pair libraries. The information about the constructed libraries (number of reads, total number of base pairs, average insert size) is summarized in Table 1. After sequencing, we assembled the reads *de novo* to obtain scaffolds. Characteristics of the scaffolds are summarized in Table 2.

### Genes expressed in the polka-dotted pattern

We compared gene expression patterns between pigmentation areas and an unpigmented area, and searched differentially expressed genes (DEGs). These areas can be distinguished by GFP label using a transgenic line which carries *eGFP* connected with an enhancer of *yellow* [17] (Fig. 1e, f). We identified genes upregulated or downregulated commonly in Area 1 (pigmentation area around a campaniform sensillum, Fig. 2a) and Area 2 (vein tip, Fig. 2a), compared with Area 3 (unpigmented, Fig. 2b). Comparison of the gene expression between Area 1 and Area 3 showed that 2333 genes were differentially expressed. Among them, 1390 genes were upregulated (Fig. 3a) and 943 genes were downregulated (Fig. 3b) in Area 1 in comparison to Area 3. 2582 genes were differentially expressed between Area 2 and Area 3. Among them, 1593 genes were upregulated (Fig. 3a) and 989 genes were downregulated (Fig. 3b) in Area 2 in comparison to Area 3. Integrating these data, the number of common DEGs was 1035. Among them, 615 genes were upregulated both in Area 1 and Area 2 (Fig. 3a), while 420 genes were downregulated in Area 1 and Area 2 (Fig. 3b). Consistent with previously reported findings, *wingless* and *yellow* were expressed in the pigmentation areas [17,19], indicating the high sensitivity and accuracy of the present method of analysis. *wingless* and *yellow* were included in the 615 commonly upregulated DEGs (Fig. 4, Table 3, Table 4, (the complete table is available at [https://figshare.com/articles/Genes\\_differentially\\_expressed\\_in\\_the\\_pigmentation\\_areas\\_of\\_Drosophila\\_guttifera\\_/11888898](https://figshare.com/articles/Genes_differentially_expressed_in_the_pigmentation_areas_of_Drosophila_guttifera_/11888898))).

### Pigmentation pattern-associated genes regulated by *wingless*

Because ectopic expression of *wingless* is known to induce pigmentation, genes

sufficient for pigmentation formation in wings must be included in the gene network downstream of *wingless*. To identify the genes that are under the control of *wingless*, we identified genes upregulated or downregulated when *wingless* was ectopically expressed. Among the 615 common upregulated (Area 1 and 2) DEGs, 78 genes were upregulated by ectopic expression of *wingless* (Fig. 3a). In 420 common downregulated (Area 1 and 2) DEGs, *wingless* downregulated 73 genes (Fig. 3b). In total, 151 genes associated with the pigmentation pattern were regulated by *wingless* gene. These 151 genes were blasted against the protein database of *Drosophila melanogaster* and 131 genes were annotated. For these 131 genes, enrichment analysis with DAVID resulted 14 functional annotation clusters, and 6 of which were significant (Table 5). In the most significant cluster, Gene Ontology (GO) terms “Glycoprotein”, “Plasma membrane”, “Disulfide bond”, “Signal peptide” and “Receptor” were included (Table 5). GO terms such as “cuticle pigmentation” and “melanin biosynthetic process” were included in the 3rd significant cluster. 20 genes that could not be annotated were reanalyzed with Blast2GO. Four genes were annotated and remaining 16 genes did not match to any gene in the database.

Among pigmentation pattern associated genes regulated by *wingless*, six genes can be categorized as melanin synthesis-related genes [22-24]. Among them, *yellow*, *laccase2*, and *tan* were upregulated in the pigmentation areas (Fig. 4a, b, Table 3, Table 4) and also upregulated by *wingless* (Fig. 5a, Table 6). *yellow-e*, *yellow-h*, and *silver (svr)* were downregulated both in the pigmentation areas (Fig. 4a, b, c, d, Table 3, Table 4) and in the area where *wingless* was ectopically expressed (Fig. 5a, b, Table 6).

Regulatory genes, such as transcription factors and genes involved in signaling pathways are important to understand the regulatory network controlling the pigmentation pattern. Eight transcription factors were associated with pigmentation and regulated by *wingless*. *Zinc-finger protein interacting with CP190 (ZIPIC)*, *zinc finger protein 28-like*, and Enhancer of split complex genes such as *E(spl)m3-HLH*, *E(spl)m5-HLH*, *E(spl)m7-HLH*, and *lethal (3) malignant brain tumor (l(3)mbt)* were upregulated in the pigmentation areas (Fig. 4c, d, Table 3, Table 4) and by *wingless* (Fig. 5b, Table 6). *Mothers against dpp (Mad)*, and *Mediator complex subunit 18 (MED18)* were downregulated in the pigmentation areas and by *wingless*. Two signal ligands, *Delta (Dl)* and *Wnt oncogene analog 4 (DWnt4)*, ortholog of human *WNT9A/B*, [25]) were upregulated in the pigmentation areas and by *wingless* (Fig. 4c, d, Fig. 5b, Table 3, Table 4, Table 6). As receptors of ligands, *saxophone (sax)* was upregulated and *frizzled 2 (fz2)*

was downregulated in the pigmentation areas and by *wingless* (Fig. 4c, d, Fig. 5b, Table 3, Table 4, Table 6). *wingless* itself was not detected in DEG analysis with ectopic *wingless* expression, which is reasonable in our experimental design. Our transgenic line ectopically drove the *wingless* gene originated from *D. melanogaster* [17] and its transcripts were not mapped on *D. guttifera* genome in the analysis.

## **Discussion**

### **A large number of genes were specifically regulated in the area of pigmentation formation**

Transcriptome analyses revealed that a large number of genes were specifically regulated in the area of pigmentation formation: 78 genes were upregulated commonly in the pigmentation area and by *wingless*, and 73 genes were downregulated commonly in the pigmentation area and by *wingless*. In the butterfly *Bicyclus anynana*, 132 genes were upregulated and 54 genes were downregulated in relation to eye spot formation [26]. In comparison with these numbers of genes in butterflies, the number of genes identified by our results seem reasonable. However, *Drosophila* pigmentation has been thought to be a simple trait, and researchers have tried to explain the evolution of pigmentation by changes of expression of a small number of genes [16,27-29]. In the abdominal tergite of *Drosophila melanogaster*, the combination of *ebony* mutation and ectopic expression of the *yellow* gene can induce ectopic pigmentation [30]. In *D. melanogaster* wings, however, the same combination resulted in scarcely any ectopic pigmentation [16,21]. Those findings are consistent with those of the present study, in which we found many genes that were specifically regulated in the areas of pigmentation formation. This also suggests that experimental reproduction of the gain of pigmentation patterns through overexpression of genes is not trivial.

Although ectopic expression of *wingless* can induce pigmentation, the intensity of induced pigmentation is weaker than that of the natural spotted pigmentation. 537 genes were upregulated and 347 genes were downregulated commonly in Area 1 and Area 2 but not regulated by *wingless* (Fig. 3). Thus, they were not essential to make pigmentation, but might have supplemental roles, or they might be unrelated to pigmentation but have a structural role unique to the pigmented area.

### **Known gene regulatory networks of *Drosophila* pigmentation were not responsible**

### **for *D. guttifer* wing pigmentation**

Regulation of pigmentation has been studied in multiple *Drosophila* models. The best-studied case was abdominal pigmentation in *D. melanogaster*. Male-specific pigmentation was controlled positively by *Abd-B* and negatively by *bab* genes, and pigmentation common to the two sexes was positively controlled by *omb* [31]. In *D. biarmipes* wings, the pigmentation is controlled positively by *Dll*, and negatively by *en* [13,16]. Neither of these genes was included in DEGs in our analysis. Thus, regulatory mechanisms of pigmentation in *D. guttifer* are not a simple co-option of the known gene regulatory network of pigmentation. Then, what kinds of genes are responsible for the emergence of the novel pigmentation pattern in *D. guttifer*?

### ***wingless* regulated neural development genes and other signaling genes specifically in the pigmentation areas**

We identified 12 regulatory genes (transcription factor genes and signaling factor genes) that were regulated by *wingless* in the pigmentation areas. Among them, four (*Dl*, *E(spl)m3-HLH*, *E(spl)m5-HLH* and *E(spl)m7-HLH*) genes belonged to neurogenesis genes of GO terms (Fig. 6). Because Area 2 (vein tip) did not include any tissue of neural origin, these genes were considered to be expressed in epidermal cells, which is the only cell type present in Area 2. In wing discs of *D. melanogaster*, *Dl* and *E(spl)* complex genes are involved in neurogenesis mediated by Notch signaling [32]. During this process, the expression of *wingless* is necessary and sufficient for the expression of *Dl* [33].

Decapentaplegic (Dpp) and Hedgehog (Hh) signaling can work cooperatively or antagonistically with Wingless signaling in various aspects of *Drosophila* development [34,35]. In the present study, genes involved in Dpp signaling, *saxophone (sax)* and *Mothers against dpp (Mad)*, were regulated by *wingless* in the pigmentation areas. *hh* and *patched* of the Hh signaling pathway were up-regulated in the area of pigmentation, but not regulated by *wingless* (Table 7).

Wnt signaling genes *fz2* and *DWnt4* were regulated in the pigmentation areas and by *wingless* (Fig. 6). Downregulation of *fz2* in the pigmentation areas of *D. guttifer* might contribute to achieving the proper gradient of Wingless protein, as it is known to do in *D. melanogaster* wing discs [36]. *DWnt4* was expressed in the area surrounding the *wingless* expression region in wings of *D. guttifer* [19], suggesting that it might play a

role in pigmentation pattern formation.

For the formation of the wing pigmentation pattern of *D. guttifer*, the necessity for a transcription factor gene or signaling factor gene other than *wingless* was suggested in a previous study. When putative binding sites of T-cell factor (Tcf) family proteins [37] in the *yellow* enhancer were artificially mutated, EGFP expression driven by the mutated enhancer was not altered, indicating the canonical Wnt signaling does not directly give input to the enhancer, and another factor does [17]. It is possible that the gene which regulates the expression of *yellow* downstream of *wingless* is included in the genes identified in this study. We also found multiple putative binding sites of Tcf family proteins around most of the genes in Table 6, both in *D. guttifer* and in *D. melanogaster* by using GenePalette [38], suggesting potential connections between *wingless* and these genes, although putative binding sites do not guarantee actual binding or function.

### **Known functions of pigmentation genes regulated by *wingless***

In *D. melanogaster*, *yellow*, *tan*, and *laccase2* are known to be effector genes for pigmentation and have been proven to promote melanin pigmentation in *D. melanogaster* [21,30,39]. Association of these genes with melanin pigmentation patterns was also reported in other insects [23,40,41]. Therefore it was not surprising to detect these three genes in the present study. The *svr* gene, which encodes a carboxypeptidase and is involved in metabolism of N-acetyl dopamine (NADA) in *D. melanogaster* [22,42], was downregulated in the area of pigmentation of *D. guttifer*. Expression of Yellow family protein genes such as *yellow-e* and *yellow-h* was also downregulated in the pigmentation areas on the wings of *D. guttifer*, but the molecular functions of these genes have not been identified in any insect. In wings of the butterflies (*Vanessa caudui* and *Heliconius* spp.), another Yellow family protein gene, *yellow-d*, was upregulated at red pigmentation areas compared with black pigmentation areas [41,43]. Our results together with these previously reported studies suggest that these Yellow protein family genes might play a role in inhibiting melanin pigmentation.

### **Evolutionary scenario of pigmentation pattern evolution**

The genes co-opted for the pigmentation formation are highly likely to have been included in the 151 genes identified in this study. This raises the question: What kind

of genetic change enabled the evolution of pigmentation patterns? There is an ongoing discussion about whether a large gene circuit is recruited or many genes are individually recruited to form a circuit during the evolution of a novel trait [44,45]. Examples of co-option of pre-existing circuits consisting of multiple regulatory genes are known from animals and plants [11,26,46-49]. In the present case, we can ask whether the large gene network for pigmentation regulated by a master control gene, *wingless*, was co-opted all at the same time, or whether individual genes were co-opted one by one to form the current network. To our knowledge, there is no functional evidence of a pigmentation pattern mainly regulated by *wingless* in other *Drosophila* species. Therefore, a "large gene circuit model" does not seem reasonable to explain emergence of the novel pigmentation pattern in *Drosophila guttifera*.

Taking our findings altogether, we suggest that the novel pigmentation pattern of *D. guttifera* was not caused by co-options of a large gene circuit of regulatory genes and effector genes. To test whether the pigmentation pattern has evolved by multi-step co-option, we will have to compare multiple species with and without pigmentation, as well as to test individual gene functions in *D. guttifera*. These investigations will further our understanding of the evolution of pigmentation pattern formation in *D. guttifera*, as an example of the emergence of evolutionary novelties.

## **Materials and methods**

### **Flies**

*Drosophila guttifera* is a North American species that belongs to (or is closely related to) the *quinaria* group of subgenus *Drosophila* [50,51]. The inbred line (A5) was made by ten successive sibling crosses of a wildtype (stock no. 15130-1971.10) obtained from the *Drosophila* Species Stock Center at the University of California, San Diego. Two lines (transgenic lines No. 1 and No.2) of *D. guttifera* were used for transcriptome analyses. Transgenic line No. 1 was established by five successive backcrosses (introgression) of a transgenic line that carries *nuclear eGFP* connected with a *yellow* enhancer (*vein spot* CRE-*nuclear eGFP*, gut 1c+R GFP#12) [17] with the A5 inbred line. Transgenic line No. 2 was established by two successive backcrosses of a UAS-*wg* line [17] with transgenic line No. 1. These backcrosses aimed to unify the genetic backgrounds to improve the mapping efficiency in the transcriptome analyses. Flies were reared with standard cornmeal/sugar/yeast/agar food at 25 °C [52].

## Genome sequencing and gene prediction

Genomic DNA was extracted from adults of the inbred line (A5) with a Genra Puregene Tissue Kit (Qiagen). A paired-end library with an average insert size of 450 bp was constructed with a TruSeq DNA PCR-free Library Prep Kit (Illumina) and two different mate-pair libraries (3 kb and 5 kb) were prepared with a Nextera Mate Pair Library Prep Kit (Illumina). Sequencing libraries were run on the Illumina HiSeq 2500 sequencer with a read length of 150 bp. Paired-end and mate pair reads were *de novo* assembled using Platanus v1.2.1.1 [53] after removal of adaptors and error correction with SOAPec [54]. The assembly sequences had 767 scaffolds with a total length of 168.4 Mb and a scaffold N50 length of 1.8 Mb.

Gene prediction was conducted with Augustus [55] on the scaffolds of the *D. guttifera* inbred line (A5). The option used in the analysis with Augustus was "--species=fly".

## Collecting samples for transcriptome analysis

Pupae at stage P12 (i) or P12 (ii) were used for two successive transcriptome analyses. These stages are just after the stage when *yellow* expression and pigmentation process have started [56]. The transcriptomes were compared by performing the combination of utilization of fluorescence-marked tissues, repetitive microsurgical samplings, and a sensitive RNA sequencing technology. For the first experiment, individuals from the transgenic line No. 1 which carries *eGFP* connected with an enhancer of *yellow* were used. An EGFP-positive area around a campaniform sensillum on 3rd longitudinal vein (Area 1, Fig. 2a), an EGFP-positive area at the tip of 3rd longitudinal vein (Area 2, Fig. 2a), and an EGFP-negative area on 3rd longitudinal vein in wings of flies (Area 3, Fig. 2b) were separated with a surgical knife under a stereo microscope SZX-16 (Olympus). By using those areas on the same wing vein, we aimed to exclude gene expression changes in regions between wing veins. The width of these tissues was about 50  $\mu\text{m}$ . For the second experiment, an EGFP-positive area and an EGFP-negative area with a width of about 75  $\mu\text{m}$  at the same place in individuals from transgenic lines No. 1 and No. 2 were dissected (Fig. 2c, d). From dissected tissues, RNA was collected with an RNeasy Micro Kit (Qiagen) and stored at  $-80^{\circ}\text{C}$ . For RNA extraction, 20 dissected tissues were used for one replicate. Five biological replicates for

each area were prepared (total: 20 x 5 = 100 tissues). The quality of extracted RNA was examined with an Agilent 2100 bioanalyzer (Agilent Technology).

### **RNA sequencing**

The library for RNA sequencing was constructed according to the protocol of Quartz-Seq, a highly sensitive method of RNA sequencing [57]. This protocol includes two PCR steps. Twenty-one cycles were performed for the first PCR, and eight cycles were performed for the second PCR. RNA sequencing was performed with NextSeq 550 (Illumina).

### **Transcriptome analysis and enrichment analysis**

The sequenced transcriptome was mapped to the genome of *D. guttifer* with HISAT2 [58]. Transcriptome assembly was conducted with StringTie [59]. Differentially expressed genes were identified with edgeR [60]. An FDR (false discovery rate) of 0.05 was chosen as the threshold to identify DEGs.

DEGs were blasted against the protein database of *D. melanogaster*, obtained from Ensembl [61,62]. BLAST analysis was performed with Blastx using an E-value < 1e-3. The top hit outcome for each gene was taken as the result of gene annotation. Based on the obtained gene annotation, enrichment analysis was conducted with DAVID [63]. Genes that could not be annotated was reanalyzed with Blast2GO (database: nr, E-value < 1e-3) [64].

### **References**

1. Shubin N, Tabin C, Carroll S. 2009 Deep homology and the origins of evolutionary novelty. *Nature*. 457, 818-823. (doi: 10.1038/nature07891)
2. Wagner GP, Lynch VJ. 2010 Evolutionary novelties. *Curr Biol*. 20, R48-R52. (doi: 10.1016/j.cub.2009.11.010)
3. Moczek AP, Rose DJ. 2009 Differential recruitment of limb patterning genes during development and diversification of beetle horns. *Proc Natl Acad Sci U S A*. 106, 8992-8997. (doi: 10.1073/pnas.0809668106)
4. Hu Y, Linz DM, Moczek AP. 2019 Beetle horns evolved from wing serial homologs. *Science*. 366, 1004-1007. (doi: 10.1126/science.aaw2980)
5. Nijhout HF. 1980 Pattern formation on lepidopteran wings: determination of an eyespot.

- Dev Biol.* 80, 267-274. (doi: 10.1016/0012-1606(80)90403-0)
6. Carroll SB, Gates J, Keys DN, Paddock SW, Panganiban GE, Selegue JE, Williams JA. 1994 Pattern formation and eyespot determination in butterfly wings. *Science*. 265, 109-114. (doi: 10.1126/science.7912449)
  7. Brakefield PM, Gates J, Keys D, Kesbeke F, Wijngaarden PJ, Monteiro A, French V, Carroll SB. 1996 Development, plasticity and evolution of butterfly eyespot patterns. *Nature*. 384, 236-242. (doi: 10.1038/384236a0)
  8. Monteiro A, Glaser G, Stockslager S, Glansdorp N, Ramos D. 2006 Comparative insights into questions of lepidopteran wing pattern homology. *BMC Dev Biol.* 6, 52. (doi: 10.1186/1471-213X-6-52)
  9. Zhang L, Reed RD. 2016 Genome editing in butterflies reveals that *spalt* promotes and *Distal-less* represses eyespot colour patterns. *Nat Commun.* 7, 11769. (doi: 10.1038/ncomms11769)
  10. Connahs H, Tlili S, van Creijl J, Loo TYJ, Banerjee TD, Saunders TE, Monteiro A. 2019 Activation of butterfly eyespots by *Distal-less* is consistent with a reaction-diffusion process. *Development*. 146, dev169367. (doi: 10.1242/dev.169367)
  11. Glassford WJ, Johnson WC, Dall NR, Smith SJ, Liu Y, Boll W, Noll M, Rebeiz M. 2015 Co-option of an ancestral Hox-regulated network underlies a recently evolved morphological novelty. *Dev Cell*. 34, 520-531. (doi: 10.1016/j.devcel.2015.08.005)
  12. Williams TM, Selegue JE, Werner T, Gompel N, Kopp A, Carroll SB. 2008 The regulation and evolution of a genetic switch controlling sexually dimorphic traits in *Drosophila*. *Cell*. 134, 610-623. (doi: 10.1016/j.cell.2008.06.052)
  13. Arnoult L, Su KF, Manoel D, Minervino C, Magriña J, Gompel N, Prud'homme B. 2013 Emergence and diversification of fly pigmentation through evolution of a gene regulatory module. *Science*. 339, 1423-1426. (doi: 10.1126/science.1233749)
  14. Rebeiz M, Williams TM. 2017 Using *Drosophila* pigmentation traits to study the mechanisms of *cis*-regulatory evolution. *Curr Opin Insect Sci.* 19, 1-7. (doi: 10.1016/j.cois.2016.10.002)
  15. Koshikawa S. 2020 Evolution of wing pigmentation in *Drosophila*: Diversity, physiological regulation, and *cis*-regulatory evolution. *Dev Growth Diff.* (doi: 10.1111/dgd.12661)
  16. Gompel N, Prud'homme B, Wittkopp PJ, Kassner VA, Carroll SB. 2005 Chance caught on the wing: *cis*-regulatory evolution and the origin of pigment patterns in

- Drosophila*. *Nature*. 433, 481-487. (doi: 10.1038/nature03235)
17. Werner T, Koshikawa S, Williams TM, Carroll SB. 2010 Generation of a novel wing colour pattern by the Wingless morphogen. *Nature*. 464, 1143-1148. (doi: 10.1038/nature08896)
  18. Koshikawa S, Fukutomi Y, Matsumoto K. 2017 *Drosophila guttifera* as a model system for unraveling color pattern formation. In: Sekimura T, Nijhout HF (eds) Diversity and evolution of butterfly wing patterns: an integrative approach. 287-301. Springer, New York. (doi: 10.1007/978-981-10-4956-9\_16)
  19. Koshikawa S, Giorgianni MW, Vaccaro K, Kassner VA, Yoder JH, Werner T, Carroll SB. 2015 Gain of *cis*-regulatory activities underlies novel domains of *wingless* gene expression in *Drosophila*. *Proc Natl Acad Sci U S A*. 112, 7524-7529. (doi: 10.1073/pnas.1509022112)
  20. Koshikawa S. 2015 Enhancer modularity and the evolution of new traits. *Fly*. 9, 155-159. (doi: 10.1080/19336934.2016.1151129)
  21. Riedel F, Vorkel D, Eaton S. 2011 Megalin-dependent Yellow endocytosis restricts melanization in the *Drosophila* cuticle. *Development*. 138, 149-158. (doi: 10.1242/dev.056309)
  22. Walter MF, Zeineh LL, Black BC, McIvor WE, Wright TR, Biessmann H. 1996 Catecholamine metabolism and in vitro induction of premature cuticle melanization in wild type and pigmentation mutants of *Drosophila melanogaster*. *Arch Insect Biochem Physiol*. 31, 219-233. (doi: 10.1002/(SICI)1520-6327(1996)31:2<219::AID-ARCH9>3.0.CO;2-U)
  23. Wittkopp PJ, Beldade P. 2009 Development and evolution of insect pigmentation: genetic mechanisms and the potential consequences of pleiotropy. *Semin Cell Dev Biol*. 20, 65-71. (doi: 10.1016/j.semcdb.2008.10.002)
  24. Ferguson LC, Green J, SurrIDGE A, Jiggins CD. 2011 Evolution of the insect *yellow* gene family. *Mol Biol Evol*. 28, 257-272. (doi: 10.1093/molbev/msq192)
  25. Janssen R, Le Gouar M, Pechmann M, Poulin F, Bolognesi R, Schwager EE, Hopfen C, Colbourne JK, Budd GE, Brown SJ, Prpic NM, Kosiol C, Vervoort M, Damen WG, Balavoine G, McGregor AP. 2010 Conservation, loss, and redeployment of Wnt ligands in protostomes: implications for understanding the evolution of segment formation. *BMC Evol Biol*. 10, 374. (doi: 10.1186/1471-2148-10-374)
  26. Özsü N, Monteiro A. 2017 Wound healing, calcium signaling, and other novel

- pathways are associated with the formation of butterfly eyespots. *BMC Genomics*. 18, 788. (doi: 10.1186/s12864-017-4175-7)
27. Jeong S, Rokas A, Carroll SB. 2006 Regulation of body pigmentation by the Abdominal-B Hox protein and its gain and loss in *Drosophila* evolution. *Cell*. 125:1387-1399. (doi: 10.1016/j.cell.2006.04.043)
  28. Jeong S, Rebeiz M, Andolfatto P, Werner T, True J, Carroll SB. 2008 The evolution of gene regulation underlies a morphological difference between two *Drosophila* sister species. *Cell*. 132, 783-793. (doi: 10.1016/j.cell.2008.01.014)
  29. Rebeiz M, Pool JE, Kassner VA, Aquadro CF, Carroll SB. 2009 Stepwise modification of a modular enhancer underlies adaptation in a *Drosophila* population. *Science*. 326, 1663-1667. (doi:10.1126/science.1178357)
  30. Wittkopp PJ, True JR, Carroll SB. 2002 Reciprocal functions of the *Drosophila* Yellow and Ebony proteins in the development and evolution of pigment patterns. *Development*. 129, 1849-1858.
  31. Kopp A, Duncan I, Godt D, Carroll SB. 2000 Genetic control and evolution of sexually dimorphic characters in *Drosophila*. *Nature*. 408, 553-559. (doi: 10.1038/35046017)
  32. de Celis JF, de Celis J, Ligoxygakis P, Preiss A, Delidakis C, Bray S. 1996 Functional relationships between *Notch*, *Su(H)* and the bHLH genes of the *E(spl)* complex: the *E(spl)* genes mediate only a subset of *Notch* activities during imaginal development. *Development*. 122, 2719-2728.
  33. Micchelli CA, Rulifson EJ, Blair SS. 1997 The function and regulation of cut expression on the wing margin of *Drosophila*: *Notch*, *Wingless* and a dominant negative role for *Delta* and *Serrate*. *Development*. 124, 1485-1495.
  34. Theisen H, Haerry TE, O'Connor MB, Marsh JL. 1996 Developmental territories created by mutual antagonism between *Wingless* and *Decapentaplegic*. *Development*. 122, 3939-3948.
  35. Basler K, Struhl G. 1994 Compartment boundaries and the control of *Drosophila* limb pattern by *hedgehog* protein. *Nature*. 368, 208-214. (doi: 10.1038/368208a0)
  36. Cadigan KM, Fish MP, Rulifson EJ, Nusse R. 1998 *Wingless* repression of *Drosophila frizzled 2* expression shapes the *Wingless* morphogen gradient in the wing. *Cell*. 93, 767-777. (doi: 10.1016/s0092-8674(00)81438-5)
  37. Gustavson MD, Crawford HC, Fingleton B, Matrisian LM. 2004 Tcf binding

- sequence and position determines beta-catenin and Lef-1 responsiveness of MMP-7 promoters. *Mol Carcinog.* 41, 125-139. (doi: 10.1002/mc.20049)
38. Smith, AF, Posakony, JW, Rebeiz, M. 2017 Automated tools for comparative sequence analysis of genic regions using the GenePalette application. *Developmental Biology*, 429, 158-164. (doi: 10.1016/j.ydbio.2017.06.033)
39. True JR, Yeh SD, Hovemann BT, Kemme T, Meinertzhagen IA, Edwards TN, Liou SR, Han Q, Li J. 2005 *Drosophila tan* encodes a novel hydrolase required in pigmentation and vision. *PLoS Genet.* 1, e63. (doi: 10.1371/journal.pgen.0010063)
40. Futahashi R, Banno Y, Fujiwara H. 2010 Caterpillar color patterns are determined by a two-phase melanin gene pre patterning process: new evidence from *tan* and *laccase2*. *Evol Dev.* 12, 157-167. (doi: 10.1111/j.1525-142X.2010.00401.x)
41. Zhang L, Martin A, Perry MW, van der Burg KR, Matsuoka Y, Monteiro A, Reed RD. 2017 Genetic basis of melanin pigmentation in butterfly wings. *Genetics.* 205, 1537-1550. (doi: 10.1534/genetics.116.196451)
42. Wright TR. 1987 The genetics of biogenic amine metabolism, sclerotization, and melanization in *Drosophila melanogaster*. *Adv Genet.* 24, 127-222. (doi: 10.1016/S0065-2660(08)60008-5)
43. Hines HM, Papa R, Ruiz M, Papanicolaou A, Wang C, Nijhout HF, McMillan WO, Reed RD. 2012 Transcriptome analysis reveals novel patterning and pigmentation genes underlying *Heliconius* butterfly wing pattern variation. *BMC Genomics.* 13, 288. (doi: 10.1186/1471-2164-13-288)
44. True JR, Carroll SB. 2002 Gene co-option in physiological and morphological evolution. *Annu Rev Cell Dev Biol.* 18, 53-80. (doi: 10.1146/annurev.cellbio.18.020402.140619)
45. Carroll SB, Grenier JK, Weatherbee SD 2013 From DNA to diversity: molecular genetics and the evolution of animal design. John Wiley & Sons.
46. Keys DN, Lewis DL, Selegue JE, Pearson BJ, Goodrich LV, Johnson RL, ... Carroll SB 1999 Recruitment of a hedgehog regulatory circuit in butterfly eyespot evolution. *Science.* 283, 532-534. (doi: 10.1126/science.283.5401.532)
47. Gao F, Davidson EH. 2008 Transfer of a large gene regulatory apparatus to a new developmental address in echinoid evolution. *Proc Natl Acad Sci U S A.* 105, 6091-6096. (doi: 10.1073/pnas.0801201105)
48. Lemons D, Fritzenwanker H, Gerhart J, Lowe CJ, McGinnis W. 2010 Co-option of

- an anteroposterior head axis patterning system for proximodistal patterning of appendages in early bilaterian evolution. *Dev Biol.* 344, 358-362. (doi: 10.1016/j.ydbio.2010.04.022)
49. Nakayama H, Yamaguchi T, Tsukaya H. 2012 Acquisition and diversification of cladodes: leaf-like organs in the genus *Asparagus*. *Plant Cell.* 24, 929-940. (doi: 10.1105/tpc.111.092924)
50. Izumitani HF, Kusaka Y, Koshikawa S, Toda MJ, Katoh T. 2016 Phylogeography of the Subgenus *Drosophila* (Diptera: Drosophilidae): evolutionary history of faunal divergence between the Old and the New Worlds. *PloS One.* 11, e0160051. (doi: 10.1371/journal.pone.0160051)
51. Chialvo CHS, White BE, Reed LK, Dyer KA. 2019 A phylogenetic examination of host use evolution in the *quinaria* and *testacea* groups of *Drosophila*. *Mol Phylogenet Evol.* 130, 233-243. (doi: 10.1016/j.ympev.2018.10.027)
52. Fukutomi Y, Matsumoto K, Funayama N, Koshikawa S. 2018 Methods for staging pupal periods and measurement of wing pigmentation of *Drosophila guttifera*. *J Vis Exp.* 131, e56935. (doi: 10.3791/56935)
53. Kajitani R, Toshimoto K, Noguchi H, Toyoda A, Ogura Y, Okuno M, Yabana M, Harada M, Nagayasu E, Maruyama H, Kohara Y, Fujiyama A, Hayashi T, Itoh T. 2014 Efficient *de novo* assembly of highly heterozygous genomes from whole-genome shotgun short reads. *Genome Res.* 24, 1384-1395. (doi: 10.1101/gr.170720.113)
54. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu SM, Peng S, Xiaoqian Z, Liu G, Liao X, Li Y, Yang H, Wang J, Lam TW, Wang J. 2012 SOAPdenovo2: an empirically improved memory-efficient short-read *de novo* assembler. *Gigascience.* 1,18. (doi: 10.1186/2047-217X-1-18)
55. Stanke M, Steinkamp R, Waack S, Morgenstern B. 2004 AUGUSTUS: a web server for gene finding in eukaryotes. *Nucleic Acids Res.* W309-W312. (doi: 10.1093/nar/gkh379)
56. Fukutomi Y, Matsumoto K, Agata K, Funayama N, Koshikawa S. 2017 Pupal development and pigmentation process of a polka-dotted fruit fly, *Drosophila guttifera* (Insecta, Diptera). *Dev Genes Evol.* 227, 171-180. (doi: 10.1007/s00427-017-0578-3)

57. Sasagawa Y, Nikaido I, Hayashi T, Danno H, Uno KD, Imai T, Ueda HR. 2013 Quartz-Seq: a highly reproducible and sensitive single-cell RNA sequencing method, reveals non-genetic gene-expression heterogeneity. *Genome Biol.* 14, 3097. (doi: 10.1186/gb-2013-14-4-r31)
58. Kim D, Langmead B, Salzberg SL. 2015 HISAT: a fast spliced aligner with low memory requirements. *Nat Methods.* 12, 357–360. (doi: 10.1038/nmeth.3317)
59. Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015 StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol.* 33, 290–295. (doi: 10.1038/nbt.3122)
60. Robinson MD, McCarthy DJ, Smyth GK. 2010 edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics.* 26, 139–140. (doi: 10.1093/bioinformatics/btp616)
61. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009 BLAST+: architecture and applications. *BMC Bioinform.* 10, 421. (doi: 10.1186/1471-2105-10-421)
62. Howe KL, Contreras-Moreira B, De Silva N, Maslen G, Akanni W, Allen J, Alvarez-Jarreta J, Barba M, Bolser DM, Cambell L, Carbajo M, Chakiachvili M, Christensen M, Cummins C, Cuzick A, Davis P, Fexova S, Gall A, George N, Gil L, Gupta P, Hammond-Kosack KE, Haskell E, Hunt SE, Jaiswal P, Janacek SH, Kersey PJ, Langridge N, Maheswari U, Maurel T, McDowall MD, Moore B, Muffato M, Naamati G, Naithani S, Olson A, Papatheodorou I, Patricio M, Paulini M, Pedro H, Perry E, Preece J, Rosello M, Russell M, Sitnik V, Staines DM, Stein J, Tello-Ruiz MK, Trevanion SJ, Urban M, Wei S, Ware D, Williams G, Yates AD, Flicek P. 2019 Ensembl Genomes 2020—enabling non-vertebrate genomic research. *Nucleic Acids Res.* gkz890. (doi: 10.1093/nar/gkz890)
63. Huang DW, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R, Baseler MW, Lane HC, Lempicki RA. 2007 The DAVID Gene Functional Classification Tool: a novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol.* 8, R183. (doi: 10.1186/gb-2007-8-9-r183)
64. Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. 2005 Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics.* 21, 3674-3676. (doi: 10.1093/bioinformatics/bti610)

## Tables

**Table 1**

The data of the libraries of *D. guttifera* constructed for genome sequencing. PE400, MP3kbp, and MP5kbp respectively indicate the paired-end library, the mate-pair library (3kb), and the mate-pair library (5kb).

Library	Number of reads	Total number of base pairs (bp)	Average insert size (bp)
PE400	151,389,704	22,708,455,600	450
MP3kbp	41,484,414	6,222,662,100	3,169
MP5kbp	30,474,280	4,571,142,000	5,227

**Table 2**

Characteristics of the *D. guttifera* genome sequence obtained by *de novo* assembly.

Characteristics	Measures
Total number of base pairs (bp)	168,421,893
Number of scaffolds	767
Scaffold N50 (bp)	1,784,351
Contig N50 (bp)	182,614

**Table 3**

Genes differentially expressed in a pigmentation area around a campaniform sensillum. FC, CPM, and FDR indicate fold change, counts per million, and false discovery rate,

respectively.

	logFC	logCPM	FDR
<i>y</i>	4.97989524	13.1819431	2.38E-36
<i>t</i>	2.76380354	12.0058893	3.29E-32
<i>laccase2</i>	2.49456691	11.8424645	2.15E-33
<i>yellow-e</i>	-1.6407385	10.3188765	1.81E-15
<i>yellowe-h</i>	-0.5731037	10.5879713	0.02343591
<i>svr</i>	-0.7948767	9.41379963	0.00438384
<i>ZIPIC</i>	4.53189996	0.48568358	0.0208437
<i>zinc finger protein 28-like</i>	6.75852343	1.85690936	0.0005392
<i>E(spl)m3-HLH</i>	1.23731928	8.47601255	0.0003062
<i>E(spl)m5-HLH</i>	6.77199195	3.84590398	6.05E-10
<i>E(spl)m7-HLH</i>	3.36253299	4.92651899	0.0000107
<i>Mad</i>	-1.5333172	5.94437046	0.02335105
<i>Dl</i>	1.70066765	8.69275453	3.05E-07
<i>DWnt4</i>	7.88039218	2.78881957	0.00610713
<i>sax</i>	1.84041069	5.74292161	0.04887547
<i>fz2</i>	-1.7268879	8.84471432	1.56E-08
<i>wg</i>	4.90156885	0.66703408	0.00181758
<i>MED18</i>	-7.5365154	2.49510051	0.00031397
<i>l(3)mbt</i>	4.03476224	4.54864458	0.00016924

**Table 4**

Genes differentially expressed in a pigmentation area at the tip of a vein. FC, CPM, and FDR indicate fold change, counts per million, and false discovery rate, respectively.

	logFC	logCPM	FDR
<i>y</i>	4.21635634	12.5960958	5.72E-28
<i>t</i>	3.17677306	12.5066497	3.28E-46

<i>laccase2</i>	1.9708477	11.544914	8.32E-28
<i>yellow-e</i>	-2.5486142	10.2771931	4.54E-32
<i>yellowe-h</i>	-0.6835507	10.671958	0.00134155
<i>svr</i>	-0.6447186	9.60117515	0.00423173
<i>ZIPIC</i>	4.73395417	0.69571295	0.0256575
<i>zinc finger protein 28-like</i>	5.82784953	1.30102744	0.00000242
<i>E(spl)m3-HLH</i>	1.3827603	8.72212278	0.0000125
<i>E(spl)m5-HLH</i>	5.06729935	2.44510302	0.00069965
<i>E(spl)m7-HLH</i>	3.60181175	5.29016258	0.00000958
<i>Mad</i>	-2.0456884	5.94769958	0.01764714
<i>Dl</i>	0.92496862	8.28076591	0.01422465
<i>DWnt4</i>	7.3910185	2.48248069	0.00010193
<i>sax</i>	2.05209481	6.05221034	0.01234576
<i>fz2</i>	-2.6177129	8.81769459	2.23E-19
<i>wg</i>	6.33393063	1.65109029	0.00062794
<i>MED18</i>	-7.5507612	2.61383665	0.00022139
<i>l(3)mbt</i>	3.71774754	4.41678381	0.03790313

**Table 5**

The results from enrichment analysis with DAVID, showing clusters with enrichment score > 1.3 and Gene ontology terms with *p* value < 0.05 in each cluster (Count indicates the number of genes for each Gene ontology term).

<b>Cluster 1</b>		
<b>enrichment score: 2.42</b>	<b>Count</b>	<b><i>p</i> value</b>
Glycosylation site: N-linked	16	1.90E-05
Glycoprotein	16	4.90E-05
Plasma membrane	18	8.00E-04

Cell membrane	11	2.90E-03
Topological domain: Extracellular	10	6.10E-03
Disulfide bond	14	7.60E-03
Topological domain: Cytoplasmic	11	9.70E-03
Transmembrane region	13	1.70E-02
Signal peptide	10	4.20E-02
Receptor	9	4.60E-02

---

**Cluster 2**

	<b>Count</b>	<b>p value</b>
<b>enrichment score: 2.38</b>		
Extracellular matrix	8	3.40E-04
Chitin-based cuticle development	7	2.60E-03
Structural constituent of cuticle	6	3.90E-03
Structural constituent of chitin-based larval cuticle	5	1.70E-02
Insect cuticle protein	5	2.10E-02

---

**Cluster 3**

	<b>Count</b>	<b>p value</b>
<b>enrichment score: 2.18</b>		
Cuticle pigmentation	4	6.60E-04
Melanin biosynthetic process	3	7.40E-03

Major royal jelly	3	7.80E-03
<b>Cluster 4</b>		
<b>enrichment score: 1.39</b>	<b>Count</b>	<b>p value</b>
Membrane	42	1.20E-02
<b>Cluster 5</b>		
<b>enrichment score: 1.34</b>	<b>Count</b>	<b>p value</b>
ANK repeat	4	3.60E-02
Ankyrin repeat	4	4.70E-02
<b>Cluster 6</b>		
<b>enrichment score: 1.33</b>	<b>Count</b>	<b>p value</b>
Carbohydrate metabolic process	5	1.40E-02

**Table 6**

Genes differentially expressed in an area where *wingless* is ectopically expressed. FC, CPM, and FDR indicate fold change, counts per million, and false discovery rate, respectively.

	logFC	logCPM	FDR
<i>y</i>	4.54116336	12.8494004	3.6E-148
<i>t</i>	2.40367879	12.0035577	1.25E-74
<i>laccase2</i>	2.04478128	11.4946868	3.67E-39
<i>yellow-e</i>	-1.2740677	11.392683	1.96E-25
<i>yellowe-h</i>	-0.4516609	9.7843299	0.00750851
<i>svr</i>	-0.6215228	9.01017126	0.00382779
<i>ZIPIC</i>	5.0147075	-0.3005216	0.0284088
<i>zinc finger protein 28-like</i>	5.61711109	2.5793453	0.00012374

<i>E(spl)m3-HLH</i>	0.53145394	8.24183597	0.00380777
<i>E(spl)m5-HLH</i>	5.25665975	1.58014564	0.00043059
<i>E(spl)m7-HLH</i>	1.67127368	5.65074921	1.23E-15
<i>Mad</i>	-0.8277961	4.89916451	0.01393637
<i>Dl</i>	1.02257261	7.82859549	8.12E-11
<i>DWnt4</i>	8.32686641	2.16749637	4.35E-09
<i>sax</i>	0.91143441	5.58205955	0.00742031
<i>fz2</i>	-0.4058316	8.27179547	0.02026252
<i>wg</i>	0	-1.2863604	1
<i>MED18</i>	-6.8364613	0.89990996	0.00079404
<i>l(3)mbt</i>	1.93217694	4.83757955	0.01514216

**Table 7**

Data of expression of *hedgehog* (*hh*), *patched* (*ptc*), and *smoothened* (*smo*) in a pigmentation area around a campaniform sensillum, at the tip of a vein, and in an area where *wingless* is ectopically expressed.

		logFC	logCPM	FDR
	Campaniform sensillum	4.85624359	0.63602448	0.04417015
<i>hh</i>	Vein tip	7.03066917	2.18407496	0.01422465
	<i>wingless</i> overexpression	-4.2417977	-0.6349855	0.0705383
	Campaniform sensillum	6.41499537	3.16486805	0.0012656
<i>ptc</i>	Vein tip	5.79689401	2.74718533	0.02202927
	<i>wingless</i> overexpression	-0.6229317	3.57406688	1
	Campaniform sensillum	-1.4152405	5.26637498	0.51827634

<i>smo</i>	Vein tip	-6.8976128	4.95229132	1.01E-08
	<i>wingless</i> overexpression	0.52985214	4.07845493	0.95456002

---

## Figure legends

### Fig. 1

Wings of *Drosophila guttifer* and the expression pattern of *wingless*. **a:** A wing of *Drosophila guttifer*. This picture shows wild type pigmentation pattern. **b:** The wing pigmentation pattern of an individual in which *wingless* is ectopically expressed. **c:** The expression pattern of *wingless* (blue) in a wing of a wild type individual. **d:** The position where *wingless* is ectopically expressed (blue) in the fly shown in b (Modified from [17]). **e:** The expression pattern of EGFP driven by an enhancer of *yellow*. In this individual, the expression pattern of *wingless* is the same as wild type. **f:** The expression pattern of EGFP driven by an enhancer of *yellow* in an individual with ectopic *wingless* expression shown in d. Individuals shown in e and f were from transgenic lines No. 1 and No. 2 for each (see Materials and methods). For experiments, 100 individuals from No. 1 and No. 2 for each were used. The expression pattern of EGFP of 100 individuals from No. 1 and No. 2 were the same as the patterns shown in e and f for each. The scale bar indicates 250  $\mu$ m.

### Fig. 2

Pupal wing tissues used for transcriptome analyses. Dissected areas are indicated with white boxes. **a:** Area 1 (an EGFP-positive area around a campaniform sensillum on the 3rd longitudinal vein) and Area 2 (an EGFP-positive area at the tip of the 3rd longitudinal vein, which does not include bristles or other tissue of neural origin). **b:** Area 3 (an EGFP-negative area on the 3rd longitudinal vein). **c:** An EGFP-positive area where *wingless* is ectopically expressed. **d:** An EGFP-negative area where *wingless* is not ectopically expressed. For each area, 20 tissues were dissected for one replicate and 5 biological replicates were used for experiments. Scale bars indicate 250  $\mu$ m.

### Fig. 3

The number of differentially expressed genes (DEGs) detected in transcriptome analyses.

**a:** The number of DEGs upregulated at pigmentation areas and an area where *wingless* is ectopically expressed. **b:** The number of DEGs downregulated at the pigmentation areas and an area where *wingless* is ectopically expressed. The circle labeled with “Campaniform sensillum” indicates the result from the comparison of transcriptome between Area 1 (a pigmentation area around campaniform sensillum) and Area 3 (unpigmented). The circle labeled with “Vein tip” indicates the result from the comparison of transcriptome between Area 2 (a pigmentation area at the tip of 3rd longitudinal vein) and Area 3. The circle labeled with “*wingless* overexpression” indicates the number of genes differentially expressed where *wingless* is ectopically expressed.

#### **Fig. 4**

Volcano plots of the results from the first transcriptome analysis. The horizontal axis indicates fold changes and the vertical axis indicates significance calculated as FDR with edgeR. Orange points indicate upregulated DEGs and blue points indicate downregulated DEGs. **a:** This plot shows genes differentially expressed in Area 1 (a pigmentation area around a campaniform sensillum) compared with gene expression of Area 3 (unpigmented). **b:** This plot shows genes differentially expressed in Area 2 (a pigmentation area at the tip of 3rd longitudinal vein) compared with gene expression of Area 3. **c:** Genes with  $-\log_{10}(\text{FDR}) < 20$  are extracted from a. **d:** Genes with  $-\log_{10}(\text{FDR}) < 20$  are extracted from b. In c and d, *m3*, *m5*, and *m7* indicate *E(spl)m3-HLH*, *E(spl)m5-HLH*, and *E(spl)m7-HLH* respectively.

#### **Fig. 5**

Volcano plots of the results from the second transcriptome analysis. The horizontal axis indicates fold changes and the vertical axis indicates significance calculated as FDR with edgeR. Orange points indicate DEGs upregulated and blue points indicate downregulated DEGs. **a:** This plot shows genes differentially expressed where *wingless* is ectopically expressed. **b:** Genes with  $-\log_{10}(\text{FDR}) < 20$  are extracted from a.

#### **Fig. 6**

The putative gene regulatory network for formation of the novel wing pigmentation pattern. Green box: Notch signaling gene, grey boxes: melanin synthesis genes, yellow

boxes: Wnt signaling genes, and blue boxes: Dpp signaling genes.

### **Acknowledgements**

We thank Sean B. Carroll and Thomas Werner for providing fly stocks; Tsuyoshi Katahata and Machiko Teramoto for fly stock maintenance; Arnaud Martin, Kiyokazu Agata, Noriko Funayama, Keiji Matsumoto, Naoyuki Fuse, Takeshi Inoue, Norito Shibata, Katsushi Yamaguchi, and Takahiro Bino for scientific advice; two anonymous referees for helpful comments; and Elizabeth Nakajima for English editing. This work was supported by Functional Genomics Facility, NIBB Core Research Facilities. This work was funded by NBRP Genome Information Upgrading Program (*Drosophila*) to S.Kondo, NIBB Collaborative Research Program (15-835, 16-426, 17-423, 18-402, 19-403) to S.Koshikawa, KAKENHI (17K19427, 18H02486) to S.Koshikawa, KAKENHI (18J20452) to Y.F., and Yamada Science Foundation to S.Koshikawa. Y.F. is a JSPS Research Fellow.

### **Authors' contributions**

Y.F., S.Kondo, S.S. and S.Koshikawa conceived of and designed the study; Y.F., S.Kondo, and A.T. collected data and conducted the analysis; Y.F., A.T. and S.Koshikawa drafted the manuscript, with input from S.Kondo and S.S.; Y.F. and S.Koshikawa edited and wrote the final version; all authors gave final approval for publication and agree to be held accountable for the work performed therein.

### **Competing interests**

We declare we have no competing interests.




Fig. 1




Fig. 2





Fig. 3




Fig. 4





Fig. 5




Fig. 6