



Title	Immersive 3-Dimensional Virtual Reality Modeling for Case-Specific Presurgical Discussions in Cerebrovascular Neurosurgery
Author(s)	Sugiyama, Taku; Clapp, Tod; Nelson, Jordan et al.
Citation	Operative neurosurgery, 20(3), 289-299 <a href="https://doi.org/10.1093/ons/opaa335">https://doi.org/10.1093/ons/opaa335</a>
Issue Date	2021-03
Doc URL	<a href="https://hdl.handle.net/2115/84221">https://hdl.handle.net/2115/84221</a>
Rights	This is a pre-copyedited, author-produced version of an article accepted for publication in Operative Neurosurgery following peer review. The version of record Taku Sugiyama, Tod Clapp, Jordan Nelson, BS, Chad Eitel, BS, Hiroaki Motegi, Naoki Nakayama, Tsukasa Sasaki, RT, Kikutaro Tokairin, Masaki Ito, Ken Kazumata, Kiyohiro Houkin, Immersive 3-Dimensional Virtual Reality Modeling for Case-Specific Presurgical Discussions in Cerebrovascular Neurosurgery, Operative Neurosurgery, Volume 20, Issue 3, March 2021, Pages 289-299 is available online at: <a href="https://doi.org/10.1093/ons/opaa335">https://doi.org/10.1093/ons/opaa335</a>
Type	journal article
File Information	Oper Neurosurg 20 289-299.pdf



1 **Title: Immersive Three-Dimensional Virtual Reality Modeling for Case-Specific**  
2 **Presurgical Discussions in Cerebrovascular Neurosurgery**

3

4 Taku Sugiyama, MD, PhD<sup>1)</sup>, Tod Clapp, PhD<sup>2)</sup>, Jordan Nelson, BS<sup>2)</sup>, Chad Eitel, BS<sup>2)</sup>, Hiroaki  
5 Motegi, MD, PhD<sup>1)</sup>, Naoki Nakayama, MD, PhD<sup>1)</sup>, Tsukasa Sasaki, RT<sup>3)</sup>, Kikutaro Tokairin, MD,  
6 PhD<sup>1)</sup>, Masaki Ito, MD, PhD<sup>1)</sup>, Ken Kazumata, MD, PhD<sup>1)</sup>, and Kiyohiro Houkin, MD, PhD<sup>4)</sup>

7

8 **Affiliations:**

9 1) Department of Neurosurgery, Hokkaido University Graduate School of Medicine, Sapporo, Japan

10 2) Department of Biomedical Sciences, Colorado State University, Colorado, USA

11 3) Department of Radiology, Hokkaido University Hospital, Sapporo, Japan

12 4) Department of Emergent Neurocognition, Faculty of Health Sciences, Hokkaido University,  
13 Sapporo, Japan

14

15 **Correspondence:** Taku Sugiyama, M.D., Ph.D.

16 Department of Neurosurgery, Hokkaido University Graduate School of Medicine, North 15 West 7,  
17 Kita-ku, Sapporo 060-8638, Japan

18 Tel: +81-11-706-5987, Fax: +81-11-708-7737

19 E-mail: [takus1113@med.hokudai.ac.jp](mailto:takus1113@med.hokudai.ac.jp)

20

21 **Disclosure:** The authors have no personal, financial, or institutional interest in any of the drugs,  
22 materials, or devices described in this manuscript.

23 **Sources of funding:** None

24 **Acknowledgements:** We would like to thank Katsuhiko Hieda and Yoko Matsuura for their prominent  
25 assistance of this study.

26

27 **Abstract**

28 **Background:** Adequate surgical planning includes a precise understanding of patient-specific anatomy  
29 and is a necessity for neurosurgeons. Although the use of virtual reality (VR) technology is emerging  
30 in surgical planning and education, few studies have examined the effectiveness of immersive VR  
31 during surgical planning using a modern head-mounted display.

32 **Objective:** To investigate if and how immersive VR aids presurgical discussions of cerebrovascular  
33 surgery.

34 **Methods:** A multiuser immersive VR system, BananaVision™, was developed and used during  
35 presurgical discussions in a prospective patient cohort undergoing cerebrovascular surgery. A  
36 questionnaire/interview was administered to multiple surgeons after the surgeries to evaluate the  
37 effectiveness of the VR system compared to conventional imaging modalities. An objective assessment  
38 of the surgeon's knowledge of patient-specific anatomy was also conducted by rating surgeons' hand-  
39 drawn presurgical illustrations.

40 **Results:** The VR session effectively enhanced surgeons' understanding of patient-specific anatomy in  
41 the majority of cases (83.3%). An objective assessment of surgeons' presurgical illustrations was  
42 consistent with this result. The VR session also effectively improved the decision making process  
43 regarding minor surgical techniques in 61.1% of cases and even aided surgeons in making critical  
44 surgical decisions about cases involving complex and challenging anatomy. The utility of the VR  
45 system was rated significantly higher by trainees than by experts.

46 **Conclusion:** Although rated as more useful by trainees than by experts, immersive 3D VR modeling  
47 increased surgeons' understanding of patient-specific anatomy and improved surgical strategy in  
48 certain cases involving challenging anatomy.

49

50 **Running Title:** Immersive VR for Cerebrovascular Surgery

51 **Keywords:** presurgical planning, simulation, rehearsal, VR, microsurgery, aneurysm, arteriovenous  
52 malformation

53 **Introduction**

54 A precise understanding of microsurgical anatomy is essential for neurosurgeons. Surgical trainees  
55 spend years learning intricate neuroanatomy through case observations and conventional two-  
56 dimensional (2D) representations such as textbook illustrations.<sup>1, 2</sup> Moreover, since neurovascular  
57 anatomy varies significantly from patient to patient, a precise understanding of patient-specific  
58 anatomy is required for cerebrovascular surgery.

59 To date, patient-specific three-dimensional (3D)-printed physical models and virtual reality  
60 (VR) models have been utilized to enhance surgical planning and education as they convey  
61 comprehensive 3D anatomical information to surgeons.<sup>2-4</sup> Advanced simulation models have also been  
62 combined with mixed reality technology.<sup>5</sup> VR models require no extra cost to render models and allow  
63 the user to repeat sessions. Recent study confirmed that VR models have advantages over 3D-printed  
64 models including the ability to zoom, greater image resolution, and greater model durability.<sup>3</sup> Although  
65 3D-printed models allow the use of depth perception and are easier to manipulate,<sup>3</sup> “immersive” VR  
66 environments with modern head-mounted displays (HMDs) provide more convincing stereopsis than  
67 stereoscopic viewing of 2D images, allowing the user to perceive depth in a similar way to 3D-printed  
68 models.<sup>6-9</sup> Gerig et al. have demonstrated that HMDs are superior to 2D screens in terms of depth  
69 perception, regardless of additional artificially recreated depth cues.<sup>6</sup>

70 Although studies examining the utility of VR systems have shown promising results, there have  
71 been few prospective studies utilizing “immersive” VR technology including HMDs. Additionally, VR  
72 systems have not yet been widely integrated into clinical practice, possibly due to concerns about cost  
73 and model construction times.<sup>7-10</sup> Therefore, in this study, we introduce the newly developed VR  
74 modeling program, BananaVision™ (Colorado State University), which has the features of fast  
75 processing time, multiuser functionality, and high usability. We evaluated this system during the  
76 presurgical planning of cerebrovascular surgeries.

77

78

79 **Materials and methods**

80 *VR software (Video 1)*

81 The VR modeling program BananaVision™ was developed at the Department of Biomedical  
82 Sciences, Colorado State University to improve anatomical learning through digitization. This  
83 platform rapidly (<1–2 minutes) generates 3D models using patient’s medical images such as  
84 computed tomography (CT) and magnetic resonance imaging (MRI) scans in VR space. The image  
85 processing procedure is provided in **Supplementary Data 1**.

86 Wearing typical HMDs, the users can maneuver and observe these 3D models in VR space.  
87 Using handheld controllers, the user can easily adjust the images to visualize the skull, brain  
88 parenchyma, and vasculature. The user can change the size of objects, mobilize objects in any  
89 direction, and slice objects on any plane. An “eraser” function enables the user to simulate  
90 craniotomies. Because two surgeons wearing HMDs could simultaneously access the same VR  
91 space, surgical teams could share patient-specific anatomy and surgical views while discussing a  
92 surgical plan or teaching trainees (**Figure 1A**).

93  
94 *Facility and surgeons*

95 The VR system utility for neurosurgery was independently assessed in Hokkaido University  
96 Hospital. Two interconnected personal computers with the BananaVision™ were placed in a  
97 conference room beside the neurosurgical ward in the Hokkaido University Hospital. Surgeons could  
98 freely use this system at any time.

99 Fourteen surgeons participated in this study and were sorted into two groups based on their  
100 level of neurosurgical experience. The trainee group included 10 neurosurgical residents  
101 (postgraduate years 3–6) and one clinical fellow, who had not performed an entire microsurgery  
102 independently. The expert group included 3 board-certified neurosurgeons, who had independently  
103 performed over 500 cerebrovascular microsurgeries.

104

105 ***Study design***

106 This study was approved by the institutional review board of Hokkaido University Hospital (No.  
107 018-0291). Informed consent was obtained from all participants, including surgeons and patients.  
108 The study design is summarized in **Figure 1B**.

109 All patients with cerebrovascular disease (CVD) scheduled or under consideration for  
110 microsurgery between April 2019 and December 2019 were enrolled in this study. Patients with CVD  
111 included patients with cerebral aneurysm, arteriovenous malformation (AVM), and occlusive CVD  
112 such as moyamoya disease (MMD). After the patients were admitted, staff surgeons and attending  
113 residents routinely discussed and created surgical plans using conventional imaging methods,  
114 including MRI, CT, CTA, digital subtraction angiography (DSA), and perfusion imaging. The  
115 attending residents presented the surgical plans during our weekly case meetings prior to surgery, and  
116 the plans were discussed by all neurosurgical staff. Presurgical schematic illustrations were routinely  
117 presented for educational purposes. Attending surgeons then freely decided whether or not to use the  
118 VR system.

119 Outcomes were evaluated 3–4 months after surgery. A favorable outcome was defined as a  
120 modified Rankin Scale of either 0 or 1 or as neurological improvement after surgery relative to the  
121 patient's preoperative state.<sup>11, 12</sup>

122  
123 ***Questionnaire/interview survey***

124 When presurgical planning sessions with the VR system were conducted before surgery, a simple  
125 questionnaire/interview was administered to both primary and assistant surgeons after surgery to  
126 evaluate the effectiveness of the VR sessions. The questionnaire included the following three  
127 questions: **(Q1)** Did you think the VR session effectively increased your understanding of patient-  
128 specific anatomy? **(Q2)** Did you think the VR session effectively enhanced your decision-making  
129 ability regarding a minor surgical technique (e.g., patient's head position, order of the dissection)?  
130 **(Q3)** Did you think the VR session effectively enhanced your decision-making process regarding a

131 critical surgical technique (e.g., surgical approach, range of craniotomy)? The surgeons indicated  
132 their agreement with each question using a 5-point Likert scale ranging from 1 to 5 (“1: Strongly  
133 disagree,” “2: Disagree,” “3: Neutral,” “4: Agree,” and “5: Strongly agree”). The effectiveness of the  
134 VR sessions in each case was represented by the primary surgeon’s evaluation.

135

### 136 ***Objective assessment of surgeons’ anatomical understanding***

137 To objectively evaluate whether the VR session improved surgeons’ understanding of patient-specific  
138 anatomy, we evaluated presurgical schematic illustrations drawn by surgeons who had attended  
139 presurgical planning sessions. Presurgical illustrations were compared with the actual surgical videos,  
140 and the precision of the presurgical illustrations was rated by an independent expert surgeon who did  
141 not take part in performing aneurysm surgeries. They were rated from 0 to 10 in terms of six categories:  
142 (1) projection of aneurysms (0–2 points), (2) location of parent arteries (0–2 points), (3) location of  
143 branch arteries (0–2 points), (4) the surrounding anatomical structure (brain, cranial nerve, or dura  
144 matter/tentorium) that was mainly attached to the aneurysm (0–2 points), (5) other anatomical  
145 structures that were attached to the aneurysm (0 or 1 point), and (6) other anatomical structures in the  
146 surgical view (0 or 1 point). The artistic element was not considered in this rating system. Information  
147 about when (pre- or post-VR) and by whom the presurgical illustration was drawn was blinded to the  
148 rater for evaluation purposes.

149

### 150 ***Statistical analyses***

151 Statistical analyses were performed using the SPSS Statistics® (IBM). The unpaired population rates  
152 between the two groups were compared using the chi-squared or Fisher’s exact tests as appropriate.  
153 The paired ordinal scale data were compared using the Wilcoxon matched-pair signed-rank tests. A p-  
154 value <0.05 was considered statistically significant.

155

156

157 **Results**

158 *Patients' characteristics*

159 During the study period, 38 patients (15 aneurysms, 10 AVMs, and 13 occlusive CVDs) were  
160 candidates for cerebrovascular surgery, and surgeons chose to conduct presurgical VR sessions in 20  
161 (52.6%) patients (12 aneurysms, 7 AVMs, and 1 MMD) (**Figure 2**). In the majority of patients with  
162 occlusive CVD who underwent extracranial-intracranial (EC-IC) bypass, VR sessions were not  
163 conducted because the manner in which EC-IC and indirect bypasses were performed was relatively  
164 standardized. VR was used during presurgical planning for only one patient with occlusive CVD  
165 (MMD), for whom VR was used to identify and preserve the middle meningeal artery. In contrast, VR  
166 was used during presurgical planning for the majority of aneurysm and AVM cases (75% and 70%,  
167 respectively).

168 Of the 20 patients for whom VR was used during presurgical planning, one patient's surgery  
169 for cervical arteriovenous fistula was abandoned because of a complication during the preoperative  
170 embolization session. Of note, one patient with multiple aneurysms was selected for observation rather  
171 than surgery after estimation of their surgical risk during a VR session (**Supplementary Data 2**).  
172 Therefore, the remaining 18 patients underwent surgery and were analyzed. Patient information is  
173 summarized in **Table 1**. The patients comprised 10 men and 8 women aged 22–76 years.

174 All surgeries except one (Case 2) went as planned during the presurgical VR sessions. In the  
175 exceptional case, surgeons planned to clip the patient's aneurysm; however, they found a perforator  
176 strongly attached to the aneurysm intraoperatively and changed the plan to aneurysmal coating. We  
177 observed three surgical complications: two cases of perforator injury (ischemic stroke) and one case  
178 of wound infection. Seventeen patients (94.4%) eventually experienced favorable outcomes.

179

180 *Effectiveness of presurgical VR sessions*

181 Thirty-six responses to questionnaires were obtained from both primary and assistant surgeons across  
182 18 surgical cases. The VR sessions were evaluated as “effective” for increasing surgeons’

183 understanding of patient-specific anatomy in 83.3% of cases (**Figure 3A**) and as “effective” at  
184 enhancing the decision-making process concerning minor surgical techniques in 61.1% of cases  
185 (**Figure 3B**). In some cases (27.8%), VR sessions were also evaluated as “effective” for enhancing the  
186 decision-making process regarding critical surgical techniques (**Figure 3C**). Surgeons never answered  
187 “Disagree” or “Strongly disagree” in response to any question. However, 3 surgeons (21.4%)  
188 experienced mild motion sickness after a VR session.

189 The trainee group was significantly more likely to answer “Strongly agree” in response to Q1  
190 than the expert group ( $p < 0.01$ ) (**Figure 3D**). The trainee group was also more likely to answer  
191 “Strongly agree” or “Agree” in response to both Q2 and Q3 than the expert group ( $p < 0.01$  and  $p = 0.01$ ,  
192 respectively) (**Figure 3E, F**).

193 The cases in which the primary and assistant surgeons agreed with all questions (Q1–3) were  
194 challenging cases involving complex anatomy such as (1) multiple aneurysms (Case 9), (2) a recurrent  
195 basilar artery (BA)-superior cerebellar artery (SCA) aneurysm after coiling (Case 11), (3) a medial  
196 temporal AVM fed by the passing anterior choroidal artery (Case 14), and (4) an occipital AVM fed by  
197 a passing artery (Case 17).

198

### 199 ***Objective assessment using presurgical schematic illustrations***

200 After surgery, 24 total illustrations (14 pre-VR and 10 post-VR) drawn by 9 surgeons (1 expert and 8  
201 trainees) for cases of 9 anterior circulation aneurysms (MCA;  $n = 5$ , internal carotid artery;  $n = 2$ , anterior  
202 communicating artery;  $n = 2$ ) were collected and rated. The score of the illustrations significantly  
203 increased after VR sessions relative to before VR sessions ( $p < 0.05$ ) (**Figure 4**). Interestingly, although  
204 expert surgeon’s scores were higher relative to trainee’s score even before VR sessions, their scores  
205 also increased after VR sessions.

206

### 207 ***Illustrative cases***

#### 208 ***Case 17***

209 A 26-year-old male with a left occipital AVM elected to undergo surgery (**Figure 5A–C**). Preoperative  
210 3D-DSA revealed three feeders from the posterior cerebral artery to the AVM. Of these, one of the  
211 feeders passed through the nidus and supplied blood flow to normal brain parenchyma. Although the  
212 spatial association between the feeders, drainer, and other surroundings was complex, the surgeon  
213 clearly understood the anatomy after using the surgical view during a VR session (**Figure 5D**). The  
214 surgery was performed as planned (**Figure 5E, F**).

215

### 216 *Case 11*

217 A 66-year-old male was admitted to the hospital with a rapidly regrowing BA-SCA aneurysm, which  
218 had previously ruptured and been coiled (**Figure 6A–C**). After discussing treatment options with the  
219 patient, surgeons decided to microsurgically repair the aneurysm. Although the distance between the  
220 intra-aneurysmal coils and the origin of posterior cerebral artery was sufficient to insert a clip, the  
221 distance between the coils and the origin of the SCA was significantly small to insert a clip without  
222 sacrificing the SCA (**Figure 6B**). Therefore, surgeons planned to use clipping in conjunction with a  
223 superficial temporal artery (STA)-SCA bypass. Three surgical approach options were initially  
224 discussed: (1) a transsylvian approach, (2) a subtemporal approach, and (3) a subtemporal approach  
225 with an anterior petrosectomy. A VR session was subsequently conducted involving two experts and  
226 the attending residents. In the VR session, the surgeons determined that the retro-carotid space between  
227 the internal carotid artery and anterior clinoid was significantly narrow to use as a surgical corridor to  
228 the aneurysm in the transsylvian approach (**Figure 6D**). We also simulated a subtemporal approach  
229 combined with an anterior petrosectomy (**Figure 6E**); however, we determined that the space created  
230 by the anterior petrosectomy would not be beneficial to approach both the aneurysmal neck and the  
231 proximal BA. We therefore utilized a standard subtemporal approach and successfully accomplished  
232 both an STA-SCA bypass (**Figure 6F**) and clipping (**Figure 6G, H**).

233

234

235 **Discussion**

236 We prospectively tested the efficacy of a multiuser VR system during daily presurgical planning  
237 sessions for cerebrovascular surgeries. As a result, high utilization rates were observed before  
238 aneurysmal and AVM surgeries possibly due to the strengths of this system such as fast processing  
239 time and high usability.<sup>7, 8</sup> Although the fundamental information volume was similar as the original  
240 CTA data,<sup>13</sup> the surgeons could better understand patient-specific anatomy by observing 3D patient  
241 models in immersive VR space.<sup>14</sup> As expected, the educational effect of VR was greater in the trainee  
242 group; however, even expert surgeons benefited from VR sessions. The multiuser functionality of the  
243 system also enhanced presurgical planning and education because surgeons could discuss and teach  
244 surgical strategy while sharing the same surgical view in VR space.<sup>15</sup>

245 Notably, an objective assessment revealed that VR sessions improved surgeons' anatomical  
246 understanding in both expert and trainee surgeons. This new research method allows for an objective  
247 assessment of surgeons' performance, which is emphasized in the current competency-based education  
248 paradigm.<sup>16, 17</sup>

249  
250 ***Virtual models in cerebrovascular surgery***

251 Recent trends toward increased endovascular techniques and work-hour restrictions are decreasing the  
252 surgical experience of trainees and increasing the difficulty of surgery.<sup>18, 19</sup> Presurgical simulation of  
253 patient-specific anatomy is therefore increasingly required for proper presurgical planning and  
254 education without threatening patients' safety. During aneurysm surgery, the most important strategic  
255 objective is to secure the proximal parent artery while avoiding the risk of aneurysmal rupture.  
256 Therefore, information regarding the projection of the aneurysm and the spatial association between  
257 the proximal and distal arteries are vital considerations when planning these surgeries. VR sessions  
258 could have aided the decision-making process regarding surgical approaches. Similar benefits were  
259 also observed in AVM surgeries, specifically regarding the strategy for securing the main feeding artery  
260 while avoiding the occlusion of the passing artery and draining vein before nidus resection.<sup>20</sup>

261

262 ***Limitations***

263 This study has several limitations. First, it included a limited number of cases and could not  
264 demonstrate a direct correlation with clinical outcomes such as reduction of operating time or blood  
265 loss and neurological improvement. Second, the self-selection of the user to participate in VR session  
266 could introduce a bias in the result during subjective assessment since he/she was most likely to rate  
267 the session favorably.

268 The largest shortcoming of the BananaVision™ was likely its inability to fuse multimodal  
269 images. Several investigators have successfully built more realistic 3D graphics that include  
270 information regarding cranial nerves and the brain surface.<sup>21, 22</sup> Another limitation is the current  
271 availability of other VR platforms.<sup>15, 23-25</sup> Integrating VR systems with other devices such as navigation  
272 systems<sup>26-28</sup> and augmented reality technologies<sup>29-31</sup> will likely produce the next generation of surgical  
273 planning tools. Additional technologies such as procedural simulation with haptic feedback and/or  
274 tissue deformation would also be another direction.<sup>23-25, 32</sup> However, since these advanced systems will  
275 be expensive and time consuming, only disease- or purpose-specific usage of advanced technologies  
276 should be considered in the future.<sup>7, 33</sup>

277

278 **Conclusion**

279 Although the educational impact of VR sessions was greater in surgical trainees than expert surgeons,  
280 3D modeling of patient-specific imaging effectively increased surgeons' understanding of patient-  
281 specific anatomy and helped determine surgical strategy in certain cases involving complex and  
282 challenging anatomy.

283

284 **References**

- 285 1. Chan S, Conti F, Salisbury K, Blevins NH. Virtual reality simulation in neurosurgery:  
286 technologies and evolution. *Neurosurgery*. Jan 2013;72 Suppl 1:154-164.
- 287 2. Tomlinson SB, Hendricks BK, Cohen-Gadol A. Immersive Three-Dimensional Modeling and  
288 Virtual Reality for Enhanced Visualization of Operative Neurosurgical Anatomy. *World*  
289 *Neurosurg*. Nov 2019;131:313-320.
- 290 3. Bairamian D, Liu S, Eftekhari B. Virtual Reality Angiogram vs 3-Dimensional Printed  
291 Angiogram as an Educational tool-A Comparative Study. *Neurosurgery*. Aug 1 2019;85(2):E343-  
292 E349.
- 293 4. Randazzo M, Pisapia JM, Singh N, Thawani JP. 3D printing in neurosurgery: A systematic  
294 review. *Surg Neurol Int*. 2016;7(Suppl 33):S801-S809.
- 295 5. Weinstock P, Rehder R, Prabhu SP, Forbes PW, Roussin CJ, Cohen AR. Creation of a novel  
296 simulator for minimally invasive neurosurgery: fusion of 3D printing and special effects. *J*  
297 *Neurosurg Pediatr*. Jul 2017;20(1):1-9.
- 298 6. Gerig N, Mayo J, Baur K, Wittmann F, Riener R, Wolf P. Missing depth cues in virtual reality  
299 limit performance and quality of three dimensional reaching movements. *PLoS One*.  
300 2018;13(1):e0189275.
- 301 7. Kin T, Nakatomi H, Shono N, et al. Neurosurgical Virtual Reality Simulation for Brain Tumor  
302 Using High-definition Computer Graphics: A Review of the Literature. *Neurol Med Chir (Tokyo)*.  
303 Oct 15 2017;57(10):513-520.
- 304 8. Li Y, Zhao Y, Zhang J, et al. Low-Cost Interactive Image-Based Virtual Endoscopy for the  
305 Diagnosis and Surgical Planning of Suprasellar Arachnoid Cysts. *World Neurosurg*. Apr  
306 2016;88:76-82.
- 307 9. Mashiko T, Otani K, Kawano R, et al. Development of three-dimensional hollow elastic model for  
308 cerebral aneurysm clipping simulation enabling rapid and low cost prototyping. *World*  
309 *Neurosurg*. Mar 2015;83(3):351-361.
- 310 10. Coburn JQ, Freeman I, Salmon JL. A Review of the Capabilities of Current Low-Cost Virtual  
311 Reality Technology and Its Potential to Enhance the Design Process. *Journal of Computing and*  
312 *Information Science in Engineering*. 2017;17(3).
- 313 11. Sugiyama T, Nakayama N, Ushikoshi S, et al. Complication rate, cure rate, and long-term  
314 outcomes of microsurgery for intracranial dural arteriovenous fistulae: a multicenter series and  
315 systematic review. *Neurosurg Rev*. Jan 2 2020.
- 316 12. Darsaut TE, Findlay JM, Magro E, et al. Surgical clipping or endovascular coiling for unruptured  
317 intracranial aneurysms: a pragmatic randomised trial. *J Neurol Neurosurg Psychiatry*. Aug  
318 2017;88(8):663-668.
- 319 13. Stewart N, Lock G, Hopcraft A, Kanesarajah J, Coucher J. Stereoscopy in diagnostic radiology  
320 and procedure planning: does stereoscopic assessment of volume-rendered CT angiograms lead to  
321 more accurate characterisation of cerebral aneurysms compared with traditional monoscopic

- 322 viewing? *J Med Imaging Radiat Oncol.* Apr 2014;58(2):172-182.
- 323 14. Wake N, Wysock JS, Bjurlin MA, Chandarana H, Huang WC. "Pin the Tumor on the Kidney:" An  
324 Evaluation of How Surgeons Translate CT and MRI Data to 3D Models. *Urology.* Sep  
325 2019;131:255-261.
- 326 15. Kockro RA, Stadie A, Schwandt E, et al. A collaborative virtual reality environment for  
327 neurosurgical planning and training. *Neurosurgery.* Nov 2007;61(5 Suppl 2):379-391; discussion  
328 391.
- 329 16. Sugiyama T, Lama S, Gan LS, Maddahi Y, Zareinia K, Sutherland GR. Forces of Tool-Tissue  
330 Interaction to Assess Surgical Skill Level. *JAMA Surg.* Mar 1 2018;153(3):234-242.
- 331 17. Sugiyama T, Nakamura T, Ito Y, et al. A Pilot Study on Measuring Tissue Motion During Carotid  
332 Surgery Using Video-Based Analyses for the Objective Assessment of Surgical Performance.  
333 *World J Surg.* Sep 2019;43(9):2309-2319.
- 334 18. Kalani MYS, Wanebo JE, Martirosyan NL, Nakaji P, Zabramski JM, Spetzler RF. A raised bar  
335 for aneurysm surgery in the endovascular era. *J Neurosurg.* May 2017;126(5):1731-1739.
- 336 19. Quillin RC, 3rd, Cortez AR, Pritts TA, Hanseman DJ, Edwards MJ, Davis BR. Operative  
337 Variability Among Residents Has Increased Since Implementation of the 80-Hour Workweek. *J*  
338 *Am Coll Surg.* Jun 2016;222(6):1201-1210.
- 339 20. Sugiyama T, Gan LS, Zareinia K, Lama S, Sutherland GR. Tool-Tissue Interaction Forces in  
340 Brain Arteriovenous Malformation Surgery. *World Neurosurg.* Jun 2017;102:221-228.
- 341 21. Oishi M, Fukuda M, Hiraishi T, Yajima N, Sato Y, Fujii Y. Interactive virtual simulation using a  
342 3D computer graphics model for microvascular decompression surgery. *J Neurosurg.* Sep  
343 2012;117(3):555-565.
- 344 22. Kin T, Nakatomi H, Shojima M, et al. A new strategic neurosurgical planning tool for brainstem  
345 cavernous malformations using interactive computer graphics with multimodal fusion images. *J*  
346 *Neurosurg.* Jul 2012;117(1):78-88.
- 347 23. Alaraj A, Luciano CJ, Bailey DP, et al. Virtual reality cerebral aneurysm clipping simulation  
348 with real-time haptic feedback. *Neurosurgery.* Mar 2015;11 Suppl 2:52-58.
- 349 24. Gmeiner M, Dirnberger J, Fenz W, et al. Virtual Cerebral Aneurysm Clipping with Real-Time  
350 Haptic Force Feedback in Neurosurgical Education. *World Neurosurg.* Apr 2018;112:e313-e323.
- 351 25. Wong GK, Zhu CX, Ahuja AT, Poon WS. Craniotomy and clipping of intracranial aneurysm in a  
352 stereoscopic virtual reality environment. *Neurosurgery.* Sep 2007;61(3):564-568; discussion 568-  
353 569.
- 354 26. Kockro RA, Reisch R, Serra L, Goh LC, Lee E, Stadie AT. Image-guided neurosurgery with 3-  
355 dimensional multimodal imaging data on a stereoscopic monitor. *Neurosurgery.* Jan 2013;72  
356 Suppl 1:78-88.
- 357 27. Mert A, Buehler K, Sutherland GR, et al. Brain tumor surgery with 3-dimensional surface  
358 navigation. *Neurosurgery.* Dec 2012;71(2 Suppl Operative):ons286-294; discussion ons294-285.
- 359 28. Schulze F, Buhler K, Neubauer A, Kanitsar A, Holton L, Wolfsberger S. Intra-operative virtual

- 360 endoscopy for image guided endonasal transsphenoidal pituitary surgery. *Int J Comput Assist*  
361 *Radiol Surg*. Mar 2010;5(2):143-154.
- 362 **29.** Besharati Tabrizi L, Mahvash M. Augmented reality-guided neurosurgery: accuracy and  
363 intraoperative application of an image projection technique. *J Neurosurg*. Jul 2015;123(1):206-  
364 211.
- 365 **30.** Incekara F, Smits M, Dirven C, Vincent A. Clinical Feasibility of a Wearable Mixed-Reality  
366 Device in Neurosurgery. *World Neurosurg*. Oct 2018;118:e422-e427.
- 367 **31.** Lovo EE, Quintana JC, Puebla MC, et al. A novel, inexpensive method of image coregistration for  
368 applications in image-guided surgery using augmented reality. *Neurosurgery*. Apr 2007;60(4  
369 Suppl 2):366-371; discussion 371-362.
- 370 **32.** Kockro RA, Killeen T, Ayyad A, et al. Aneurysm Surgery with Preoperative Three-Dimensional  
371 Planning in a Virtual Reality Environment: Technique and Outcome Analysis. *World Neurosurg*.  
372 Dec 2016;96:489-499.
- 373 **33.** McClelland TJ, Ford K, Dagash H, Lander A, Lakhoo K. Low-fidelity Paediatric Surgical  
374 Simulation: Description of Models in Low-Resource Settings. *World J Surg*. May 2019;43(5):1193-  
375 1197.
- 376
- 377
- 378
- 379

380 **Figure Legends**

381 **Fig. 1**

382 **A:** Virtual reality (VR) station with BananaVision™ installed together with hand controller and a  
383 head-mounted display. Collaborative case discussions were conducted while sharing the same  
384 immersive VR models.

385 **B:** Study design.

386

387 **Fig. 2** Flow chart of study participants during study period.

388 **Abbreviations** AN; aneurysm, AVM; arterio-venous malformation, AVF; arteriovenous fistula, CVD;  
389 cerebrovascular disease, CTA; computed tomography angiography, MMD; moyamoya disease

390

391 **Fig. 3**

392 **A-C:** Bar graphs of primary surgeons' responses to the three questions for each case.

393 **D-F:** Bar graphs of the rates at which all surgeons answered "Strongly agree" or "Agree" to the three  
394 questions. \*p<0.05, \*\*p<0.01.

395

396 **Fig. 4** Presurgical schematic illustrations and scores (Case 5)

397 **A:** A conventional three-dimensional (3D) computed tomography angiogram (CTA).

398 **B:** A VR session. The users could view 3D models at any angle, simulate craniotomies, and experience  
399 surgical approaches in immersive VR space.

400 **C-E:** Intraoperative photographs of the secured parent artery (**E**), the exposed aneurysm (**F**), and the  
401 surgical site after clipping (**G**).

402 **F, G:** An example of presurgical illustrations before and after VR sessions. The scores tended to be  
403 higher in experts than in trainees, and all scores increased after VR sessions. \*p<0.05

404

405

406 **Fig. 5** An arteriovenous malformation (AVM) case (Case 17)

407 An AVM was located in the left medial occipital lobe (A). A conventional digital subtraction  
408 angiogram (DSA, B) and rotational DSA (C) showing three feeders and one drainer (asterisk). One  
409 of the feeders (F-2) was passing through the nidus and was supplying blood flow to normal brain  
410 parenchyma. Through VR models, surgeons could experience an operative 3D view and understand  
411 the spatial relationships between the nidus, drainer, and feeders (D). Intraoperative photographs  
412 showing that the feeders and drainer were precisely identified (E), and the nidus was removed as  
413 planned without injuring either the passing artery or drainer (F).

414

415 **Fig. 6** A case of a regrowing BA-SCA aneurysm after coiling (Case 11).

416 A conventional DSA (A), rotational DSA (B), and 3D-CTA (C) showing regrowth of a BA-SCA  
417 aneurysm after coiling. The VR simulation of a transsylvian approach (D) showed that the postero-  
418 inferior space of the internal carotid artery was too narrow to approach the aneurysm (asterisk). The  
419 simulation of a subtemporal approach (E) combined with an anterior petrosectomy (arrow) showed  
420 that the surgical corridor through the anterior petrosectomy was not beneficial for approaching the  
421 aneurysm or for securing the proximal basilar artery. Intraoperative photographs (F-H) showed that  
422 the SCA was anastomosed with the STA. The aneurysm was clipped as planned using the  
423 subtemporal approach.

424 **Abbreviations** PComA; posterior communicating artery, PCA; posterior cerebral artery, SCA;  
425 superior cerebellar artery, ACA; anterior cerebral artery, ICA; internal carotid artery, MCA; middle  
426 cerebral artery, CP; clinoid process, BA; basilar artery, STA; superficial temporal artery, CN IV; fourth  
427 nerve

428

429

430 **Supplemental Digital Content**

431 **Supplementary Data 1** BananaVision™ and image processing


432 **Supplementary Data 2** A case of multiple aneurysms who selected for observation

433

434


435 **Video Legends**

436 **Video 1** Multiuser immersive virtual reality system



**B**

**Surgery**



**Presurgical planning session**

with  
conventional  
imaging


with  
VR system  
(optional)

**Postsurgical evaluation**


- Questionnaire/interview survey
- Rating of presurgical illustration

**Outcome  
evaluation**




**Fig. 1**




**Fig. 2**




**Fig. 3**



**Fig. 4**



**Fig. 5**



**Fig. 6**

Case No.	Age /sex	Disease	Location	Feature	Surgical plan	Results		Questionnaire		
						Surgery	Complication	Q1	Q2	Q3
1	71/F	AN	Rt. ICA-OpthA	Unrup./6 mm/superior proj.	Clipping via TSA with AC	Clipped as planned		4	4	3
2	62/M	AN	Rt. ICA-top	Unrup./5 mm/posterior proj.	Clipping via TSA	Coated d/t perforator adhesion		5	4	3
3	69/F	AN	Lt. ICA-top	Unrup. (growing)/4 mm/ posterior proj.	Clipping via TSA	Clipped as planned		4	4	3
4	76/F	AN	Lt. pMCA-ATA	Unrup. (growing)/4 mm/ inferior proj.	Clipping via TSA	Clipped as planned		4	3	3
5	67/F	AN	Lt. MCA	Unrup./9 mm/lateral proj.	Clipping via TSA	Clipped as planned		4	4	3
6	43/F	AN	Lt. MCA	Unrup. (growing)/4 mm/ antero-medial proj.	Clipping via TSA	Clipped as planned		4	3	3
7	67/F	AN	Rt. MCA	Unrup./4 mm/supero-lateral proj.	Clipping via TSA	Clipped as planned		5	4	3
8	41/F	AN	AComA	Unrup./4 mm/supero-posterior proj.	Clipping via IHA	Clipped as planned		5	4	3
9	70/M	Multiple ANs	Rt. MCA	Unrup./6 mm/inferior proj.	Clipping via rt. TSA	Clipped as planned		5	4	5
			Rt. ICA-AChoA	Unrup./5 mm/posterior proj.	Clipping via rt. TSA	Clipped as planned				
			AComA	Unrup./3 mm/supero-posterior proj.	Probably could not be clipped via rt. TSA	Could not be clipped				
			Lt. pACA	Unrup./2 mm/superior proj.	Probably could be clipped via rt. TSA	Clipped as expected				
10	66/M	AN	Lt. VA	Symptomatic/fusiform/thrombosed/34 mm	Trapping via LSA	Trapped as planned		5	3	4
11	62/M	AN	Rt. BA-SCA	Regrowth after coiling/10 mm/postero-lateral proj.	STA-SCA bypass + Clipping via SubT	Clipped as planned	Perforator injury (mRS;1)	5	5	5
12	58/M	AVM/dAVF	Lt. frontal	Post-embolization and radiation/S-M III/Borden III	Removal via frontal craniotomy	Removed as planned		3	3	3
13	30/M	AVM	Lt. temporal	Unrup. (seizure)/S-M IV	Removal via fronto-temporal craniotomy	Removed as planned		5	3	3
14	32/M	AVM	Rt. medial temporal	Ruptured/S-M III	Removal via TSA	Removed as planned	Perforator injury (mRS; 2)	4	4	4
15	22/M	AVM	Lt. parietal	Unrup./S-M III	Removal via parietal craniotomy	Removed as planned		5	5	3
16	26/M	AVM	Lt. parietal	Ruptured/S-M II	Removal via parietal craniotomy	Removed as planned		3	3	3
17	26/M	AVM	Lt. occipital	Unrup./S-M III	Removal via occipital craniotomy	Removed as planned	Infection	5	5	4
18	46/F	MMD	Lt. side	Hemorrhagic onset (contralateral side)/Suzuki IV	STA-MCA bypass + indirect bypass	Revascularized as planned		3	3	3

**Table 1. Summary of patients included in the study**

**Abbreviation** F; female, AN; aneurysm, Rt.; right, ICA; internal carotid artery, OpthA; ophthalmic artery, Unrup.; unruptured, proj.; projection, TSA; transsylvian approach, AC; anterior clinoidectomy, M; male, d/t; due to, pMCA: proximal middle cerebral artery, ATA; anterior temporal artery, MCA; middle cerebral artery, AComA; anterior communicating artery, IHA; interhemispheric approach, AChoA; anterior choroidal artery, pACA; proximal anterior cerebral artery, VA; vertebral artery, LSA; lateral suboccipital approach, BA; basilar artery, SCA; superior cerebellar artery, STA; superficial temporal artery, SubT; subtemporal approach, mRS; modified Rankin Scale, AVM; arteriovenous malformation, dAVF; dural arteriovenous fistula, S-M; Spetzler-Martin grade, Borden; Borden type, MMD; moyamoya disease, Suzuki; Suzuki stage