



HOKKAIDO UNIVERSITY

Title	Responses of ectomycorrhizal diversity of larch and its hybrid seedlings and saplings to elevated CO ₂ , O ₃ , and high nitrogen loading
Author(s)	QU, Laiye; WANG, Xiaona; MAO, Qiaozhi et al.
Citation	Eurasian Journal of Forest Research, 22, 23-27
Issue Date	2022
DOI	https://doi.org/10.14943/EJFR.22.23
Doc URL	https://hdl.handle.net/2115/84968
Type	departmental bulletin paper
File Information	6)EJFR-Qu-Koike_final2.pdf



Responses of ectomycorrhizal diversity of larch and its hybrid seedlings and saplings to elevated CO₂, O₃, and high nitrogen loading

QU Laiye¹, WANG Xiaona^{2,3}, MAO Qiaozhi^{2,4}, AGATHOKLEOUS Evgenios^{2,5},
CHOI DongSu^{2,6}, TAMAI Yutaka², WATANABE Toshihiro², KOIKE Takayoshi^{1,2*}

¹ Research Center for Eco-Environmental Science, CAS, Beijing 100085 and
University of Chinese Academy of Science, Beijing 100049, China

² School of Agriculture, Hokkaido University, Sapporo 060-8589, Japan

³ Present address: Hebei Agricultural University, Hebei 071001, China,

⁴ Present address: College of Resources and Environment, Southwest University,
Chongqing 400715, China,

⁵ Present address: Department of Ecology, School of Applied Meteorology,
Nanjing University of Information Science and Technology, Nanjing 210044, China,

⁶ Present address: Institute of Agriculture, Tokyo University of Agriculture and Technology,
Fuchu-Tokyo 183-8509, Japan

Abstract

The genus *Larix* (larch) is broadly distributed in northern Eurasia, including Japan and Korea. Larch species are a typical afforestation tree in northeastern China, Russia, Korea, and Japan. Larch and red pine seedlings inoculated with *Pisolithus arhizus* (earth-ball fungus) have increased photosynthetic rates as well as being acid-soil tolerant. Infecting *P. arhizus* with a variety of ectomycorrhizae (ECM) accelerated these functions. Under elevated CO₂ (about 700 ppm) conditions, larch and pine seedlings infected with ECM showed no down-regulation in photosynthesis and displayed higher growth rates than those grown in ambient CO₂ (380 ppm) conditions. Simulated nitrogen (N) deposition (50 kg N hr⁻¹yr⁻¹) increased photosynthesis and growth of F₁ hybrid larch for the initial two years. There were no significant effects for the next three to eight years on Brown Forest Soil. Among the three species (Japanese, Dahurian and F₁ hybrid larch), the dominant ECM species preferred N. Elevated O₃ decreased growth, ECM infection rates, and ECM species diversity; however, elevated CO₂ moderated or increased them in hybrid larch F₁. After elevated CO₂ and O₃ fumigation on F₁ seedlings, we found *Suillus* sp., a larch specialist ECM in F₁, which means these ECM will be effective materials for larch plantations.

Key words: Elevated CO₂, Ozone, nitrogen deposition, larch, hybrid larch, ectomycorrhiza (ECM).

Introduction

The larch is a typical ectomycorrhizal (ECM) tree species and dominant deciduous conifer in the northern hemisphere, especially dominating permafrost ecosystems (Osawa *et al.* 2010, Qu *et al.* 2010, 2014; Figure 1). Some larch species are representative afforestation species in northeast China, Russian Far East, Korea, and Japan. Most larches can tolerate cold




Figure 1. Japanese larch seedlings inoculated with Ectomycorrhiza; *Pisolithus arhizus* (A) and No-inoculated as Control (B). Diameter of the coin in photo is 2.3mm. (Qu L,

and late frost (Kurahashi 1988), thus attaining significant biomass with a high growth rate in cold regions.

Due to these good growth traits, larch (Japanese larch) was exported to Europe as a pollen resource. This high growth results from its high photosynthetic rate and unique arrangement of two different

types of needles, i.e., short-shoot and long-shoot needles (Kitaoka *et al.* 2020). For breeding practices, paternal inheritance in chloroplast DNA in larch species (Szmidt *et al.* 1987) has surely been contributing to improving photosynthesis via pollen father.

From 1999, the Chinese government decided to reforest farmland and employed larch in northeast China to increase forests (Zhang *et al.* 2000). In northern Japan, hybrid larch F₁ (*Larix gemelinii* var. *japonica* × *L. kaempferi*; hereafter F₁) was produced to improve tolerance to grazing damage by voles and stem straightness, thus enhancing timber quality (Hokkaido 2007; Ryu *et al.* 2009). Nowadays propagation methods have improved, i.e., stem cutting, so the F₁ is becoming a principal afforestation tree species in Japan (Kita *et al.* 2018).

Since the 1960s, with the rapid economic development, air pollutants (NO_x, SO_x, Ozone: O₃) and increasing CO₂ have impacted forest health and vigor. SO_x pollutants were reduced by desulfurization equipment during the 1970s; however, NO_x has hardly decreased because it is mainly produced by diesel cars (Izuta 2017). NO₂ is transformed to O₃ and NO under UV radiation (Gregg *et al.* 2003). In addition,

atmospheric CO₂ concentration [CO₂] has increased since the Industrial Revolution and has reached around 418 ppm at the current rate of 2.2 ppm yr⁻¹. The photosynthetic rate is usually reduced by elevated CO₂ under root restricting conditions. In addition, lockdowns applied amid the Covid-19 pandemic may decrease the




Figure 2. Yearly trend of NO₂ emission of 3 regions. Data are cited from Akimoto (2003) and Kurokawa and Ohara (2020). 1990~: EU-LRTAP convention; USA-EPA Air Pollutant Emission and NIES.

rate to about 7 ppm yr⁻¹ between 2019 and 2020.

Sufficient CO₂ and adequate N are regarded as the productive atmospheric environment for forest trees. We summarize the effects of changing environment (CO₂, N deposition, and O₃) on the growth of larch and larch-ECM interactions.

Elevated CO₂

In many cases, we found down-regulation of photosynthesis, even in a FACE system (Free Air CO₂ Environment; Koike *et al.* 2015). We expected ECM to act as a carbon sink and moderate down-regulation in photosynthesis, Japanese red pine seedlings inoculated with a kind of commercial ECM (*Pisolithus arhizus*) showed no down-regulation (Choi *et al.* 2005). The same trend was expected in larch species (Qu *et al.* 2004). Hybrid larch F₁ was planted in the FACE for three years and its aboveground tended to fall down by wind because of increased above-ground biomass (Watanabe *et al.* 2013). After five years of CO₂ fumigation in FACE, Japanese larch decreased biomass allocation to branches and increased it by about 20% in the stem compared with ambient CO₂ (Watanabe *et al.* 2016). Birch (*Betula platyphylla* var. *japonica*) and kalopanax (*Kalopanax septemlobus*) allocate about 10% less biomass to their stems. However, no clear anatomical structure of them changed with elevated CO₂.

Elevated O₃

Ozone levels have been increasing around the northern hemisphere in the past several decades (Sitch *et al.* 2007). With the GIS method and OTC experiment, Watanabe *et al.* (2010) predicted that the growth of Japanese larch would be reduced by elevated O₃ around the Kanto plain, however, it will be not as large in northern Japan. As mentioned above, O₃ concentration is

generally high in the suburbs due to the reaction of NO₂ with UV radiation (exhaust gas: NO from diesel cars plus UV=NO₂) (Gregg *et al.* 2003). This cycling of NO_x under UV can lead to O₃ in suburban green areas around big cities.

According to recent NO₂ trend in Asia (Kurokawa and Ohara 2020), we should pay attention to rapid an increase of NO₂ emission as a precursor of O₃. Based on the statistics of EU (LRTAP) and US (EPA), the NO₂ emission from Asia reached to about 43 Tg NO₂ yr⁻¹ which is 4 times larger than that from EU or UAS (Figure 2). Therefore, we should know this evidence and try to give O₃ tolerance to larch plantation as suggested by




Figure 3. Growth of F₁ under different O₃ (Modified from Kita *et al.* 2018)

Watanabe *et al.* (2006).

Based on screening experiments using open top chambers (OTCs), Yamaguchi *et al.* (2011) summarized the O₃ sensitivity of 18 tree species with use of OTC and potted seedlings in Japan, and among them, the Japanese larch showed a moderate sensitivity. What about the F₁ under elevated O₃?

The specific difference in O₃ sensitivity was examined between Japanese and F₁ seedlings planted on the ground of OTCs (<5, 25, 45, and 80 ppb). The growth of both larches was significantly suppressed by 80 ppb (Figure 3; Sugai *et al.* 2017). The biomass of F₁ seedlings decreased under 25 ppb, compared to <5 ppb, but due to its heterosis it maintained a similar biomass with Japanese larch seedlings in elevated O₃ treatments. The biomass of Japanese larch decreased at 80 ppb, but it was lower at low O₃ (<5 ppb) compared to 25 ppb. This phenomenon is regarded as “hormesis” (Agathokleous 2018). Namely, mild stresses often accelerate growth and some growth function. As almost all practical production of larch seedlings is done in the suburbs, we tried to use ethylenediurea (EDU) to moderate the adverse effects of elevated O₃ on larch seedlings (Agathokleous *et al.* 2020, 2021). An effective concentration of EDU is

400 mg EDU L⁻¹ applied as soil drench it protects both Japanese and F₁ seedlings against toxicities induced by exposure to elevated O₃ for up to 3-4 years.

Nitrogen deposition and elevated O₃

The combined effects of N and elevated O₃ were studied in seedlings of two broadly distributed species, Siebold's beech and Japanese larch, in OTCs. The beech is classified as highly sensitive to O₃ (Yamaguchi *et al.* 2011). With increasing N (NH₄NO₃), O₃ sensitivity of the beech increased in terms of Accumulated exposure Over Threshold of 40 ppb O₃ (AOT40) (Yamaguchi *et al.* 2007). In contrast, O₃ sensitivity of Japanese larch decreased with increasing N up to 50 kg N ha⁻¹ yr⁻¹ (Watanabe *et al.* 2006) as suggested in above-section. However, F₁ had slightly increased O₃ sensitivity with 50 kg N ha⁻¹ yr⁻¹ under free-air O₃ exposure (Dong-Gyu *et al.* 2016), which may be attributed to decreased (or too much accelerated) leaf life-span with N application (Sugai *et al.* 2019).

In general, phosphorous (P) is the second most important nutrient after N; for the growth of F₁ an adequate supply P and N is required. Magnesium (Mg) was the limiting element in the nursery of Hokkaido University (Mao *et al.* 2014). In this edaphic condition, we examined the effects of N deposition (NH₄NO₃) on the growth of hybrid larch F₁ for eight years. However, except for N application to F₁ by the 2nd year, almost no difference in growth of F₁ was found between the N treatment and the control (=no N application). Based on DNA analysis of the ITS region in symbiotic ECM (Wang *et al.* 2015), most of them infecting Japanese larch, Dahurian larch, and F₁ were nitrogenous species (Wang *et al.* 2018).

Elevated CO₂ and O₃

Plants usually close stomata under elevated CO₂, and consequently, plants can reduce the absorption of O₃. We examined the effects of elevated O₃ (80 ppb) on the growth and ECM infection and diversity of F₁ seedlings under elevated CO₂ in OTCs (Wang *et al.* 2015). Under elevated O₃, ECM infection rate and species diversity were reduced; however, these trends were moderated by elevated CO₂ (600 ppm) (Figure 4). Early successional types of ECMs were only found at ambient and elevated CO₂. However, larch specialist *Suillus* sp. dominated at elevated O₃. This evidence suggests that *Suillus* sp. may support the growth of the host plant: F₁.

As Qu *et al.* (2004) suggested the photosynthetic rate in larch species infected with multiple ECM species was higher than that with a single ECM species (Figure 2). This phenomenon is recognized as follow: most ECM activity depends on soil pH; some ECM prefer low pH




Figure 4. Combination effect of elevated O₃ and CO₂ on ECM infection and diversity (*H'*). Modified from Wang *et al.* (2015) and the *H'* is the Shannon index showing biodiversity. The larger *H'*, the higher biodiversity is. Illustration of fruit body indicates *Suillus* sp.

but some require neutral or high pH (6–8), such as *Rhizopogon rubescens* (Furusawa *et al.* 2020). Aluminum (Al) is released below pH<4.5 and inhibits root growth. These are species-specific traits, and multiple ECM infections may benefit the host plants.

Conclusion

Larch trees planted in China in 1999 to return farmland to forests should be thinned. In northern Japan, Japanese larch is intensively used after considerable improvements in timber utilization. After harvest, we should make plantations with container-grown seedlings to save labor and attain high plantation efficiency. We also expect a new CO₂ sink when planting a new plantation. If we make new plantations with hybrid larch F₁, we should ensure they do not increase N deposition under elevated O₃. To make planting stocks, we should inoculate larch seedlings with ECM (*Suillus* sp.); larch specialist for increased tolerance against environmental stress.

Acknowledgements:

We deeply appreciate the staff of Hokkaido Forestry Research Institute for their continuous support of our researches. Financial support in part by JST (No. JPMJSC18HB; representative researcher, M. Watanabe of TUAT and T. Watanabe of HU) and by the National Key Research and Development Program of China (2017YFE0127700; LY. Qu) are acknowledged.

Reference

- Agathokleous E, Kitao M, Wang XN, Mao QZ, Harayama H, Manning WJ, Koike T. (2020) Ethylenediurea (EDU) effects on Japanese larch: an one growing season experiment with simulated regenerating communities and a four growing season application to individual saplings. *Journal of Forestry Research*, doi.org/10.1007/s11676-020-01223-6
- Agathokleous, E., Kitao, M., Koike, T. (2021) Ethylenediurea (EDU) effects on hybrid larch saplings exposed to ambient or elevated ozone over three growing seasons. *Journal of Forestry Research*, 10.1007/s11676-021-01352-6
- Choi, D.S., Quoreshi, A.M., Maruyama, Y., Jin, H.O., Koike, T. (2005) Effect of ectomycorrhizal infection on growth and photosynthetic characteristics of *Pinus densiflora* seedlings grown under elevated CO₂ concentrations. *Photosynthetica* 43: 223-229.
- Choi, D.S., Jin, H.O., Chung, D. J. Sasa, K., Koike, T. (2008) Growth and physiological activity in *Larix kaempferi* seedlings inoculated with ectomycorrhizae affected by soil acidification. *Trees - Structure and Function* 22:729-735.
- Dong-Gyu Kam, Shi, C., Watanabe, M., Kita, K., Satoh, F., Koike, T. (2015) Growth of Japanese and hybrid larch seedlings grown under free-air O₃ fumigation—an initial assessment of the effects of adequate and excessive nitrogen. *Journal of Agricultural Meteorology* 71: 239-244.
- Izuta, T. (2017) *Air Pollution Impacts on Plants in East Asia*, Springer-Tokyo. Pp.496.
- Kayama, M., Qu, L.Y., Koike, T. (2015) Elements and ectomycorrhizal symbiosis affecting the growth of Japanese larch seedlings regenerated on slopes of an active volcano in northern Japan. *Trees* 29: 1567-1579.
- Kita, T., Sugai, T., Fujita, S., Koike, T. (2018) Breeding effort on hybrid larch F₁ and its responses to environmental stresses. *Forest genetics and Tree Breeding* 7: 107-114.
- Kitaoka, S, Qu, LY., Watanabe, Y, Watanabe, M, Watanabe, T, Koike, T. (2020) Heterophyllous shoots of Japanese larch trees: the seasonal and yearly variation in CO₂ assimilation capacity of the canopy top with changing environment. *Plants* 9 (10), <https://doi.org/10.3390/plants9101278>
- Koike, T., Mao, QZ., Inada, N., Kawaguchi, K., Hoshika, Y., Kita, K. and Watanabe, M. (2012) Growth and photosynthetic responses of cuttings of a hybrid larch (*Larix gmelinii* var. *japonica* x *L. kaempferi*) to elevated ozone and/or carbon dioxide. *Asian Journal of Atmospheric Environment* 6:104-110.
- Kurokawa, J., Ohara, T. (2020) Long-term historical trends in air pollutant emissions in Asia: Regional emission inventory in Asia (REAS) version 3. *Atmospheric Chemistry and Physics* 20: 12761-12793.
- Kurahashi, A. (1988) Improvement of larches by species hybridization, *Bulletin of The Tokyo Univ. Forests*, 79: 1-96 (in Japanese with English summary).
- Mao, QZ., Watanabe, M., Makoto, K., Kita, K., Koike, T. (2014) High nitrogen deposition may enhance growth of the new hybrid larch F₁ growing at two phosphorus levels. *Landscape and Ecological Engineering* 10:1-8.
- Mochizukia, T., Watanabe, M., Koike, T., Tani, A. (2017) Monoterpene emissions from needles of hybrid larch F₁ (*Larix gmelinii* var. *japonica* × *Larix kaempferi*) grown under elevated carbon dioxide and ozone. *Atmospheric Environment* 148: 197-202.
- Qu. LY., Quoreshi, A.M., Koike, T. (2003) Root growth characteristics, biomass, and nutrient dynamics of seedlings of two larch species raised under different fertilization regimes. *Plant and Soil* 255: 293-302.
- Qu, LY., Shinano, T., Quoreshi, A.M., Tamai, Y., Osaki, M., Koike, T. (2004) Allocation of ¹⁴C-Carbon in infected with ectomycorrhizae of two species of larch seedlings. *Tree Physiology* 24: 1369–1376.
- Qu, LY., Kitaoka, S., Kuromaru, M., Osaki, M., Sasa, K., Koike, T. (2009) Root-shoot communication of the seedlings of Japanese larch and its hybrid grown in different temperature regimes. *Landscape and Ecological Engineering* 5: 115-123.
- Qu, LY., Makoto, K., Choi, DS., Quoreshi, AM., Koike, T. (2010) The Role of Ectomycorrhiza in Boreal Forest Ecosystem, In: Osawa, A. et al. (eds) *Permafrost Ecosystem - Siberian Larch Forest-*, Springer Ecological Studies 209. Pp. 413-425.
- Ryu K, Watanabe M, Shibata H, Takagi K, Nomura M, Koike, T. (2009) Ecophysiological responses of the larch species in northern Japan to environmental changes as a base of afforestation. *Landscape and Ecological Engineering* 5: 99 -106.
- Shinano, T., Yamamoto, T., Tawaraya, T., Tadokoro, M., Koike, T., Osaki, M. (2007) Effects of enriched CO₂ levels on the nutrient uptake characteristics of *Larix kaempferi*. *Tree Physiology*, 27: 97-104.
- Sitch, S., Cox, P.M., Collins, W.J., Huntingford, C. (2007) Indirect radiative forcing of climate change through ozone effects on the land-carbon sink. *Nature* 448: 791-794, doi:10.1038/nature/06059
- Sugai, T, Kam, D-G, Agathokleous, E, Watanabe, M, Kita K, Koike, T (2017) Growth and photosynthetic response of two larches exposed to O₃ mixing ratios ranging from pre-industrial to near future. *Photosynthetica* 56: 901-910.
- Sugai, T, Watanabe, T, Kita, K, Koike, T. (2019) Nitrogen loading increases the ozone sensitivity of larch seedlings with higher sensitivity to nitrogen loading. *Science of the Total Environ.*, 663: 587-595
- Szmidt, A.E., Aldén, T., Hällgren, J.E. (1987) Paternal inheritance of chloroplast DNA in *Larix*. *Plant Molecular Biology* 9: 59–64.
- Watanabe M, Yamaguchi M, Iwasaki M, Matsuo N, Naba J, Tabe C, Matsumura H, Kohno Y, Izuta T (2006) Effects of ozone and/or nitrogen load on the growth of *Larix kaempferi*, *Pinus densiflora* and *Cryptomeria japonica* seedlings. *Journal of Japanese Society Atmospheric Environment* 41:320-334
- Watanabe, M., Matsuo, N., Yamaguchi, M., Matsumura, H., Kohno, Y., Izuta, T. (2010) Risk assessment of ozone impact on the carbon absorption of Japanese

- representative conifers. *European Journal of Forest Research* 129: 421-430.
- Watanabe, M., Mao, Q., Novriyanti, E., Kita, K., Takagi, K., Satoh, F., Koike, T. (2013) Elevated CO₂ enhances the growth of hybrid larch F₁ (*Larix gmelinii* var. *japonica* × *L. kaempferi*) seedlings and changes its biomass allocation. *Trees* 27: 1647-1655
- Watanabe, Y., Wakabayashi, K., Kitaoka, S., Satomura, T., Eguchi, N., Watanabe, M., Nakaba, S., Takagi, K., Sano, Y., Funada, R., Koike, T. (2016) Response of tree growth and wood structure of *Larix kaempferi*, *Kalopanax septemlobus* and *Betula platyphylla* saplings to elevated CO₂ concentration for 5 years exposure in a FACE system. *Trees* 30: 1569-1579
- Wang, XN, Qu, LY., Mao, Q., Watanabe, M., Hoshika, Y., Koyama, A., Kawaguchi, K., Tamai, Y. Koike, T. (2015) Ectomycorrhizal colonization and growth of the hybrid larch F₁ under elevated CO₂ and O₃. *Environmental Pollution* 197: 116-126.
- Wang, XN., Agathokleous, E., Qu, LY., Fujita, S., Watanabe, M., Tamai, Y., Mao, Q., Koyama, A., Koike, T. (2018) Effects of simulated nitrogen deposition on ectomycorrhizae community structure in hybrid larch and its parents grown in volcanic ash soil: The role of phosphorous. *Science of the Total Environment* 618: 905-915.
- Zhang, P., Shao, G., Zhao, G., Le Master, D.C., Parker, G.P., Dunning, Jr. J.B., Li, Q. (2000) China's Forest Policy for the 21st Century. *Science* 288: 2135-2136.