HOKKAIDO UNIVERSITY

Title	Biological notes on Zeuxo ezoensis Okamoto et al., 2020(Crustacea : Peracarida : Tanaidacea)
Author(s)	Hayakawa1, Yoshinobu; Kakui, Keiichi
Citation	Aquatic Animals, AA2022, AA2022-10 https://doi.org/10.34394/aquaticanimals.2022.0_AA2022-10
Issue Date	2022-06
Doc URL	https://hdl.handle.net/2115/85805
Rights	Copyright(c)2022 Yoshinobu Hayakawa1 & Kakui Keiichi
Type	journal article
File Information	Azutic Animals AA2022-10.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Aquatic Animals

水生動物 第 2022 巻
令和 4 年 6 月

Notes and News

Biological notes on *Zeuxo ezoensis* Okamoto et al., 2020 (Crustacea: Peracarida: Tanaidacea)

Yoshinobu Hayakawa[1†], Keiichi Kakui[2*]

[1]School of Science, Hokkaido University, Sapporo 060-0810, Japan. [2]Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan. [†]Deceased.

[*]Corresponding author; e-mail: kakui@eis.hokudai.ac.jp, Tel: +81-11-706-2750.

Abstract

We report microhabitats used by the tanaidid tanaidacean *Zeuxo ezoensis* Okamoto et al., 2020 on the marine alga *Sargassum confusum* C. Agardh, as well as its molting behavior, based on observation of captive individuals. *Zeuxo ezoensis* constructed its tube-shaped nests on *S. confusum* (1) at the bases of lateral branches, (2) on main branches in the apical region, (3) in cracked air bladders, and (4) in clumps of a filamentous alga (Chordariaceae sp.) epiphytic on *S. confusum*. The behavioral sequence of a single molt observed for a male *Z. ezoensis* was as follows. (1) A break occurred along the ventrolateral and posterodorsal edges of the carapace, creating a dorsal opening in the old exoskeleton; (2) the animal began to extricate itself from the old exoskeleton through this opening; (3) the cephalothorax and its appendages shed the old exoskeleton; and (4) the animal completed extrication from the old exoskeleton. The entire molt took about five minutes.

Key words: benthos, epibiont, Tanaidoidea, Tanaidomorpha, tube dweller

Our current knowledge of the biological aspects of tanaidacean crustaceans is fragmentary. This is true even for the family Tanaididae, representatives of which are commonly found in shallow waters, and to which perhaps most relevant attention has been given. Tanaididae contains about 90 shallow-water species (cf. Anderson 2020), but biological information is available for only a few of them and is limited to a few genera (e.g., Johnson and Attramadal 1982a, b; Matsumasa 1994; Nakaoka 2002; Morales-Vela et al. 2008; Chim et al. 2016; Tanabe et al. 2017).

Zeuxo ezoensis Okamoto et al., 2020 is a shallow-water tanaidid reported from Hokkaido, Japan (Okamoto et al. 2020), from among algae such as the brown *Sargassum* spp. and red *Neorhodomela aculeata* (Perestenko) Masuda (Kakui et al. 2017). Using silk, it constructs tube-shaped nests (Kakui et al. 2021), that may function to provide shelter from predators such as gobiid fish (Kakui 2015), staurozoan jellyfish (Kakui 2022), and anthuroid isopods (Shiraki and Kakui 2022). In this study, based on YH's observations of captive animals in 1980, we report (1) microhabitats on *Sargassum confusum* C. Agardh used by *Z. ezoensis* and (2) the behavioral sequence of its molt.

Microhabitats on *Sargassum confusum*

Several dishes (50 or 80 mm in diameter; kept at about 15°C under ambient light), each containing one female and one male *Z. ezoensis*, a branch of *S. confusum*, and seawater, were prepared and observed more than once a day. Tanaidids constructed their tubes (1) on the bases of lateral branches, (2) on main branches in the apical region,

(3) in cracked air bladders, and (4) among a filamentous brown alga Elachistaceae sp. [= Chordariaceae sp. in the recent classification] epiphytic on *S. confusum* (Fig. 1). We note that in the wild, *S. confusum* thalli with many clumps of epiphytic algae attached tended to harbor more *Z. ezoensis* than those with only a few.

Molting behavior

Molting typically occurred within nest, making observation impossible. Only one molt by a male outside a nest was observed; it involved the following behavioral sequence (Fig. 2). (1) A dorsal transverse break occurred in the old exoskeleton between the carapace and pereonite 1. This break extended anteriorly to the base of the eye lobe and along the ventrolateral edges of the carapace, creating a dorsal opening in the old exoskeleton. While this was occurring, the animal occasionally shook the posterior body dorsally and ventrally, actively moved its pereopods, and frequently touched its antennules with its chelipeds (Fig. 2A). (2) The animal began to extricate itself from the old exoskeleton through the dorsal opening, as its posterior body moved peristaltically (Fig. 2B). (3) The cephalothorax and its appendages soon shed the old exoskeleton, with the posterior part of the body remaining in the old exoskeleton. The animal continuously shook the posterior end of the body in the dorso-ventral plane



Fig. 1. Microhabitats on *Sargassum confusum* used by *Zeuxo ezoensis* Okamoto et al., 2020. A–D, tube-shaped nests (shown in blue) constructed by *Z. ezoensis* at the base of a lateral branch (A), on a main branch in the apical region (B), in a cracked air bladder (C), and among algae epiphytic on *S. confusum* (D). Drawing by Yoshinobu Hayakawa.



Fig. 2. Behavioral sequence of molting in *Zeuxo ezoensis* Okamoto et al., 2020. A, a dorsal opening (arrowhead) appears in the old exoskeleton. B, the animal begins extrication from the old exoskeleton. C, the cephalothorax and its appendages shed the old exoskeleton. D, the animal completes extrication from the old exoskeleton. Arrows indicate movements of parts of the body. Details of the sequence are provided in the main text. Drawing by Yoshinobu Hayakawa.

(Fig. 2C). (4) The animal completed extrication from the old exoskeleton, continuously shaking the posterior part of the body dorsoventrally, possibly to help extend the new exoskeleton (Fig. 2D). The molt was completed in about five minutes. To our knowledge, this is the second study describing the behavioral sequence of molting in Tanaidacea. The sequence reported here resembles another example observed by Bückle Ramírez (1965) in the leptocheliid *Heterotanais oerstedii* (Krøyer, 1842).

The natural history of tanaidaceans remains largely unknown. Recent observations of living tanaidaceans under both wild and captive conditions have brought new discoveries, such as a tanaidid species dwelling in barnacle tests (Chim et al. 2016; see also Chim and Tong 2019), mucoid tube construction in a parapseudid species (Kakui and Hiruta 2017), and the second example of sex change in a nototanaid species (Kakui and Hiruta 2022). Future observations will undoubtedly make further discoveries.

Acknowledgments

This study was carried out in 1980 by the late YH under the guidance of the late Tatsunori Itô, with help in sampling from the late Kazuro Shinta. KK thanks Reiko Hayakawa, YH's wife, for permission to use of YH's drawings, and Matthew H. Dick for reviewing the manuscript and editing the English.

References

Anderson, G. (2020). Tanaidacea—Forty Years of Scholarship, Version 3.0. https://aquila.usm.edu/tanaids30/5/ (accessed 2 May 2022).

Bückle Ramírez, L. F. (1965). Untersuchungen über die Biologie von *Heterotanais oerstedi* Kröyer (Crustacea, Tanaidacea). Z. Morphol. Ökol. Tiere 55: 714–782.

Chim, C. K., Wong, H. P.-S., Tan, K. S. (2016). *Tetraclita* (Cirripedia, Thoracica) tests as an important habitat for intertidal isopods and other marine and semi-terrestrial fauna on tropical rocky shores. Crustaceana 89: 985–1040.

Chim, C. K., Tong, S. J. W. (2019). *Xenosinelobus balanocolus*, a new tanaidid genus and species (Crustacea: Peracarida: Tanaidacea) from barnacles on intertidal rocky shores and seawalls in the Singapore Strait. Zootaxa 4629: 413–427.

Johnson, S. B., Attramadal, Y. G. (1982a). A functional-morphological model of *Tanais cavolinii* Milne-Edwards (Crustacea, Tanaidacea) adapted to a tubicolous life-strategy. Sarsia 67: 29–42.

Johnson, S. B., Attramadal, Y. G. (1982b). Reproductive behaviour and larval development of *Tanais cavolinii* (Crustacea: Tanaidacea). Mar. Biol. 71: 11–16.

Kakui, K. (2015). First report of *Zeuxo* sp. (Crustacea: Tanaidacea) as prey for the forktongue goby, *Chaenogobius annularis* Gill, 1859. Rishiri Stud 34: 1–6. (in Japanese with English abstract).

Kakui, K., Hiruta, C. (2017). Tube construction by a tanaidacean crustacean using a novel mucus secretion system involving the anal opening. Zool. Lett. 3: 20.

Kakui, K., Hayakawa, Y., Katakura, H. (2017). Difference in size at maturity in annual and overwintering generations in the tanaidacean *Zeuxo* sp. in Oshoro Bay, Hokkaido, Japan. Zool. Sci. 34: 129–136.

Kakui, K., Fleming, J. F., Mori, M., Fujiwara, Y., Arakawa, K. (2021). Comprehensive transcriptome sequencing of Tanaidacea with proteomic evidences for their silk. Genome Biol. Evol. 13: evab281.

Kakui, K. (2022). Predation of tanaidacean crustaceans by *Haliclystus tenuis* (Cnidaria: Staurozoa) in captivity. Bull. Otaru Mus. 35: 9–12. (in Japanese with English abstract).

Kakui, K., Hiruta, C. (2022). Protogynous hermaphroditism in Crustacea: a new example from Tanaidacea. Can. J. Zool. doi.org/10.1139/cjz-2022-0033.

Matsumasa, M. (1994). Effect of secondary substrate on associated small crustaceans in a brackish lagoon. J. Exp. Mar. Biol. Ecol. 176: 245–256.

Morales-Vela, V., Suárez-Morales, E., Padilla-Saldívar, J., Heard, R. W. (2008). The tanaid *Hexapleomera robusta* (Crustacea: Peracarida) from the Caribbean manatee, with comments on other crustacean epibionts. J. Mar. Biol. Assoc. U.K. 88: 591–596.

Nakaoka, M. (2002). Predation on seeds of seagrasses *Zostera marina* and *Zostera*

caulescens by a tanaid crustacean *Zeuxo* sp. Aquat. Bot. 72: 99–106.
Okamoto, N., Oya, Y., Kakui, K. (2020). A new species of *Zeuxo* (Crustacea: Peracarida: Tanaidacea) from Japan, with remarks on carapace pigmentation as a potentially useful taxonomic character. Mar. Biol. Res. 16: 411–422.
Shiraki, S., Kakui, K. (2022). Observations on predation in *Paranthura japonica* Richardson, 1909 (Isopoda: Cymothoida: Paranthuridae). Zool. Sci. 39: 270–274.
Tanabe, Y., Hayashi, R., Tomioka, S., Kakui, K. (2017). *Hexapleomera urashima* sp. nov. (Crustacea: Tanaidacea), a tanaidid epibiotic on loggerhead sea turtles at Yakushima Island, Japan. Zootaxa 4353: 146–160.

Received: 13 May 2022 | Accepted: 6 June 2022 | Published: 9 June 2022