



HOKKAIDO UNIVERSITY

Title	Regional differences in the fatty acid composition, and vitamin and carotenoid concentrations in farm bulk milk in Hokkaido, Japan
Author(s)	Mitani, Tomohiro; Kobayashi, Kuniyuki; Ueda, Koichiro et al.
Citation	Animal science journal, 92(1), e13570 https://doi.org/10.1111/asj.13570
Issue Date	2021-06-20
Doc URL	https://hdl.handle.net/2115/86118
Rights	This is the peer reviewed version of the following article: Mitani, T, Kobayashi, K, Ueda, K, Kondo, S. (2021). Regional differences in the fatty acid composition, and vitamin and carotenoid concentrations in farm bulk milk in Hokkaido, Japan. Anim Sci J, 92, e13570., which has been published in final form at https://doi.org/10.1111/asj.13570 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Type	journal article
File Information	asj.13570.pdf



Title:

Regional differences in the fatty acid composition, and vitamin and carotenoid concentrations in farm bulk milk in Hokkaido, Japan

Category: Research Articles

Authors:

Tomohiro Mitani¹, Kuniyuki Kobayashi², Koichiro Ueda³, and Seiji Kondo³

Institutions:

¹ Field Science Center for Northern Biosphere, Hokkaido University, Sapporo 060-0811, Japan

² Graduate School of Global Food Resources, Hokkaido University, Sapporo, 060-8589, Japan

³ Research Faculty of Agriculture, Hokkaido University, Sapporo, 060-8589, Japan

Corresponding Author:

Tomohiro MITANI

Postal address: Field Science Center for Northern Biosphere, Hokkaido University, Kita 11, Nishi 10, Kita-ku, Sapporo 060-0811, Japan

E-mail: tmitani@fsc.hokudai.ac.jp

ABSTRACT

In this study, we investigated the regional differences in the composition of farm bulk milk produced at three different dairy areas in Hokkaido, Japan. A field survey was conducted at Central, Tokachi, and North areas of Hokkaido, three or four times a year. At each farm, an interview questionnaire for farm basal data was conducted, and 500 mL of bulk tank milk sample was obtained. Fatty acid composition, and vitamin and carotenoid concentrations in the milk samples were determined. In Central and Tokachi areas, corn silage was used as the main forage. In North area, fresh herbage was the dominant feed in the summer season, and grass hay was the main feed in the winter season. Discriminant analysis revealed that the composition of milk samples differed among the areas and seasons. Milk from Central and Tokachi areas contained a higher ratio of linoleic acid compared with that from North area, but there were only slight differences in the composition of milk between Central and Tokachi areas. The concentrations of carotenoids and α -tocopherol were higher in samples from North area and the ratios of *trans*-vaccenic acid and conjugated linoleic acid were higher in the summer season than in the indoor season.

Key words: feeding management, milk compositions, regional difference, Hokkaido

1 INTRODUCTION

In the Japanese dairy industry, Hokkaido is the most important region accounting for more than half of the milk produced in Japan (Ministry of Agriculture, Forestry, & Fisheries, 2020). Dairy farms in Hokkaido can be divided into three typical feeding management types (Harada, Gonoï & Takahashi, 1983). These types correspond to geographical regions differing in climate, soil, and social environment for dairy farmers (Figure 1). Tokachi and Okhotsk areas are typical upland-cropping dairy areas. Dairy farmers in these areas can use corn silage (CS) as self-making forage owing to the temperate climate. North and Kosen areas are typical grassland dairy areas because the applicable forage in these areas is limited to grass forage owing to low cumulative temperature and short sunshine duration. Central and South areas are typical sub-urban dairy areas where farms for dairy production are small owing to the high price of farmland although CS can be used as forage in these areas.

Feeding management for dairy cows strongly affects the composition of milk. It not only affects the major components of milk, such as milk fat, protein, and lactose, but also the minor components, such as fatty acids (FAs), vitamins, and carotenoids (e.g., Bauman & Griinari, 2003; Chilliard et al., 2007; Palmquist, Beaulieu & Barbano, 1993; Walker, Dunshea & Doyle, 2004). These components strongly influence the physical property and visual aspects of dairy products (Couvreur et al., 2006; Larsen et al., 2014). In addition, feeding management for dairy cows influences the sensory properties of milk and dairy products (Croissant et al., 2007; Martin et al., 2005). Therefore, milk composition could differ among the different regions of Hokkaido, reflecting the differences in the feeding management.

Recently, the interests of consumers in dairy products, especially in Western countries, have changed, not only with regard to taste and functional ingredients

beneficial for human health, but also in respect of how and where the milk used in the products is produced, for examples, issues related to the organic nature of product, animal welfare, sustainable development goals (SDGs), and protected denominations of origin (PDO) (Capuano et al., 2013; Elgersma, Tamminga & Ellen, 2006; Trubek & Bowen, 2008). To cater to these demands, the characteristics of milk produced in each region of Hokkaido have been investigated. Wide range on-farm surveys covering standard farmers are essential to determine the differences in the composition of milk resulting from regional differences. However, such on-farm surveys targeting regional differences have been scarce worldwide (Gaspardo et al., 2010; Larsen et al., 2010), as well as in Japan (Yayota et al., 2013), and none has been conducted in Hokkaido. In Europe, a study was conducted to clarify the relationship between the FA composition of milk and farm feeding systems using a dataset of 1,248 farm bulk milk samples (Coppa et al. 2013 and 2015).

In this study, we conducted an on-farm survey in three typical dairy areas of Hokkaido, namely Tokachi, North, and Central, to determine regional differences in the composition of farm bulk milk, with respect to the FA composition, and concentrations of vitamins and carotenoids which strongly influenced on the physical property and visual aspects of dairy products. Based on these results, characteristics of milk and different production systems can be described, and the diversity of Hokkaido milk products can be indicated.

2 MATERIALS AND METHODS

2.1 Field survey

The field survey was conducted at three regions in Hokkaido (Tokachi area located around 42°92'N and 143°20'E, North area located around 45°12'N and 142°36'E, and Central

area located around 43°06'N and 141°34'E,) during the winter (December) season in 2008, and during the spring (April or May) and summer (August or September) season in 2009. For North area, an additional survey was conducted in the early summer (June) season in 2009. In each season, 47 dairy farmers at 25, 13, and 9 farms in Tokachi, North, and Central areas, respectively, were surveyed. The farmers in Tokachi and Central areas were selected by an introduction of the agricultural cooperative in each area. The all of farmers in North adopted grazing management in the summer season and were selected from the farmers surveyed in the previous study (Mitani et al., 2016). The survey method was similar to one described in our previous study, in which an interview survey was conducted and 500 mL of bulk tank milk was collected (Mitani et al., 2016). The data of milk yield, and the amount and type of supply feeds were obtained from the interview survey. The collected milk samples were brought to a laboratory in cold storage and divided into sub-samples. The sub-sample for analysis of milk composition was immediately sent to the Laboratory of Hokkaido Dairy Milk Recording and Testing Association. The concentrations of milk fat, milk protein, lactose, and solids not fat (SNF) were determined using a Fourier transform infrared device (MilkoScan FT+; Foss Electric, Hillerød, Denmark). The sub-samples used for other analysis were stored at - 80 °C until use.

2.2 Chemical analysis of milk

The milk sample used for FA analysis was thoroughly thawed under tap water, and then warmed in a water bath to dissolve the fat. FAs were extracted using a modified version of the Roese–Gottlieb method (ISO and IDF, 2001), and were methylated using a 2 N solution of NaOCH₃ in methanol and a 14% boron trifluoride solution in methanol (Christie, 2001). The FA methyl esters, thus formed, were analyzed using a gas

1 chromatograph equipped with a flame ionization detector (GC-2010; Shimadzu Co.,
2 Kyoto, Japan). The FA methyl esters were detected under the conditions described in our
3 previous study (Mitani et al. 2016).

4 The concentrations of carotenoids and retinol were analyzed using a modified
5 version of a previously described method (Hulshof et al., 2006). The milk fat was
6 extracted using a modified version of the Roese–Gottlieb method (ISO and IDF, 2001),
7 and was saponified using a 5% KOH solution in ethanol. Echinenone was used as an
8 internal standard. The concentrations of carotenoids and retinol in extracted samples were
9 analyzed by high performance liquid chromatography (LaCrome Elite; Hitachi High-
10 Tech Science Corporation, Tokyo, Japan). Carotenoids and retinol were separated on a
11 C18-ODS column (Cadenza CD-C18, 150 × 2 mm; Imtakt, Kyoto, Japan) with a gradient
12 elution method, and detected using a diode array detector (retinol at 325 nm, and lutein
13 and β -carotene at 450 nm).

14 The concentration of α -tocopherol was determined with a direct saponification
15 method. The milk sample was saponified using 2N KOH solution in methanol, containing
16 ascorbic acid, and α -tocopherol was extracted with hexane. α -tocotrienol was used as an
17 internal standard. The extracted sample was analyzed by high performance liquid
18 chromatography (LaCrome Elite; Hitachi High-Tech Science Corporation). α -Tocopherol
19 was separated on a C18-ODS column (Cadenza CD-C18, 150 × 2 mm; Imtakt), and
20 detected using a diode array detector at 292 nm.

22 **2.3 Statistical analysis**

23 All the farmers in North area conducted grazing management during the summer season.
24 In this area, winter and spring seasons, and early summer and summer seasons were
25 regarded as indoor and grazing seasons, respectively, and results are therefore described

1 for these two seasons. Most of the farmers in Central and Tokachi areas did not conduct
2 grazing management. Therefore, results from these areas were not categorized by seasons.

3 Statistical analysis was conducted using the JMP Pro 14.3.0 (SAS Institute Inc.,
4 Cary, NC, USA) and SIMCA 13.0.3 (Umetrics AB., Umeå, Sweden) software. Data were
5 analyzed with the one-way analysis of variance model using the Fit Model Platform in
6 JMP. The regional differences in milk composition were analyzed using the orthogonal
7 partial least squares discrimination analysis model in SIMCA. The analyses were
8 conducted using standardized data and the full cross-validation method (leave one out).
9 Predictive and orthogonal parameters in the model were estimated by the auto-fitting
10 procedure in SIMCA.

12 **3 RESULTS**

13 **3.1 Farm basal data and feeding management**

14 The mean values and the range of farm basal data and feeding management are shown in
15 Table 1. The farm characteristics were distinct for each area. In Tokachi area, the herd
16 size was big and milk yield per cow and per lactation was high. In Central area, the herd
17 size was not large but milk yield per cow and per lactation was high as in the farms of
18 Tokachi area. In North area, the herd size was small and milk yield per cow and per
19 lactation was not high. Milk yield per cow and per lactation in Central and Tokachi areas
20 was 1.2-times higher than that in North area.

21 The supply of primary and secondary forage also differed among the regions
22 (Table 2). In Central and Tokachi areas, CS was used as the main forage, but the supply
23 of fresh herbage was less. Although farmers conducted similar feeding management in
24 Central and Tokachi areas, those in Central area supplied relatively more hay, and those
25 in Tokachi area supplied relatively more grass silage (GS). Forage in North area was

1 limited to grass forage, such as a GS, hay, and fresh herbage. Fresh herbage was used as
2 the main forage during the grazing season in North area. The supply of concentrate in
3 Central and Tokachi areas was higher than that in North area, which mostly ranged
4 between 5 and 8 kg/day. Half of the farmers in North area supplied less than 5 kg/day of
5 concentrate.

6 7 **3.2 Regional differences in milk composition**

8 The milk composition differed significantly among regions and seasons, except for the
9 concentration of milk protein, the ratio of C10:0 FA, and retinol concentration (Table 3).
10 For visualization of the regional differences in milk composition, a discriminant analysis
11 was conducted using 23 parameters, which included 4 of major milk composition, 15 of
12 FAs, and 4 of vitamins and carotenoids (Figure 2). The discriminant model included three
13 predictive parameters (P1, P2, and P3) and seven orthogonal parameters by auto-fitting.
14 The cumulative R^2 value was 0.578 (0.269, 0.197, and 0.112 for P1, P2, and P3,
15 respectively), and the cumulative Q^2 value was 0.488 (0.259, 0.171, 0.058 for P1, P2, and
16 P3, respectively). The results showed that the model significantly indicated the regional
17 differences in milk composition. The model also correctly distinguished each area using
18 milk composition. The percentage of correct discrimination was high, being 96% in
19 Tokachi (72/75), 88.5% in North-indoor (23/26), and 88.5% in North-grazing (23/26);
20 however, the percentage in Central area was low, being 55.6% (15/27).

21 Predictive parameter 1 separated Tokachi and Central areas (positive), and North
22 area (negative). Milk composition clearly differed between Tokachi and Central area, and
23 North area. Predictive parameter 1 correlated positively with the ratio of *cis*-9,12 C18:2
24 (linoleic acid: LA) and lactose concentration, and negatively with β -carotene, lutein, and
25 α -tocopherol concentrations, and ratios of *trans*-11 C18:1 (*trans*-vaccenic acid: TVA),

cis-9, *trans*-11 C18:2 (conjugated linoleic acid: CLA), and *cis*-9,12,15 C18:3 (α -linolenic acid: ALA).

Predictive parameter 2 separated the indoor (positive) and grazing seasons (negative) in North area. Milk composition clearly differed between the indoor and grazing seasons in North area. Predictive parameter 2 correlated positively with milk fat concentration, the ratios of C16:0, C16:1, C17:0, and C20:0, and negatively with the ratio of C18:0, *cis*-9 C18:1 (Oleic acid: OA), TVA, LA, and CLA.

Predictive parameter 3 separated Central (positive) and Tokachi (negative) areas. However, the separation was insufficient, and half of the milk samples in Central area were included in a similar range of milk from Tokachi area. Predictive parameter 3 correlated positively with the ratio of C12:0 and C14:0, and negatively with the ratio of C16:1, ALA, C20:0, and retinol concentration.

4 DISCUSSIONS

In the present study, we analyzed, for the first time, the regional differences in milk composition in Hokkaido. Hokkaido is very interesting island with regard to the rich diversity in the feeding management of cows. The rich diversity in each region was originated from the difference in the available forage. The differences in the supplied forage among the regions of Hokkaido have not changed much from the 1980s (Harada et al., 1983). Therefore, the differences in feeding management among the regions in Hokkaido essentially resulted from the differences in regional characteristics, such as climate and social environment.

There have been few studies investigating the regional differences in milk composition, with regard to FA composition, and vitamin and carotenoid concentrations (Gaspardo et al., 2010; Larsen et al., 2010; Yayota et al., 2013). The results of the

discriminant analysis (Figure 2) indicate that the milk derived from farms with specific feeding management in each area could be distinguished using the milk composition. This also indicates that the distinct milk composition in each area represents characteristics for selected farms located in each area. Gaspardo et al. (2010) showed that the origin of bulk milk from Italy or Slovenia could be discriminated using milk the FA composition. Larsen et al. (2010) reported that the FA composition, and vitamin and carotenoid concentrations in farm bulk milk differed among geographical areas and seasons in Sweden.

The milk FA composition in Tokachi and Central areas showed a higher ratio of LA (positive P1 in Figure 2). This was because farmers in these areas supplied more CS and grain feeds than those in North area. A high LA intake in Tokachi and Central areas was easily expected because grain feeds, including CS, contain higher amount of LA compared with grass forage (Walker et al., 2004). Not only the high LA intake, the high grain intake, including CS, promotes an uptake of LA from the duodenum. In general, high grain intake tends to depress the rumination function and reduces the ruminal pH. Under these circumstances, the function of biohydrogenation by rumen microbes reduces and the amount of LA escaping from the rumen increases (Chilliard et al., 2007; Dewhurst et al., 2006).

The milk composition differed slightly between Tokachi and Central areas because of similar feeding management in these areas. The slightly higher ratios of C12:0 and C14:0 in Central area compared with those in Tokachi area would have resulted from the difference in secondary forage such as hay and low moisture GS in Central area and GS in Tokachi area. This would probably be caused by a high intake of rough forage, such as hay and low moisture GS, which induces not only the rumination behavior in cows but also increases the production of acetate in the rumen, a substrate in the *de novo* synthesis of FAs in the mammary glands (Chilliard et al., 2000).

1 The samples from North area were high in TVA, CLA, and ALA (negative P1 in
2 Figure 2). This was because of the high intake of grass forage in this area. The main FA
3 in grass forage is ALA and that in grain feeds is LA (Walker et al., 2004). The ingested
4 ALA is converted to TVA and CLA upon biohydrogenation and isomerization by rumen
5 microbes, and these FAs are then absorbed from the duodenum and excreted into the milk
6 after transformation (Chilliard et al., 2007). Therefore, a high ratio of these FAs in milk
7 samples from North areas was a natural consequence of feeding.

8 Because FAs, including ALA in GS or hay, decreased during ensiling or drying
9 (Dewhurst et al., 2006; Kalač & Samková, 2010; Villeneuve et al., 2013), the ratio of
10 TVA, CLA, and ALA in milk was higher in the grazing season than in the indoor season.
11 The milk FA composition between grazing and indoor feeding has been compared in
12 several studies (e.g., Croissant et al., 2007; Kay et al., 2005; Kelly et al., 1998; Schroeder
13 et al., 2003). In a previous study on milk FA composition of dairy farmers in grassland
14 dairy areas of Hokkaido (Mitani et al., 2016), a high ratio of CLA, TVA, and OA during
15 the grazing season and a high ratio of C16:0 during the indoor season was confirmed. Our
16 results are in agreement with those of the previous study.

17 The content of fat-soluble antioxidants, such as carotenoids and α -tocopherol, is
18 higher in grasses than in grains, such as corn (Noziere et al., 2006). The content of
19 carotenoids and α -tocopherol in grass forage decreases in the order fresh herbage > GS >
20 hay because these compounds were easily broken down upon exposure to sunlight
21 (Agabriel et al., 2007; Noziere et al., 2006). Therefore, the high carotenoid and α -
22 tocopherol content in samples from North area, especially during the grazing season, was
23 a natural consequence of higher intake of grass forage and grazing in the selected farms
24 in North areas than in Tokachi and Central areas.

25 To summarize, milk samples from the upland cropping and sub-urban dairy areas

1 had a white aspect with low carotenoid content, and contained melty fat with a high ratio
2 of LA. Milk produced during the grazing season in farms adopting grazing management
3 in grassland dairy areas had a strong orange aspect with high carotenoid content, and
4 contained melty fat with a high ratio of CLA and ALA, and that produced during the
5 indoor season in these areas had an orange aspect, and contained firm fat with a high ratio
6 of saturated FAs.

7 In conclusion, the characteristics of milk in each area originated strongly from
8 farm feeding management, especially of the main supply forage, which was defined as
9 climate and social environment for dairy farmers in each area. The present study is the
10 first to unravel the characteristics of farm bulk milk produced in each of the dairy areas
11 of Hokkaido. However, the results presented here also indicate that milk composition
12 varies widely within each area. Therefore, factors, such as feeding management in each
13 farm, influencing the distribution of milk composition within each area need to be
14 investigated in a future study.

16 **Acknowledgement**

17 We express gratitude to the dairy farmers who participated in this research. We would like
18 to thank Editage (www.editage.com) for English language editing.

20 **Conflict of interest**

21 Authors declare no conflict of interests for this article.

23 **References**

24 Agabriel, C., Cornu, A., Journal, C., Sibra, C., Grolier, P. & Martin, B. (2007). Tanker
25 milk variability according to farm feeding practices: vitamins A and E, carotenoids,

1 color, and terpenoids. *Journal of Dairy Science*, 90 (10), 4884-4896.

2 <https://doi.org/10.3168/jds.2007-0171>

3 Bauman, D. E. & Griinari, J. M. (2003). Nutritional regulation of milk fat synthesis.

4 *Annual Review of Nutrition*, 23, 203-227.

5 <https://doi.org/10.1146/annurev.nutr.23.011702.073408>

6 Capuano, E., Boerrigter-Eenling, R., van der Veer, G. & van Ruth, S. M. (2013).

7 Analytical authentication of organic products: an overview of markers. *Journal of*

8 *the Science of Food and Agriculture*, 93 (1), 12-28.

9 <https://doi.org/https://doi.org/10.1002/jsfa.5914>

10 Chilliard, Y., Ferlay, A., Mansbridge, R. M. & Doreau, M. (2000). Ruminant milk fat

11 plasticity: nutritional control of saturated, polyunsaturated, trans and conjugated

12 fatty acids. *Annales de Zootechnie* (pp. 181-206). Paris: Institut national de la

13 recherche agronomique, 1960-2000.

14 Chilliard, Y., Glasser, F., Ferlay, A., Bernard, L., Rouel, J. & Doreau, M. (2007). Diet,

15 rumen biohydrogenation and nutritional quality of cow and goat milk fat.

16 *European Journal of Lipid Science and Technology*, 109 (8), 828-855.

17 <https://doi.org/10.1002/ejlt.200700080>

18 Christie, W. W. (2001). A practical guide to the analysis of conjugated linoleic acid (CLA).

19 *Inform*, 12, 147-152.

20 Coppa, M., Ferlay, A., Chassaing, C., Agabriel, C., Glasser, F., Chilliard, Y., ... Martin, B.

21 (2013). Prediction of bulk milk fatty acid composition based on farming practices

22 collected through on-farm surveys. *Journal of Dairy Science*, 96 (7), 4197-4211.

23 <https://doi.org/10.3168/jds.2012-6379>

24 Coppa, M., Chassaing, C., Ferlay, A., Agabriel, C., Laurent, C., Borreani, G., ... Martin,

25 B. (2015). Potential of milk fatty acid composition to predict diet composition and

1 authenticate feeding systems and altitude origin of European bulk milk. *Journal*
2 *of Dairy Science*, 98 (3), 1539-1551. <https://doi.org/10.3168/jds.2014-8794>

3 Couvreur, S., Hurtaud, C., Lopez, C., Delaby, L. & Peyraud, J. L. (2006). The linear
4 relationship between the proportion of fresh grass in the cow diet, milk fatty acid
5 composition, and butter properties. *Journal of Dairy Science*, 89 (6), 1956-1969.
6 [https://doi.org/https://doi.org/10.3168/jds.S0022-0302\(06\)72263-9](https://doi.org/10.3168/jds.S0022-0302(06)72263-9)

7 Croissant, A. E., Washburn, S. P., Dean, L. L. & Drake, M. A. (2007). Chemical properties
8 and consumer perception of fluid milk from conventional and pasture-based
9 production systems. *Journal of Dairy Science*, 90 (11), 4942-4953.
10 <https://doi.org/10.3168/jds.2007-0456>

11 Dewhurst, R. J., Shingfield, K. J., Lee, M. R. F. & Scollan, N. D. (2006). Increasing the
12 concentrations of beneficial polyunsaturated fatty acids in milk produced by dairy
13 cows in high-forage systems. *Animal Feed Science and Technology*, 131 (3-4),
14 168-206. <https://doi.org/10.1016/j.anifeedsci.2006.04.016>

15 Elgersma, A., Tamminga, S. & Ellen, G. (2006). Modifying milk composition through
16 forage. *Animal Feed Science and Technology*, 131 (3-4), 207-225.
17 <https://doi.org/10.1016/j.anifeedsci.2006.06.012>

18 Gaspardo, B., Lavrencic, A., Levart, A., Del Zotto, S. & Stefanon, B. (2010). Use of milk
19 fatty acids composition to discriminate area of origin of bulk milk. *Journal of*
20 *Dairy Science*, 93 (8), 3417-3426. <https://doi.org/10.3168/jds.2009-2788>

21 Harada, T., Gono, Y. & Takahashi, M. (1983). Seasonal variations of the composition of
22 farm bulk milk in Hokkaido [In Japanese]. *Japanese Journal of Dairy and Food*
23 *Science*, 32 (5), 197-203.

24 Hulshof, P. J. M., van Roekel-Jansen, T., van de Bovenkamp, P. & West, C. E. (2006).
25 Variation in retinol and carotenoid content of milk and milk products in The

1 Netherlands. *Journal of Food Composition and Analysis*, 19 (1), 67-75.

2 <https://doi.org/10.1016/j.jfca.2005.04.005>

3 ISO (International Organization for Standardization), IDF (International Dairy
4 Federation) (2001). Milk and milk products - Extraction methods for lipids and
5 liposoluble compounds: ISO 14156: 2001, IDF 172:2001. Geneva and Brussels,
6 Switzerland: ISO and IDF.

7 Kalač, P. & Samková, E. (2010). The effects of feeding various forages on fatty acid
8 composition of bovine milk fat: A review. *Czech Journal of Animal Science*, 55
9 (12), 521-537. <https://doi.org/10.17221/2485-CJAS>

10 Kay, J. K., Roche, J. R., Kolver, E. S., Thomson, N. A. & Baumgard, L. H. (2005). A
11 comparison between feeding systems (pasture and TMR) and the effect of vitamin
12 E supplementation on plasma and milk fatty acid profiles in dairy cows. *Journal*
13 *of Dairy Research*, 72 (3), 322-332. <https://doi.org/10.1017/S0022029905000944>

14 Kelly, M. L., Kolver, E. S., Bauman, D. E., Van Amburgh, M. E. & Muller, L. D. (1998).
15 Effect of intake of pasture on concentrations of conjugated linoleic acid in milk
16 of lactating cows. *Journal of Dairy Science*, 81 (6), 1630-1636.
17 [https://doi.org/10.3168/jds.S0022-0302\(98\)75730-3](https://doi.org/10.3168/jds.S0022-0302(98)75730-3)

18 Larsen, M. K., Andersen, K. K., Kaufmann, N. & Wiking, L. (2014). Seasonal variation
19 in the composition and melting behavior of milk fat. *Journal of Dairy Science*, 97
20 (8), 4703-4712. <https://doi.org/10.3168/jds.2013-7858>

21 Larsen, M. K., Nielsen, J. H., Butler, G., Leifert, C., Slots, T., Kristiansen, G. H. &
22 Gustafsson, A. H. (2010). Milk quality as affected by feeding regimens in a
23 country with climatic variation. *Journal of Dairy Science*, 93 (7), 2863-2873.
24 <https://doi.org/10.3168/jds.2009-2953>

25 Martin, B., Verdier-Melz, I., Buchin, S., Hurtaud, C. & Coulon, J. B. (2005). Invited

1 paper: How do the nature of forages and pasture diversity influence the sensory
2 quality of dairy livestock products? *Animal Science* (81), 205-212.
3 <https://doi.org/https://doi.org/10.1079/ASC50800205>

4 Ministry of Agriculture, Forestry and Fisheries. (2020). The situation over the livestock
5 farming, statistics data, press release [homepage on the Internet]. Tokyo, Japan:
6 Ministry of Agriculture, Forestry and Fisheries; [cited 2020 2 April]. Available
7 from URL: https://www.maff.go.jp/j/chikusan/kikaku/lin/1_hosin/

8 Mitani, T., Kobayashi, K., Ueda, K. & Kondo, S. (2016). Discrimination of “grazing milk”
9 using milk fatty acid profile in the grassland dairy area in Hokkaido. *Animal*
10 *Science Journal*, 87 (2), 233-241. <https://doi.org/https://doi.org/10.1111/asj.12422>

11 Noziere, P., Graulet, B., Lucas, A., Martin, B., Grolier, P. & Doreau, M. (2006).
12 Carotenoids for ruminants: From forages to dairy products. *Animal Feed Science*
13 *and Technology*, 131 (3-4), 418-450.
14 <https://doi.org/10.1016/j.anifeedsci.2006.06.018>

15 Palmquist, D. L., Beaulieu, A. D. & Barbano, D. M. (1993). Feed and animal factors
16 influencing milk fat composition. *Journal of Dairy Science*, 76 (6), 1753-1771.
17 [https://doi.org/https://doi.org/10.3168/jds.S0022-0302\(93\)77508-6](https://doi.org/https://doi.org/10.3168/jds.S0022-0302(93)77508-6)

18 Schroeder, G. F., Delahoy, J. E., Vidaurreta, I., Bargo, F., Gagliostro, G. A. & Muller, L.
19 D. (2003). Milk fatty acid composition of cows fed a total mixed ration or pasture
20 plus concentrates replacing corn with fat. *Journal of Dairy Science*, 86 (10), 3237-
21 3248. [https://doi.org/10.3168/jds.S0022-0302\(03\)73927-7](https://doi.org/10.3168/jds.S0022-0302(03)73927-7)

22 Trubek, A. B. & Bowen, S. (2008). Creating the taste of place in the United States: can
23 we learn from the French? *Geojournal*, 73 (1), 23-30.
24 <https://doi.org/https://doi.org/10.1007/s10708-008-9175-3>

25 Villeneuve, M. P., Lebeuf, Y., Gervais, R., Tremblay, G. F., Vuillemand, J. C., Fortin, J. &

1 Chouinard, P. Y. (2013). Milk volatile organic compounds and fatty acid profile
2 in cows fed timothy as hay, pasture, or silage. *Journal of Dairy Science*, 96 (11),
3 7181-7194. <https://doi.org/10.3168/jds.2013-6785>

4 Walker, G. P., Dunshea, F. R. & Doyle, P. T. (2004). Effects of nutrition and management
5 on the production and composition of milk fat and protein: a review. *Australian*
6 *Journal of Agricultural Research*, 55 (10), 1009-1028.
7 <https://doi.org/10.1071/Ar03173>

8 Yayota, M., Tsukamoto, M., Yamada, Y. & Ohtani, S. (2013). Milk composition and flavor
9 under different feeding systems: a survey of dairy farms. *Journal of Dairy Science*,
10 96 (8), 5174-5183. <https://doi.org/10.3168/jds.2012-5963>

1 **Figure legends**

2 Figure 1. Map of the dairy areas and farming types in Hokkaido, Japan.

3

4 Figure 2. Results of the orthogonal partial least squares discrimination analysis based on

5 milk composition and area (-season) of Hokkaido, Japan. Upper panels indicate

6 predictive parameters 1 and 2 (P1 and P2). Lower panels indicate P1 and P3. Left and

7 right panels indicate the eigen values and eigenvectors, respectively.

8

1 和文抄録：

2 表題：

3 北海道における酪農家バルク乳中の脂肪酸組成、ビタミンおよびカロテノイド
4 含量の地域間の違い

6 著者名：

7 三谷朋弘¹・小林国之²・上田宏一郎³・近藤誠司³

9 所属機関：

10 ¹ 北海道大学 北方生物圏フィールド科学センター，札幌市北区，060-0811


11 ² 北海道大学 大学院国際食資源学院，札幌市北区，060-0809

12 ³ 北海道大学 大学院農学研究院，札幌市北区，060-8589

14 抄録本文：

15 本研究では、北海道における酪農家バルク乳成分の地域間の差異を調査した。
16 北海道の道央、十勝および道北地域において年3回もしくは4回、現地調査を
17 実施した。それぞれの農家で農家基礎情報についてのアンケート調査を実施
18 し、500 mLのバルクタンク乳サンプルを採取した。各乳サンプル中の脂肪酸組
19 成、ビタミンおよびカロテノイド含量を測定した。道央および十勝地域ではコ
20 ーンサイレージが主な粗飼料であり、道北地域では夏期に放牧草が、冬期には
21 乾牧草が主な粗飼料であった。乳成分23パラメータを用いた判別分析の結
22 果、各地域および季節を正確に判別可能であった。道央および十勝地域の乳は
23 これらの地域間の差異は小さかったが、道北地域と比較して乳中リノール酸割
24 合が高かった。道北地域の乳はカロテノイドおよび α -トコフェロール含量が高

- 1 く、舎飼い期の牛乳と比較して夏期の牛乳はトランス-バクセン酸および共役リ
- 2 ノール酸割合が高かった。






Figure 2

Table 1. Mean and range of management parameters for dairy farms in three areas of Hokkaido, Japan

	Area in Hokkaido		
	Central	Tokachi	North
	Mean $\pm$ SD	Mean $\pm$ SD	Mean $\pm$ SD
Number of farms	9	25	13
Herd size, lactating cows/farm	71 $\pm$ 52	105 $\pm$ 57	52 $\pm$ 17
Farming area, ha/farm	52 $\pm$ 31	45 $\pm$ 21	63 $\pm$ 15
Milk yield, kg/cow/day	26.5 $\pm$ 4.1	27.9 $\pm$ 3.1	23.0 $\pm$ 4.8
Milk production, kg/lactation	8067 $\pm$ 724	8570 $\pm$ 1271	6631 $\pm$ 2816

Table 2. Number of primary and secondary forage intake and concentrate supply level for farms in three areas (-seasons) of Hokkaido, Japan

	Area (-season)							
	Central		Tokachi		North-Indoor		North-Grazing	
Number	27		75		26		26	
Supply forage, number of farms	Primary	Secondary	Primary	Secondary	Primary	Secondary	Primary	Secondary
Grass silage (GS)	5	2	15	40	10	-	2	2
Corn silage	11	9	50	22	-	-	-	-
Hay + Low moisture GS	11	10	10	9	16	5	-	19
Pasture	-	-	-	-	-	-	24	2
Concentrate supply level, number of farms								
Low (< 5 kg/day)	4		11		14		12	
Midium (5 - 8 kg/day)	20		36		10		14	
High (> 8 kg/day)	3		28		2		-	

Table 3. Mean of milk compositions, milk fatty acid (FA) profile, and vitamin and carotenoid concentration for three areas (-season) of Hokkaido, Japan

	Area (-season)				Adusted SEM [†]	<i>P</i> value
	Central	Tokachi	North-Indoor	North-Grazing		
Milk compositions, %						
Milk fat	4.03	3.93	4.06	3.85	0.04	< 0.01
Milk protein	3.25	3.26	3.21	3.26	0.02	0.32
Lactose	4.50	4.51	4.42	4.44	0.01	< 0.01
Solid not fat	8.75	8.77	8.62	8.70	0.03	< 0.01
FA profile, % of total FA methyl esters						
C10:0	2.16	2.19	2.03	2.21	0.06	0.19
C12:0	3.36	3.21	2.98	3.20	0.06	< 0.01
C14:0	12.2	11.8	12.0	11.5	0.1	< 0.01
C14:1	1.25	1.22	1.38	1.30	0.03	< 0.01
C15:0	1.17	1.13	1.18	1.20	0.02	0.06
C16:0	33.7	32.2	35.6	29.2	0.4	< 0.01
C16:1	1.55	1.59	1.78	1.48	0.05	< 0.01
C17:0	0.60	0.58	0.66	0.62	0.01	< 0.01
C18:0	11.0	11.2	10.2	11.8	0.2	< 0.01
C18:1, <i>cis</i> -9	21.1	21.8	20.7	22.5	0.3	< 0.01
C18:1, <i>trans</i> -11	1.49	1.60	1.54	3.28	0.09	< 0.01
C18:2, <i>cis</i> -9,12	2.22	2.56	1.58	1.57	0.10	< 0.01
C18:2, <i>cis</i> -9, <i>trans</i> -11	0.54	0.58	0.66	1.28	0.03	< 0.01
C18:3, <i>cis</i> -9,12,15	0.34	0.37	0.55	0.63	0.02	< 0.01
C20:0	0.17	0.17	0.19	0.16	0.003	< 0.01
Vitamin and Carotenoid, µg/dL						
α-Tocopherol	107	92	118	150	5	< 0.01
Retinol	36.6	39.5	42.6	38.4	2.0	0.24
β-Carotene	6.6	7.7	12.0	19.2	0.6	< 0.01
Lutein	0.49	0.32	0.68	0.83	0.04	< 0.01

[†] Standard error of the mean corrected replication (n = 31.4)