



HOKKAIDO UNIVERSITY

Title	Dairy intake and the risk of pancreatic cancer : the Japan Collaborative Cohort Study (JACC Study) and meta-analysis of prospective cohort studies
Author(s)	Arafa, Ahmed; Eshak, Ehab Salah; Dong, Jia-Yi et al.
Citation	British journal of nutrition, 128(6), 1147-1155 https://doi.org/10.1017/S0007114521004232
Issue Date	2021-10-20
Doc URL	https://hdl.handle.net/2115/86380
Type	journal article
File Information	dairy-intake-and-the-risk-of-pancreatic-cancer-the-jacc-study-and-meta-analysis-of-prospective-cohort-studies.pdf



Dairy intake and the risk of pancreatic cancer: the JACC Study and meta-analysis of prospective cohort studies

Ahmed Arafa^{1,2}; Ehab S Eshak^{1,3}; Jia-Yi Dong¹; Kokoro Shirai¹; Isao Muraki¹; Hiroyasu Iso¹; Akiko Tamakoshi⁴; the JACC Study Group

¹ Public Health, Department of Social Medicine, Osaka University Graduate School of Medicine, Suita, Japan

² Department of Public Health, Faculty of Medicine, Beni-Suef University, Beni-Suef, Egypt

³ Department of Public Health, Faculty of Medicine, Minia University, El-Minia, Egypt

⁴ Department of Public Health, Graduate School of Medicine, Hokkaido University, Sapporo, Japan

Correspondence: Hiroyasu Iso, MD, Ph.D., MPH, Professor of Public Health, Department of Social Medicine, Osaka University Graduate School of Medicine, Suita-shi, Osaka 565-0871, Japan. Phone: +81-6-6879-3911; Fax: +81-6-6879-3919; E-mail: iso@pbhel.med.osaka-u.ac.jp

Running Title: Dairy intake and pancreatic cancer

Conflict of interest: The authors declare no conflict of interest.



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114521004232

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abstract

Dairy intake was suggested to reduce the risk of gastrointestinal cancers. This study investigated the association between dairy intake and the risk of pancreatic cancer (PAC) using a prospective cohort study and meta-analysis of prospective cohort studies. First, we included 59,774 people aged 40-79 years from the Japan Collaborative Cohort Study (JACC Study). The Cox regression was used to compute the hazard ratios (HRs) and 95% confidence intervals (CIs) of incident PAC for individuals who reported the highest intakes of milk, cheese, and yogurt compared with not consuming the corresponding dairy products. Then, we combined our results with those from other four prospective cohort studies that were eligible after searching several databases, in a meta-analysis, using the fixed-effects model before evaluating publication bias and heterogeneity across studies. In the JACC study, the highest versus no intakes of milk, cheese, and yogurt were not associated with the reduced risk of PAC after a median follow-up of 13.4 years: HRs (95% CIs)= 0.93 (0.64, 1.33), 0.91 (0.51, 1.62), and 0.68 (0.38, 1.21), respectively. The results did not significantly change in the meta-analysis: 0.95 (0.82, 1.11) for milk, 1.16 (0.87, 1.55) for cheese, and 0.91 (0.79, 1.05) for yogurt. The meta-analysis showed no signs of publication bias or heterogeneity across studies. To conclude, consumption of milk, cheese, and yogurt was not associated with the risk of PAC either in the JACC study or the meta-analysis.

Keywords: Milk; Cheese; Yogurt; Cancer; Pancreas; Meta-analysis

1. Introduction

With 458,918 new cases and 432,242 deaths in 2018, pancreatic cancer (PAC) contributed to 2.5% of all-cause cancers and 4.5% of all deaths caused by cancer worldwide. In Japan, a total of 43,119 new cases of PAC and 37,358 related deaths were recorded in the same year representing 4.9% of all-cause incident cancers and 9.1% of all cancer deaths in the country; almost twice proportions as the worldwide incidence and mortality [1, 2]. It is projected that, over the period between 2018 and 2040, PAC incidence and mortality will increase worldwide by 77.7% and 79.9%, respectively [2]. Given its growing incidence and poor five-year survival rate that hardly exceeds 5%, identifying modifiable risk factors for PAC has become a public health priority to apply risk prevention programs [3, 4].

Despite the complex and multifactorial pathogenesis of PAC [5-7], previous research has suggested that dietary factors may play etiological roles [8, 9]. For example, red and processed meat consumption was shown to increase the risk of PAC due to the carcinogenic effects of *N*-nitroso compounds [10-13] while consuming fruits, vegetables, and whole grains reduced that risk because of the anticarcinogenic effects of calcium, magnesium, potassium, alpha- and beta-carotene, and vitamins A, B6, and C contents of these foods [13-17]. In this regard, it could be suggested that the intake of dairy products might be associated with the reduced risk of PAC because they are rich sources of vitamins A, B12, and D as well as calcium, magnesium, and zinc [18-20]. Those vitamins and minerals pose anticarcinogenic properties via inducing cell cycle arrest, apoptosis, and differentiation while suppressing angiogenesis, invasion, and metastasis [21-24]. Further, dairy products include large amounts of lactic acid bacteria and conjugated linoleic acids [25-27]. Lactic acid bacteria help in enhancing the host's immune response, inducing antioxidative and antiproliferative functions, and detoxifying toxicants formed during food processing; characteristics that have been proven to prevent cancer [28-30]. In animal models, lactic acid bacteria were able to alleviate pancreatic inflammation, improve glucose tolerance, and prevent pancreatic damage; factors that are closely related to the risk of PAC [31]. Alike, conjugated linoleic acids were shown in pre-clinical and human studies to have potential anticarcinogenic effects [32, 33]. Using human cells *in vitro*, diets rich in conjugated linoleic acids were shown to reduce PAC penetrance and repress its proliferation [34].

Many case-control studies were conducted to detect the retrospective association between dairy intake and PAC [35-51]. However, in addition to their conflicting findings, the case-control studies were prone to selection bias attributed to the high and rapid fatality of PAC, thus, researchers, to assess dairy intakes, either recruited the survivors who showed low response rates and posed different

sociodemographic and clinical characteristics compared with the deceased or interviewed next-of-kin whom data reliability was considered uncertain. Besides, these studies were subject to bias due to the high possibility of changes in dietary habits among cases after PAC diagnosis. Furthermore, the methodological limitations of these case-control studies did not allow the temporal association between dairy intake and PAC to be investigated [53, 54]. To avoid such biases, the associations between the intakes of different dairy products and the risk of PAC were investigated using a few prospective cohort studies [53-59]. Although the dietary habits of Asian people are substantially different from those in Western countries [60], only two prospective studies assessed the possible association between dairy consumption and the risk of PAC among Asians [58, 59]. Both studies were conducted on Japanese people and were limited by the small number of participants and the lack of representativeness; one study included 11,349 residents of 13 rural areas [58] and the other study included 3158 residents of one prefecture [59].

Since the consumption of dairy products is encouraged in Japan to ensure adequate nutrient intake [61] and given the high incidence of PAC in the country [1, 2] alongside the limitations of previous national studies [58, 59], we used the data of the Japan Collaborative Cohort Study (JACC Study) to investigate the association between the intakes of three dairy products (milk, cheese, and yogurt) and the risk of PAC incidence among a large cohort of middle-aged Japanese. This study primarily hypothesized that the intake of dairy products might be inversely associated with the risk of PAC. Then, we conducted a meta-analysis combining the results of the JACC Study with those from previously published prospective cohort studies.

2. Methods

2.1. The JACC Study

2.1.1. Study population and baseline questionnaire

The JACC Study is a prospective cohort study in which baseline data collection was carried out between 1988 and 1990 in 45 areas in Japan where 110,585 people aged 40-79 years were included. The JACC Study baseline self-administered questionnaire included data about several sociodemographic characteristics, daily walking and leisure physical activity, intakes of common foods and beverages, smoking and alcohol drinking habits, and past medical histories [62, 63]. The follow-up for cancer incidence was conducted in 24 areas using population-based and hospital registries or death certificates before it was terminated by the end of 2009 in four areas, 2008 in two areas, 2006 in two areas, 2003 in one area, 2002 in eight areas, 2000 in one area, 1999 in one area, 1997 in four areas, and 1994 in one area

[62]. Herein, we excluded people with a positive history of cancer before baseline and people who missed reporting on dairy intake. Eventually, the analysis was confined to 59,774 Japanese people who reported at least one of the three questions assessing dairy intake: 58,656 in milk, and 49,302 in cheese, 49,934 in yogurt (Figure 1).

2.1.2. Exposure, outcome, and covariates

Data on dairy intake (exposure) was collected using the self-administered food frequency questionnaire in the JACC Study baseline questionnaire: *“How frequently do you consume the following items?”*. These items included dairy products in the form of *“milk”*, *“cheese”*, and *“yogurt”* among other common foods. The available responses were as follows: *“never”*, *“one to two times/month”*, *“one to two times/week”*, *“three to four times/week”*, and *“almost every day”*. A validation study among a subsample of the JACC Study’s participants showed good validity and reproducibility of the three investigated items; the Spearman rank correlation coefficients between two frequencies assessed twice apart one year were 0.69 for milk, 0.57 for cheese, and 0.54 for yogurt (p-values <0.001) and between the frequencies and the weighed dietary record were 0.65 for milk, 0.44 for cheese, and 0.58 for yogurt (p-values <0.001) [64]. The median portion size of the intakes of the three dairy products per day was estimated in the same validation study and was found to be 146 g for milk, 17 g for cheese, and 98 g for yogurt [64]. Therefore, the five frequencies in our study could be roughly converted into the following amounts: [(milk: 0.0, 6.4, 26.8, 64.0, and 128 g/day), (cheese: 0.0, 0.9, 3.6, 8.5, and 17.0 g/day), and (yogurt: 0.0, 4.9, 21.0, 47.0, and 98.0 g/day)].

On the other hand, the incident cases of PAC (outcome) were diagnosed per the tenth revision of the International Statistical Classification of Diseases and Related Health Problems (C25). Cancer incidence was detected using population-based cancer registries supported by a systematic review of hospital-based cancer registries and inpatients' records of hospitals treating cancer patients [62].

Using the same baseline questionnaire, we collected data on participants' age, sex, weight, height, educational years, perceived stress, smoking and alcohol behaviors, leisure physical activity and walking, history of diabetes, family history of cancer, and daily intakes of several foods that served in calculating daily energy intake (covariates).

2.1.3. Statistical analyses

The age and sex-adjusted p-value for significant differences in the participants' mean values and proportions of sociodemographic characteristics and common risk factors for PAC by their intake of different dairy products were calculated using the linear and logistic regression tests. The Cox

proportional regression was used to compute the hazard ratios (HRs) and their 95% confidence intervals (CIs) of the incidence of PAC for the intakes of milk, cheese, and yogurt. To obtain statistical power, the two highest intake categories “*three to four times/week*” and “*almost every day*” were merged into one category “*≥ three times/week*”.

Person-years of follow-up were calculated from the date of responding to the JACC Study’s baseline questionnaire to the date of PAC diagnosis, death, moving out, or end of the study, whichever came first. The HRs were adjusted for the following variables: age in years, sex (men and women), BMI (<25 and ≥ 25 kg/m²), educational years (<18 and ≥ 18 years), perceived stress (no, mild, moderate, and severe stress), smoking habits (never smokers, former smoker of <20 cigarettes/day, former smoker of ≥ 20 cigarettes/day, current smoker of <20 cigarettes/day, and current smoker of ≥ 20 cigarettes/day), alcohol intake (never, former, and current), leisure sports (never, one to two, three to four, and $\geq$ five hours/week), walking (never, <30 , 30-60, and >60 minutes/day), history of diabetes (yes and no), family history of cancer (yes and no), quartiles of daily intakes of meat, vegetables, and total energy (g/day). Besides, the possibility of interaction with sex, age, smoking, and history of diabetes was examined. SAS version 9.4 software (SAS Institute Inc, Cary, NC) was used for statistical analyses.

2.1.4. Ethical consideration

The research ethics committees of Nagoya University School of Medicine and Osaka University approved the protocol of the JACC study. The study was conducted per the principles of the Declaration of Helsinki.

2.2. The meta-analysis

2.2.1. Literature search

First, we searched MEDLINE (PubMed), Embase, and Web of Science for potential studies published in English before 31/3/2021 (the last day of data search) using the following terms: (Dairy OR Milk OR Cheese OR Yogurt) AND (Cancer). A full search strategy of PubMed was provided (Supplementary file 1). Then, we conducted a manual search of the reference lists of retrieved articles and review articles to obtain additional studies. We reported this meta-analysis according to the checklist of PRISMA [65] and AMSTAR2 [66].

2.2.2. Study selection

Studies were selected for analysis if they met the following criteria: 1) the exposure was milk, cheese, or yogurt intake, 2) the outcome was PAC, and 3) the study design was a prospective cohort. No limitations

were set regarding the year of publication; however, no efforts were made to retrieve unpublished data. The following relevant information was extracted from the included studies: the last name of the first author, year of publication, study name, place of study, age and sex of participants, follow-up years, number of incident cases of PAC, and covariates included in regression models. The multivariable-adjusted HRs with 95% CIs of PAC according to the used categorizations for dairy product intake were also extracted (Supplementary file 2). The quality of studies was determined using the modified Newcastle–Ottawa Scale based on studies' selection (representativeness, selection of the non-exposed, ascertainment of exposure, and demonstration of the outcome), comparability, and outcome (assessment, follow-up length, and adequacy) [67].

2.2.3. Statistical analysis

We used the fixed-effects model to compute the pooled HR with 95% CI of the included studies [68] because the test for heterogeneity was not significant according to the I^2 statistic; a measure of inconsistency across studies [69]. Publication bias was assessed using the regression test for funnel plot asymmetry [70]. All analyses were conducted separately on the following dairy products: milk, cheese, and yogurt. To explore the impact of each study, we performed a sensitivity analysis by removing studies one by one and combining the remainders in separate analyses. R-3.2.0 statistical package (Metafor: A Meta-Analysis Package for R) was used for analysis [71].

3. Results

3.1. The JACC Study

In the JACC Study, participants who reported the intakes of milk, cheese, and yogurt were younger, with lower BMI, more educated, more physically active, and more total energy consumers than their counterparts who reported no intake of the corresponding dairy products (Table 1).

Within a mean follow-up period of 13.0 years (median 13.4 years and maximum 21.6 years), a total of 198 incident cases of PAC were diagnosed. The consumption of the highest versus the lowest amounts of milk, cheese, and yogurt was not associated with the risk of PAC in the age- and sex-adjusted regression models: HRs (95% CI): 0.91 (0.63, 1.33), 1.01 (0.58, 1.78), and 0.73 (0.41, 1.28), respectively. Adjustment for sociodemographic, clinical, and nutritional variables did not change the results: HRs (95% CIs): 0.91 (0.62, 1.33) for milk, 0.91 (0.51, 1.62) for cheese, and 0.68 (0.38, 1.21) for yogurt. The p-values for trend across the increasing frequencies of the three dairy products were statistically insignificant. Also, the p-values for sex, age, smoking, and history of diabetes interactions in the three dairy products were >0.10 (Table 2).

3.2. The meta-analysis

Herein, we combined our results, in a meta-analysis, with the results of the other prospective cohort studies assessing the associations between the intakes of dairy products and the risk of PAC. After omitting irrelevant and retrospective studies, a shortlist of seven prospective cohort studies was obtained [53-59] before three studies in the list were excluded; two studies for publishing more recent results from the same data [53, 54] and one study for defining the exposure as dairy intake as a whole, not as elements of dairy intake [55] (Figure 2). Eventually, four studies were eligible for meta-analysis [56-59] which became five after adding the current JACC Study. Of the four added studies, one study was a pooling of 14 cohorts from North America, Europe, and Oceania [56], one study was conducted in Norway [57], and the remaining two studies were conducted in Japan [58, 59]. Among the five studies included for this meta-analysis, the assessment of dairy products was distributed as follows: milk in five studies, cheese in two studies, and yogurt in four studies. Except for one study that assessed PAC deaths [58], all studies assessed the risk of PAC incidence. Only one study conducted a sex-specific analysis [59] (Table 3). All studies, according to the modified NOS, were of good quality with scores ranging between seven and nine (Supplementary file 3).

In agreement with the results of the JACC study, the pooled HRs (95% CIs) for milk, cheese, and yogurt intakes in the meta-analysis showed no association with the risk of PAC: 0.95 (0.82, 1.11), 1.16 (0.87, 1.55), and 0.91 (0.79, 1.05), respectively. The JACC Study contributed to 16.2%, 24.6%, and 5.7% of the meta-analyses' weights for milk, cheese, and yogurt, respectively. The meta-analyses of the three dairy products showed no heterogeneity across studies ($I^2\% = 0.00$ each). No signs of publication bias were detected in milk and yogurt meta-analyses while conducting the regression test for publication bias in the cheese meta-analysis was unsuitable due to including two studies only (Table 4) (Supplementary file 4). Removing the JACC study from the milk and yogurt meta-analyses did not substantially change the HRs (95% CIs): 0.96 (0.81, 1.14) and 0.93 (0.80, 1.07), respectively. The sensitivity analyses by leaving out studies one by one and combining the remainders did not affect the conclusion (Supplementary file 5).

4. Discussion

The JACC Study indicated that, within a mean follow-up period of 13.0 years (median 13.4 years), the intakes of milk, cheese, and yogurt were not associated with the risk of PAC among middle-aged Japanese, and no dose-response associations were noticed. Combining the results of the JACC Study with those from other prospective cohort studies, in a meta-analysis, did not materially change the findings.

The World Cancer Research Fund and the American Institute of Cancer Research, based on limited evidence, labeled the possible association between consumption of dairy products and the risk of PAC as “*limited/non-conclusive*”. We could not find any association between consuming dairy products and the risk of PAC [72].

Of note, the JACC Study included numerous strengths such as investigating the relationship between consuming several dairy products and the risk of PAC among a large study population, excluding participants with a history of cancer, using a prospective cohort design and lengthy follow-up period, assessing dairy intake using a validated food frequency sheet, and adjusting the results for most potential confounders. Still, the JACC Study carried some limitations that should be addressed. First, the number of incident cases of PAC was limited that it did not allow us to stratify the results by potential risk factors for PAC such as sex, age, smoking behavior, and history of diabetes. Yet, this limitation was partly solved by combining the results of the JACC Study with the results of the other four prospective cohort studies via a meta-analysis. Besides, formal interaction tests showed that age, sex, smoking, and diabetes did not affect the associations. Second, we obtained no data about the histopathological classifications of PAC cases and their treatment protocols after diagnosis. Third, data on dairy consumption was collected more than 30 years ago. Although dairy consumption per capita in Japan has been increasing since then, the current consumption of dairy products in Japan is much lower than the recommended levels [73, 74]. Fourth, it could be speculated that the variation in the PAC ascertainment time across areas because of their different termination times (1994-2009) might have affected the PAC incidence. Yet, the age-adjusted PAC incidence and attributed mortality did not significantly change during this period [75]. Fifth, this study focused on investigating the intake of dietary products as a whole rather than their nutrients, however, the intakes of vitamin D and calcium were shown in previous studies to be not associated with the reduced risk of PAC [56, 76, 77].

In addition, our meta-analysis posed several strengths such as augmenting the number of incident PAC cases, limiting the inclusion criteria to prospective cohort studies that avoided the methodological

limitations of previous case-control studies, including studies of good quality according to the modified Newcastle–Ottawa Scale, and showing no signs of heterogeneity across studies or publication bias.

However, this meta-analysis had some limitations. First, the JACC Study and Genkiger et al study [56] together contributed to most incident cases of PAC and weights of the meta-analyses that were limited by the small number of included studies especially in terms of cheese and yogurt. Second, since exposure was self-reported in all studies, the possibility of non-differential misclassification bias cannot be entirely excluded. Third, it could be argued that two included studies [57, 58] did not adjust their results for smoking; a major risk factor for PAC [78], however, both studies contributed to small fractions of the meta-analyses weights. Moreover, adjusting for smoking and other sociodemographic, clinical, and nutritional factors in the JACC study did not materially change the results. Fourth, the included studies used different categories for dairy consumption. For example, the highest consumption categories of milk, cheese, and yogurt in the Genkiger et al study [56] were ≥ 500 , ≥ 50 , and ≥ 57 g/day compared with ≥ 64 , ≥ 8.5 , and ≥ 47 g/day in the JACC Study, respectively. In the previous Japanese studies, Matsumoto et al [58] assessed the risk among everyday consumers versus not everyday consumers while Khan et al [59] compared consuming more than to equal or less than several times per month, and both studies, however, did not calculate the consumed amounts of dairy products. Lastly, our meta-analysis protocol was not *a priori* registered, although we performed the meta-analysis in a standard way.

In conclusion, consumption of milk, cheese, and yogurt was found to be not associated with the risk of PAC among middle-aged Japanese in the JACC Study and the results did not change in the meta-analyses of prospective cohort studies. Since consuming dairy products was shown to have no role in reducing the risk of PAC, identifying other modifiable risk factors for PAC is important to reduce its burden.

5. Financial Support

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT) (Monbusho); Grants-in-Aid for Scientific Research on Priority Areas of Cancer; and Grants-in-Aid for Scientific Research on Priority Areas of Cancer Epidemiology from MEXT (MonbuKagaku-sho) (Nos. 61010076, 62010074, 63010074, 1010068, 2151065, 3151064, 4151063, 5151069, 6279102, 11181101, 17015022, 18014011, 20014026, 20390156, and 26293138), Comprehensive Research on Cardiovascular and Life-Style Related Diseases (H26-Junkankitou [Seisaku]-Ippan-001 and H29-Junkankitou [Seishuu]-Ippan-003), JSPS KAKENHI Grant Number JP 16H06277, and Grants-in-Aid for China Scholarship Council (CSC file No. 201608050-113).

6. Conflict of interest

The authors declare no conflict of interest.

7 Contributions

AA (Conceptualization), HI and AT (Resources), HI and AT (Funding acquisition), AA, EE, JD, KS, IM, HI, and AT (Visualization), AA (Review literature), AA (Draft writing), AA (Data analysis), EE, JD, and HI (Supervision), AA, EE, JD, KS, IM, HI, and AT (Critical revision and editing).

8 References

1. Bray F, Ferlay J, Soerjomataram I, Siegel R, Torre L, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018, 68:394-424.
2. Rawla P, Sunkara T, Gaduputi V. Epidemiology of pancreatic cancer: global trends, etiology and risk factors. *World J Oncol* 2019, 10:10-27.
3. Del Chiaro M, Segersvard R, Lohr M, Verbeke C. Early detection and prevention of pancreatic cancer: is it really possible today? *World J Gastroenterol* 2014, 20:12118-12131.
4. Arafa A, Eshak E, Abdel Rahman T, Anwar M. Hepatitis C virus infection and risk of pancreatic cancer: A meta-analysis. *Cancer Epidemiol* 2020, 65:101691.
5. Klein A. Identifying people at a high risk of developing pancreatic cancer. *Nat Rev Cancer* 2013, 13:66-74.
6. McGuigan A, Kelly P, Turkington R, Jones C, Coleman H, McCain R. Pancreatic cancer: A review of clinical diagnosis, epidemiology, treatment and outcomes. *World J Gastroenterol* 2018, 24:4846-4861.
7. Muñoz A, Chakravarthy D, Gong J, Halff G, Ghosh R, Kumar A. Pancreatic cancer: Current status and Challenges. *Curr Pharmacol Rep* 2017, 3:396-408.
8. Casari I, Falasca M. Diet and pancreatic cancer prevention. *Cancers (Basel)*. 2015, 7:2309-2317.
9. Pericleous M, Rossi R, Mandair D, Whyand T, Caplin M. Nutrition and pancreatic cancer. *Anticancer Res* 2014, 34:9-21.
10. Sen N, Seaman S, Burgess C, Baddoo P, Weber D. Investigation on the possible formation of N-nitroso-N-methylurea by nitrosation of creatinine in model systems and in cured meats at gastric pH. *J Agric Food Chem* 2000, 48:5088-5096.
11. Larsson S, Wolk A. Red and processed meat consumption and risk of pancreatic cancer: meta-analysis of prospective studies. *Br J Cancer* 2012, 106:603-607.
12. Zhao Z, Yin Z, Pu Z, Zhao Q. Association between consumption of red and processed meat and pancreatic cancer risk: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2017, 15:486-493.
13. Paluszkiewicz P, Smolinska K, Debinska I, Turski W. Main dietary compounds and pancreatic cancer risk. The quantitative analysis of case-control and cohort studies. *Cancer Epidemiol* 2012, 36:60-67.


14. Jansen R, Robinson D, Stolzenberg-Solomon R, Bamlet W, de Andrade M, Oberg A, et al. Nutrients from fruit and vegetable consumption reduce the risk of pancreatic cancer. *J Gastrointest Cancer* 2013, 44:152-161.
15. Lei Q, Zheng H, Bi J, Wang X, Jiang T, Gao X, et al. Whole grain intake reduces pancreatic cancer risk: a meta-analysis of observational studies. *Medicine* 2016, 95:e2747.
16. Li L, Luo Y, Lu M, Xu X, Lin H, Zheng Z. Cruciferous vegetable consumption and the risk of pancreatic cancer: a meta-analysis. *World J Surg Oncol* 2015, 13:44.
17. Wu Q-J, Wu L, Zheng L-Q, Xu X, Ji C, Gong T-T. Consumption of fruit and vegetables reduces risk of pancreatic cancer: evidence from epidemiological studies. *Eur J Cancer Prev* 2016, 25:196-205.
18. Bao Y, Ng K, Wolpin B, Michaud D, Giovannucci E, Fuchs C. Predicted vitamin D status and pancreatic cancer risk in two prospective cohort studies. *Br J Cancer* 2010, 102:1422-1427.
19. Lampe J. Dairy products and cancer. *J Am Coll Nutr* 2011, 30:464-470.
20. Skinner H, Michaud D, Giovannucci E, Willett W, Colditz G, Fuchs C. Vitamin D intake and the risk for pancreatic cancer in two cohort studies. *Cancer Epidemiol Biomarkers Prev* 2006, 15:1688-1695.
21. Donaldson M. Nutrition and cancer: A review of the evidence for an anti-cancer diet. *Nutr J* 2004, 3:19.
22. Gross M. Vitamin D and calcium in the prevention of prostate and colon cancer: new approaches for the identification of needs. *J Nutr* 2005, 135:326-331.
23. Gaziano M, Sesso H, Christen W, Bubes V, Smith J, MacFadyen J, et al. Multivitamins in the prevention of cancer in men: The Physicians' Health Study II randomized controlled trial. *JAMA* 2012, 308:1871-1880.
24. Ma Y, Johnson C, Trump D. Mechanistic insights of vitamin d anticancer effects. *Vitam Horm* 2016, 100:395-431.
25. Buttriss J. Nutritional properties of fermented milk products. *Int J Dairy Tech* 1997, 50:21-27.
26. Nunes J, Silva M, Perrone D, Torres A. Seasonal variation in fat quality and conjugated linoleic acid content of dairy products from the Tropics: Evidence of potential impact on human health. *Foods* 2017, 6:61.

27. Xu S, Boylston T, Glatz B. Conjugated linoleic acid content and organoleptic attributes of fermented milk products produced with probiotic bacteria. *J Agric Food Chem* 2005, 53:9064-9072.
28. Ahmadi M, Ebrahimi M, Mehrabian S, Tafvizi F, Bahrami H, Dameshghian M. Antimutagenic and anticancer effects of lactic acid bacteria isolated from Tarhana through Ames test and phylogenetic analysis by 16S rDNA. *Nutr Cancer* 2014, 66:1406-1413.
29. Kim J, Kim J, Lee K, Lee H. Cancer chemopreventive effects of lactic acid bacteria. *J Microbiol Biotechnol* 2007, 17:1227-1235.
30. Shoukat S. Potential anti-carcinogenic effect of probiotic and lactic acid bacteria in detoxification of benzo[a]pyrene: A review. *Trends Food Sci Tech* 2020, 99:450-459.
31. Wang G, Si Q, Yang S, Jiao T, Zhu H, Tian P, et al. Lactic acid bacteria reduce diabetes symptoms in mice by alleviating gut microbiota dysbiosis and inflammation in different manners. *Food Funct* 2020, 11:5898-5914.
32. den Hartigh L. Conjugated linoleic acid effects on cancer, obesity, and atherosclerosis: a review of pre-clinical and human trials with current perspectives. *Nutrients* 2019, 11:370.
33. Whigham L, Cook M, Atkinson R. Conjugated linoleic acid: implications for human health. *Pharmacol Res* 2000, 42:503-510.
34. Ding Y, Mullapudi B, Torres C, Mascariñas E, Mancinelli G, Diaz A, et al. Omega-3 fatty acids prevent early pancreatic carcinogenesis via repression of the AKT pathway. *Nutrients* 2018, 10:1289.
35. Gold E, Gordis L, Diener M, Seltser R, Boitnott J, Bynum T, et al. Diet and other risk factors for cancer of the pancreas. *Cancer* 1985, 55:460-467.
36. Norell S, Ahlbom A, Erwald R, Jacobson G, Lindberg-Navier I, Olin R, et al. Diet and pancreatic cancer: a case-control study. *Am J Epidemiol* 1986, 124:894-902.
37. Falk R, Pickle L, Fontham E, Correa P, Fraumeni J Jr. Life-style risk factors for pancreatic cancer in Louisiana: a case-control study. *Am J Epidemiol* 1988, 128:324-336.
38. La Vecchia C, Negri E, D'Avanzo B, Ferraroni M, Gramenzi A, Savoldelli R, et al. Medical history, diet and pancreatic cancer. *Oncology* 1990, 47:463-466.
39. Baghurst P, McMichael A, Slavotinek A, Baghurst K, Boyle P, Walker A. A case-control study of diet and cancer of the pancreas. *Am J Epidemiol* 1991, 134:167-179.

40. Bueno de Mesquita H, Maisonneuve P, Runia S, Moerman C. Intake of foods and nutrients and cancer of the exocrine pancreas: a population-based case-control study in The Netherlands. *Int J Cancer* 1991, 48:540-549.
41. Olsen G, Mandel J, Gibson R, Wattenberg L, Schuman L. Nutrients and pancreatic cancer: a population-based case-control study. *Cancer Causes Control* 1991, 2:291-297.
42. Mizuno S, Watanabe S, Nakamura K, Omata M, Oguchi H, Ohashi K, et al. A multi-institute case-control study on the risk factors of developing pancreatic cancer. *Jpn J Clin Oncol* 1992, 22:286-291.
43. Lyon J, Slattery M, Mahoney A, Robison L. Dietary intake as a risk factor for cancer of the exocrine pancreas. *Cancer Epidemiol Biomarkers Prev* 1993, 2:513-518.
44. Ohba S, Nishi M, Miyake H. Eating habits and pancreas cancer. *Int J Pancreatol* 1996, 20:37-42.
45. Silverman D, Swanson C, Gridley G, Wacholder S, Greenberg R, Brown L, et al. Dietary and nutritional factors and pancreatic cancer: a case-control study based on direct interviews. *J Natl Cancer Inst* 1998, 90:1710-1719.
46. Soler M, Chatenoud L, La Vecchia C, Franceschi S, Negri E. Diet, alcohol, coffee and pancreatic cancer: final results from an Italian study. *Eur J Cancer Prev* 1998, 7:455-460.
47. Abdelrehim M, Mahfouz E, Ewis A, Seedhom A, Afifi H, Shebl F. Dietary factors associated with pancreatic cancer risk in Minia, Egypt: Principal component analysis. *Asian Pac J Cancer Prev* 2018, 19:449-455.
48. Ghorbani Z, Hekmatdoost A, Zinab H, Farrokhzad S, Rahimi R, Malekzadeh R, et al. Dietary food groups intake and cooking methods associations with pancreatic cancer: a case-control study. *Indian J Gastroenterol* 2015, 34:225-232.
49. Liu S, Chen W, Wang N, Yin M, Sun X, He Y. Dietary factors and risk of pancreatic cancer: a multi-centre case-control study in China. *Asian Pac J Cancer Prev* 2014, 15:7947-7950.
50. Nkondjock A, Krewski D, Johnson K, Ghadirian P; Canadian Cancer Registries Epidemiology Research Group. Dietary patterns and risk of pancreatic cancer. *Int J Cancer* 2005, 114:817-823.
51. Chan J, Wang F, Holly E. Pancreatic cancer, animal protein and dietary fat in a population-based study, San Francisco Bay Area, California. *Cancer Causes Control* 2007, 18:1153-1167.
52. Howe G, Burch J. Nutrition and pancreatic cancer. *Cancer Causes Control* 1996, 7:69-82.
53. Stolzenberg-Solomon R, Pietinen P, Taylor P, Virtamo J, Albanes D. Prospective study of diet and pancreatic cancer in male smokers. *Am J Epidemiol* 2002, 155:783-792.

54. Michaud D, Giovannucci E, Willett W, Colditz G, Fuchs C. Dietary meat, dairy products, fat, and cholesterol and pancreatic cancer risk in a prospective study. *Am J Epidemiol* 2003, 157:1115-1125.
55. Nöthlings U, Wilkens L, Murphy S, Hankin J, Henderson B, Kolonel L. Meat and fat intake as risk factors for pancreatic cancer: the multiethnic cohort study. *J Natl Cancer Inst* 2005, 97:1458-1465.
56. Genkinger J, Wang M, Li R, Albanes D, Anderson K, Bernstein L, et al. Dairy products and pancreatic cancer risk: a pooled analysis of 14 cohort studies. *Ann Oncol* 2014, 25:1106-1115.
57. Ursin G, Bjelke E, Heuch I, Vollset S. Milk consumption and cancer incidence: a Norwegian prospective study. *Br J Cancer* 1990, 61:454-459.
58. Matsumoto M, Ishikawa S, Nakamura Y, Kayaba K, Kajii E. Consumption of dairy products and cancer risks. *J Epidemiol* 2007, 17:38-44.
59. Khan M, Goto R, Kobayashi K, Suzumura S, Nagata Y, Sonoda T, et al. Dietary habits and cancer mortality among middle aged and older Japanese living in Hokkaido, Japan by cancer site and sex. *Asian Pac J Cancer Prev* 2004, 5:58-65.
60. Tsugane S, Sasaki S, Kobayashi M, Tsubono Y, Sobue T. Dietary habits among the JPHC study participants at baseline survey. Japan Public Health Center-based Prospective Study on Cancer and Cardiovascular Diseases. *J Epidemiol* 2001, 11: 30-43.
61. Saito A, Okada E, Tarui I, Matsumoto M, Takimoto H. The Association between milk and dairy products consumption and nutrient intake adequacy among Japanese adults: Analysis of the 2016 National Health and Nutrition Survey. *Nutrients* 2019, 11:2361.
62. Mori M, Sakauchi F, Washio M, Ozasa K, Watanabe Y, Yoshimura T, et al. Survey for incidence of cancer as a measure of outcome in the JACC study. *J Epidemiol* 2005, 15: 80-85.
63. Tamakoshi A, Ozasa K, Fujino Y, Suzuki K, Sakata K, Mori M, et al. Cohort profile of the Japan Collaborative Cohort Study at final follow-up. *J Epidemiol* 2013, 23:227-232.
64. Date C, Fukui M, Yamamoto A, Wakai K, Ozeki A, Motohashi Y, et al. Reproducibility and validity of a self-administered food frequency questionnaire used in the JACC study. *J Epidemiol* 2003, 15: 9-23.
65. Moher D, Liberati A, Tetzlaff J, Altman D; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009, 339:2535.

66. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 2017, 358:j4008.
67. Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analyses. www.ohri.ca/programs/clinical_epidemiology/oxford.htm. Accessed on 9/9/2020.
68. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986, 7:177-188.
69. Higgins J, Thompson S, Deeks J, Altman D. Measuring inconsistency in meta-analyses. *BMJ* 2003, 327:557-560.
70. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997, 315:629-634.
71. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Software* 2010, 36:1-48.
72. World Cancer Research Fund (WCRF) and American Institute for Cancer Research Continuous Update Project Report. Food, nutrition, physical activity, and the prevention of pancreatic cancer, 2012. <https://www.wcrf.org/dietandcancer/pancreatic-cancer>. Accessed on 25/2/2021.
73. Ministry of Health Labour and Welfare The National Health and Nutrition Survey in Japan. <http://www.mhlw.go.jp/bunya/kenkou/eiyou/h28-houkoku.html>. Accessed on 28/6/2021.
74. Saito A, Okada E, Tarui I, Matsumoto M, Takimoto H. The association between milk and dairy products consumption and nutrient intake adequacy among Japanese adults: Analysis of the 2016 National Health and Nutrition Survey. *Nutrients* 2019,11: 2361.
75. Lin Y, Tamakoshi A, Wakai K, Kawamura T, Aoki R, Kojima M, Ohno Y. Descriptive epidemiology of pancreatic cancer in Japan. *J Epidemiol* 1998, 8: 52-9.
76. Fan H, Yu Y, Nan H, Hoyt M, Reger MK, Prizment A, et al. Associations between intake of calcium, magnesium and phosphorus and risk of pancreatic cancer: a population-based, case-control study in Minnesota. *Br J Nutr* 2021, 1-9.
77. Zhang X, Huang XZ, Chen WJ, Wu J, Chen Y, Wu CC, et al. Plasma 25-hydroxyvitamin D levels, vitamin D intake, and pancreatic cancer risk or mortality: a meta-analysis. *Oncotarget* 2017, 8: 64395-64406.
78. Pandol S, Apte M, Wilson J, Gukovskaya A, Edderkaoui M. The burning question: Why is smoking a risk factor for pancreatic cancer? *Pancreatology* 2012, 12:344-349.

Figure legends:**Figure 1:** Flowchart of the included participants in the JACC study




Figure 2: Prisma chart of the included prospective cohort studies in the meta-analysis

Table 1. Age-sex-adjusted sociodemographic characteristics of participants according to their dairy intakes of milk, cheese, and yogurt (JACC Study)

	Never	1-2 times/month	1-2 times/week	≥3 times/week
Milk				
Study population	10,491	4627	8065	35,473
Age, years*	57.9 (10.1)	56.3 (10.3)	56.1 (10.2)	58.0 (10.1)
Men, %	44.4	47.8	41.8	38.3
Body mass index, kg/m ² *	22.8 (3.2)	22.9 (3.0)	22.9 (3.0)	22.7 (2.9)
Education, years*	16.3 (2.3)	16.6 (2.3)	16.8 (2.2)	16.9 (2.3)
Perceived high stress, %	16.5	15.2	14.3	15.5
Current smoking, %	30.1	30.3	25.9	19.6
Current drinking, %	42.2	48.8	44.0	41.7
No leisure sport, %	70.1	66.6	65.7	62.9
No walking, %	9.7	8.5	8.8	9.0
History of diabetes, %	3.9	4.2	3.8	5.8
Vegetable intake, gm/day	239.2 (307.5)	211.9 (292.0)	236.5 (301.5)	301.9 (330.1)
Meat intake, gm/day	25.6 (20.0)	26.4 (18.8)	29.2 (19.4)	31.0 (20.6)
Energy intake, Kcal/day*	1450.4 (451.5)	1478.4 (455.5)	1495.3 (439.7)	1561.9 (416.9)
Family history of cancer, %	7.2	7.5	5.3	5.6
Cheese				
Study population	25,318	13,255	7007	3722

Age, years*	58.0 (10.0)	54.7 (9.7)	55.5 (10.1)	57.8 (9.9)
Men, %	39.4	43.6	41.3	38.6
Body mass index, kg/m ² *	22.9 (3.1)	22.9 (2.9)	22.7 (2.8)	22.5 (2.9)
Education, years*	16.5 (2.3)	17.2 (2.2)	17.3 (2.3)	17.4 (2.4)
Perceived high stress, %	17.0	16.3	17.3	19.2
Current smoking, %	23.5	24.8	22.6	21.1
Current drinking, %	39.3	49.4	46.6	44.1
No leisure sport, %	73.5	67.9	64.5	61.7
No walking, %	10.2	8.6	8.3	7.8
History of diabetes, %	5.2	4.5	4.0	4.9
Vegetable intake, gm/day	247.2 (309.5)	261.4 (312.9)	336.2 (338.5)	380.6 (352.8)
Meat intake, gm/day	25.3 (18.6)	30.6 (18.4)	36.1 (20.7)	39.6 (27.1)
Energy intake, Kcal/day*	1458.0 (421.9)	1549.5 (422.9)	1637.8 (425.0)	1720.9 (455.9)
Family history of cancer, %	5.1	3.5	3.1	2.3
Yogurt				
Study population	28,615	9032	6555	5732
Age, years*	57.1 (9.9)	55.5 (10.1)	56.0 (10.4)	58.5 (10.1)
Men, %	47.4	33.4	28.8	30.6
Body mass index, kg/m ² *	22.7 (3.0)	22.8 (2.9)	22.7 (2.9)	22.6 (2.9)
Education, years*	16.5 (2.3)	17.1 (2.3)	17.2 (2.3)	17.1 (2.4)
Perceived high stress, %	14.0	12.7	13.1	15.3
Current smoking, %	28.1	18.8	15.9	15.7

Current drinking, %	46.5	41.9	37.1	37.8
No leisure sport, %	67.2	62.7	61.8	59.8
No walking, %	10.4	8.8	8.8	9.5
History of diabetes, %	4.9	4.6	4.3	5.6
Vegetable intake, gm/day	246.8 (308.9)	265.2 (314.4)	318.6 (333.4)	365.8 (349.4)
Meat intake, gm/day	27.3 (19.4)	30.3 (19.1)	33.4 (20.6)	34.2 (24.5)
Energy intake, Kcal/day*	1519.8 (446.4)	1497.4 (410.0)	1513.6 (20.6)	1568.5 (420.2)
Family history of cancer, %	7.8	5.7	4.8	4.1

*Mean (standard deviation)

Table 2. The associations between dairy intakes of milk, cheese, and yogurt and the risk of pancreatic cancer (JACC Study)

	Never	1-2 times/month	1-2 times/week	≥3 times/week	P for trend
Milk					
Person-years	135,000	60,683	108,155	459,864	---
Total population	10,491	4627	8065	35,473	---
Incident cases	37	19	22	113	---
Model I	1	1.25 (0.72, 2.18)	0.83 (0.49, 1.41)	0.91 (0.63, 1.33)	0.294
Model II	1	1.22 (0.70, 2.13)	0.81 (0.47, 1.37)	0.91 (0.62, 1.33)	0.308
Cheese					
Person-years	327,950	182,921	98,716	50,613	---
Total population	25,318	13,255	7007	3722	---
Incident cases	90	29	26	14	---
Model I	1	0.70 (0.46, 1.06)	1.08 (0.70, 1.68)	1.01 (0.58, 1.78)	0.596
Model II	1	0.67 (0.43, 1.02)	1.00 (0.64, 1.58)	0.91 (0.51, 1.62)	0.770
Yogurt					
Person-years	367,641	116,267	84,128	71,809	---
Total population	28,615	9032	6555	5732	---

Incident cases	92	20	11	14	---
Model I	1	0.78 (0.48, 1.27)	0.57 (0.30, 1.06)	0.73 (0.41, 1.28)	0.179
Model II	1	0.76 (0.47, 1.24)	0.55 (0.29, 1.03)	0.68 (0.38, 1.21)	0.137

Model I: Adjusted for age and sex

Model II: Adjusted further for body mass index, education, perceived stress, smoking behavior, alcohol, leisure sport, walking, history of diabetes, family history of cancer, and total meat, vegetables, and energy intake

P-values for sex interaction (milk= 0.80, cheese= 0.85, and yogurt= 0.25)

P-values for age interaction (milk= 0.20, cheese= 0.94, and yogurt= 0.94)

P-values for smoking interaction (milk= 0.30, cheese= 0.11, and yogurt= 0.40)

P-values for history of diabetes interaction (milk= 0.44, cheese= 0.92, and yogurt= 0.42)

Table 3: Summary of the prospective cohort studies included in the meta-analysis and investigating the associations between dairy intake and the risk of pancreatic cancer

Study ID	Description	Country	Exposure (groups)	Cancer cases	Covariates
Genkinger (2014)	A pooling of ATBC, BCDDP, CNBSS, CPS-II, CTS, COSM, HPFS, IWHS, MCCS, NLCS, NYSC, NHS, PLCO, and SMC Men and women aged 15-107 years and followed up for a maximum of 7-20 years	US, Canada, Finland, Sweden, Netherlands, Australia	Whole milk, cheese, yogurt, and ice cream (highest versus lowest amount intake)	2212 (Incidence)	Age, year of questionnaire return, sex, BMI, smoking, alcohol, diabetes, and energy intake
Ursin (1990)	Men and women aged 35-74 years and followed up for a maximum of 11 years	Norway	Milk (≥ 2 glasses/day versus < 1 glass/day)	62 (Incidence)	Age, sex, and residence
Matsumoto (2007)	JMS Men and women aged 18-90 years and followed up for an average of 9.2 years	Japan	Milk, yogurt, and butter (every day versus not every day)	13 (Mortality)	Age and sex
Khan (2004)	Hokkaido Women aged ≥ 40 and followed up for an average of 14.8 years	Japan	Milk, yogurt, and butter or margarine (several times/week and every day versus never, several times/year, and	13 (Incidence)	Age, health status, health education, health screening, and smoking
	Hokkaido Men aged ≥ 40 and followed up for an average of 13.8 years			12 (Incidence)	Age and smoking

			several times/month)		
This study (2021)	JACC Men and women aged 40-79 years and followed up for an average of 12.8 years	Japan	Milk, cheese, and yogurt (highest ≥ 3 times/week versus no intake)	198 (Incidence)	Age, sex, BMI, education, stress, smoking, alcohol, leisure physical activity, walking, diabetes, family history of cancer, and intakes of meat, vegetable, and energy

Alpha-Tocopherol Beta-Carotene Cancer Prevention Study (**ATBC**); Breast Cancer
Detection Demonstration Project Follow-up Study (**BCDDP**); Canadian National
Breast Screening Study (**CNBSS**); Cancer Prevention Study II Nutrition Cohort
(**CPS-II**); California Teachers Study (**CTS**); Cohort of Swedish Men (**COSM**);
Health Professionals Follow-up Study (**HPFS**); Iowa Women's Health Study
(**IWHS**); Japan Collaborative Cohort Study (**JACC**); Jichi Medical School Cohort
Study (**JMS**); Melbourne Collaborative Cohort Study (**MCCS**); The Netherlands
Cohort Study (**NLCS**); New York State Cohort (**NYSC**); Nurses' Health Study
(**NHS**); Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial (**PLCO**); and
the Swedish Mammography Cohort (**SMC**)

Table 4: Meta-analysis showing hazard ratios (95% confidence intervals), weights, and heterogeneity and publication bias across studies investigating the association between the intakes of milk, cheese, and yogurt and the risk of pancreatic cancer

Author	Milk		Cheese		Yogurt	
	HR (95% CI)	Weight %	HR (95% CI)	Weight %	HR (95% CI)	Weight %
Genkinger	0.98 (0.82, 1.18)	71.09	1.26 (0.91, 1.76)	75.43	0.93 (0.81, 1.08)	92.43
Ursin	0.71 (0.41, 1.25)	7.58	---	---	---	---
Matsumoto	0.97 (0.33, 2.90)	1.99	---	---	1.19 (0.15, 9.10)	0.45
Khan (Women)	1.20 (0.40, 4.10)	1.74	---	---	0.90 (0.20, 3.90)	0.87
Khan (Men)	1.40 (0.40, 5.20)	1.43	---	---	0.50 (0.10, 4.20)	0.55
This study	0.91 (0.62, 1.33)	16.17	0.91 (0.51, 1.62)	24.57	0.68 (0.38, 1.21)	5.70
Overall	0.95 (0.82, 1.11)	100.00	1.16 (0.87, 1.55)	100.00	0.91 (0.79, 1.05)	100.00
I^2 (p-value for heterogeneity)	0.00 (0.887)		0.00 (0.338)		0.00 (0.823)	
Z (p-value for publication bias)	0.025 (0.980)		---		-0.643 (0.520)	