



Title	Saponins are responsible for the anti-obesogenic activity of <i>Acacia concinna</i>
Author(s)	Zhuoyue, Zhao; Ruangaram, Wijitrapha; Kato, Eisuke
Citation	Journal of natural medicines, 75, 1005-1013 https://doi.org/10.1007/s11418-021-01530-0
Issue Date	2021-09
Doc URL	https://hdl.handle.net/2115/86646
Rights	This is a post-peer-review, pre-copyedit version of an article published in Journal of natural medicines. The final authenticated version is available online at: http://dx.doi.org/10.1007/s11418-021-01530-0
Type	journal article
File Information	Eikato_JNatProd_2021.pdf



Saponins are responsible for the anti-obesogenic activity of *Acacia concinna*

Zhao Zhuoyue¹, Wijitrapha Ruangaram¹, Eisuke Kato^{2*}

1 Division of Applied Bioscience, Graduate School of Agriculture, Hokkaido University, Kita-ku, Sapporo, Hokkaido 060-8589, Japan

2 Division of Fundamental AgriScience and Research, Research Faculty of Agriculture, Hokkaido University, Kita-ku, Sapporo, Hokkaido 060-8589, Japan

*Tel: +81-11-706-2496, E-mail: eikato@chem.agr.hokudai.ac.jp

Abstract

Acacia concinna (Willd.) DC. is a medicinal plant sourced mainly from Southeast Asia. The pods of *Acacia concinna* (*A. concinna*) are a potential candidate to treat or prevent obesity; however, these medicinal attributes have not been examined in detail. In this study, the anti-obesogenic compounds in *A. concinna* pods were investigated. Chromatographic separation of the pod extract guided by pancreatic lipase inhibitory activity led to the isolation of saponins. Decomposition analysis of the saponins revealed their chemical composition to be acacic acid, monoterpenes and five types of sugars (glucose, xylose, rhamnose, quinovose, arabinose). The predicted structures of the saponins from decomposition analysis were confirmed by LC-MS analysis, showing that these saponins are mixture of various derivatives of monoterpenes and sugar units. These saponins inhibited pancreatic lipase activity strongly with an IC_{50} of 7.9 $\mu\text{g/mL}$, and reduced lipid accumulation in 3T3-L1 adipocytes at 6.3 $\mu\text{g/mL}$. The saponins also enhanced lipolysis of 3T3-L1 adipocytes at 3.1 or 6.3 $\mu\text{g/mL}$ by mediating the activity of protein kinase A and extracellular signal-regulated kinase pathways, suggesting that this mechanism is partly responsible for the observed reduction of lipid content in adipocytes. The results underline *A. concinna* as a potential source of the anti-obesogenic candidates for the future treatment and prevention of obesity.

Keywords

Acacia concinna, saponin, obesity, pancreatic lipase, lipolysis

Introduction

Acacia concinna (Wild.) DC., also known as Soap-Pod or Shikakai, belongs to the family of Fabaceae. The plant is cultivated throughout Asian countries including India, Sri Lanka, Myanmar and Thailand [1, 2]. Several parts of *Acacia concinna* (*A. concinna*) are used as traditional medicines. Infusion of the leaves is used to treat fever caused by malaria infections, the legume fruits are used for antidandruff and hair strengthening purposes, and the pods have a laxative effect, relieve coughing and reduce fevers [3, 4]. Accordingly, studies on *A. concinna* have reported potential anti-fungal and anti-dermatophyte activity [1, 4].

The growing number of overweight and obese people represents a major health crisis. Since 1975, the global number of people who are overweight and obese has tripled. Numerous modern medicines have been developed to treat overweight and obese people, and the use of medicinal plants as a natural source for treating obesity is an option [5]. Although medicinal plants generally have milder responses when compared with those of modern medicines, the smaller number of side effects because of their milder activity and affordability are attractive attributes that make these plants potentially suitable for broad, global use in preventing and treating obesity.

Our previous screening study identified *A. concinna* as a promising source for candidate to prevent and treat obesity [6]. *A. concinna* exhibited multiple anti-obesogenic activities including pancreatic lipase inhibition, enhanced lipolysis activity and a reduction in lipid accumulation. However, the active components responsible for these activities have not been investigated in detail. In this study, we identified the active anti-obesogenic components in *A. concinna* and evaluated their anti-obesogenic potential.

Results

The dried pods of *A. concinna* were extracted, partitioned between solvents and repeatedly separated by column chromatography guided by the pancreatic lipase inhibitory activity to obtain a bioactive

fraction. Further separation of the obtained bioactive fraction was initially attempted by using HPLC; however, a single broad peak was observed, indicating that separation of the components in this fraction was not feasible by this approach. The NMR spectrum of the bioactive fraction gave a relatively well-dispersed but complex $^1\text{H-NMR}$ spectrum and a wide range of m/z ions were observed by MS (Supplementary Figs. S1 and S2). The m/z ions between 2000-2500 together with the HPLC result suggested presence of compounds with similar structures. The $^1\text{H-NMR}$ signals between 0.8-2.0 ppm and 3.0-4.0 ppm suggested a triterpene moiety and sugar units indicating that this fraction is a mixture of saponins (AC saponin).

To obtain structural information about AC saponin, this sample was subjected to decomposition and the resulting products were analyzed. Alkaline hydrolysis gave compounds **1–4** as identifiable products. NMR analysis revealed these four compounds as monoterpenes conjugated with either quinovose (Qui) or xylose (Xyl) (Fig. 1, Supplementary Figs. S3–S6 and Supplementary Table S1). Acidic methanolysis gave compounds **5** and **6** as products, which were a mixture of isomers because of double bonds present in the monoterpene moiety. Detailed analysis of these two compounds revealed them as an acacic acid conjugated with a monoterpene (Fig. 1, Tables 1 and 2, Supplementary Figs. S7–S16).

Next, AC saponin was decomposed by HCl to yield monosaccharides and GC-MS analysis was performed (Supplementary Fig. S17). Comparison with standard sugars revealed that five sugars are present in AC saponin: glucose (Glc), rhamnose (Rha), Qui, Xyl and arabinose (Ara). In addition, the alkaline hydrolysis product was analyzed by LC-MS to obtain information about the oligosaccharides attached to the 3-*O* position of acacic acid. Several peaks corresponding to m/z of acacic acid lactone with attached oligosaccharide units were observed (Supplementary Figs. S18-25). At least seven types of oligosaccharides (Glc-Ara: m/z 782.4, Glc-Rha: m/z 796.4, Ara-Rha: m/z 766.4, Glc-Glc-Glc: m/z 974.5, Glc-Glc-Ara: m/z 944.5, Glc-Glc-Rha: m/z 958.5, or Glc-Ara-Rha: m/z 928.5 all corresponding to $[\text{M}+\text{NH}_4]^+$) were presumed to be attached at the 3-*O* position of acacic acid. Based on these analyses, the structures of the compounds in AC saponin isolated in this study are shown in Figure 2.

Inhibition of pancreatic lipase, reduction in lipid accumulation and increases in lipolysis activity

in the presence of AC saponin were evaluated as anti-obesogenic activity. AC saponin inhibited porcine pancreatic lipase with an IC_{50} of 7.93 $\mu\text{g/mL}$ (Fig. 3). Stimulation of 3T3-L1 adipocytes with 6.25 $\mu\text{g/mL}$ AC saponin reduced the lipid content by 19% when compared with the control. Stimulation of 3T3-L1 adipocytes with 3.13–12.5 $\mu\text{g/mL}$ AC saponin enhanced glycerol release by 136%–144% when compared with the control, indicating an enhancement of lipolysis activity in cells with an EC_{50} of 2.3 $\mu\text{g/mL}$. AC saponin only reduced the viability of the cells at high concentrations (Fig. 3D). Pancreatic lipase inhibition and lipolysis activity was also tested for the HCl decomposed mixture of AC saponin and apparently decreased activity was observed (13% pancreatic lipase inhibition at 20 $\mu\text{g/mL}$; no lipolysis activity between 1.56–50 $\mu\text{g/mL}$) indicating the importance of saponin structure. Co-incubation with specific inhibitors suggested that both protein kinase A (PKA) and extracellular signal-regulated kinase (ERK) contribute to the observed increase in lipolytic activity (Fig. 4).

Discussion

Saponins are glycosylated triterpenes found in plants. Saponins present in *A. concinna* possess relatively complex structures (Fig. 2), and according to previous reports, these saponins are composed of triterpene, two oligosaccharide units and two monoterpene units (inner and outer) [7–9]. The triterpene was identified in this report as acacic acid, which has three hydroxy groups (3-, 16-, 21-OH) and a carboxyl group at C17. The inner monoterpene is attached to 21-OH, with variations in structure because of the stereochemistry of C-6' and the presence or absence of a hydroxyl group at C-9. The outer monoterpene is conjugated to the inner monoterpene through a sugar unit (either quinovose or xylose), which may also conjugate to an additional sugar unit. Oligosaccharides are attached to the 3-OH and carboxyl group. The oligosaccharide attached to the carboxyl group has not been studied in detail with the only example being the $\alpha\text{-L-arabinofuranosyl}(1\rightarrow4)\text{-}[\beta\text{-D-glucopyranosyl}(1\rightarrow3)]\text{-}\alpha\text{-L-rhamnopyranosyl}(1\rightarrow2)\text{-}\beta\text{-D-glucopyranosyl}$ unit. Oligosaccharides attached to 3-OH have been relatively well studied and reported to be constructed from glucose, *N*-acetylglucosamine, fucose, rhamnose, arabinose and xylose. In comparison, the saponin isolated in this report contains acacic acid,

monoterpene and sugars, showing high similarity to reported structures [7–9]. Moreover, the observed MS ions ranging from m/z 2000–2400 (Supplementary Fig. S2) indicates similar molecular weight to the previously isolated saponin Kinmoonoside A (MW 2057) supporting the prediction that the isolated saponin is similar to the compound [7]. However, the sugar analysis revealed that the sugar units identified do not include (or are at an undetectable level) *N*-acetylglucosamine and fucose, indicating the type of oligosaccharide units differ to those previously reported in saponins from *A. concinna* (Supplementary Fig. S17). This difference in sugars may be because the *A. concinna* pods were purchased in a different region to previous studies, namely Bangladeshi or Myanmar [7–9], whereas in this study they were purchased in Thailand. The saponins from *A. concinna* consist of various monoterpene and oligosaccharide moieties, which yield complex but attractive structures.

Anti-obesogenic activities of saponins have been documented; however, the activity of AC saponin from *A. concinna* was confirmed in this study for the first time. Inhibition of lipase is a well-known anti-obesogenic activity of saponins and their IC_{50} values range from tens to hundreds of $\mu\text{g/mL}$ [10–15]. Thus, the lipase inhibitory activity of AC saponin was classified into the strongest group following comparison with the activity of other saponins (Fig. 3).

Saponins also increase lipolysis activity and reduce lipid accumulation in adipocytes. Sericoside present in *Terminalia sericea* has been shown to enhance lipolysis and reduce lipid accumulation in 3T3-L1 adipocytes at 1–10 $\mu\text{g/mL}$ [16]. Capsicoside G found in pepper seeds was found to show lipolytic activity at 200 μM and reduce lipid accumulation in 3T3-L1 adipocytes at 50–200 μM (71–285 $\mu\text{g/mL}$) [17]. Ginsenoside Rb1 contained in *Panax ginseng* was shown to increase lipolysis at 20 μM (22 $\mu\text{g/mL}$) in 3T3-L1 adipocytes but lipid accumulation was not reduced by this saponin [18]. Our data also show that AC saponin enhances lipolysis which participates in decreased accumulation of lipids in 3T3-L1 adipocytes (Fig. 3). The active concentration of AC saponin was comparable to reported values for other saponins. However, a negative feature of AC saponin is its cytotoxicity against adipocytes, which is relatively high when compared with the cytotoxicity of other saponins.

An interesting feature of the lipolytic activity of AC saponin is that it may utilize both PKA and

ERK mediated pathways. PKA is regulated mainly through β -adrenergic receptor to upregulate the lipolytic pathway in adipocytes [19]. ERK activates and enhances lipolysis through stimulation by growth hormones [20]. Although often described as independent pathways, PKA and ERK function in tandem to activate lipolysis in particular cases. For example, stimulation of β 3-adrenergic receptor results in activation of both pathways, leading to ERK and PKA regulating 20%–25% and 75%–80% of lipolysis in adipocytes, respectively [19]. Activation of lipolysis by serum amyloid A is inhibited by either a PKA or ERK inhibitor [21]. Interestingly, these inhibitors also reduced the lipolytic activity of AC saponin (Fig. 4). The lipolytic activity of serum amyloid A involves downregulating expression of perilipin combined with upregulating phosphorylation of hormone sensitive lipase. A similar mechanism may also be responsible for the lipolytic activity of AC saponin and characterizing this mechanism is a future research endeavor.

In conclusion, AC saponin was identified as the anti-obesogenic component of *A. concinna*. Inhibition of pancreatic lipase reduces lipid absorption from food sources and prevents weight gain [22]. Reducing lipids in adipocytes requires absorption of the large saponin molecules, which may not be adequately absorbed from the intestine, thereby not reaching adipose tissue in the body. However, a number of saponins have been reported to display anti-obesity effects in *in vivo* studies, indicating that sufficient absorption of saponins is possible [12, 14, 15, 18, 23, 24]. Thus, the stronger lipase inhibitory activity and comparable enhancement of lipolytic and reduced lipid accumulation activities of AC saponin when compared with other reported saponins indicate that the *A. concinna* pod or AC saponin from these pods will be a valuable resource for preventing obesity.

Materials and methods

General

Commercially available chemicals were purchased from Fujifilm Wako Pure Chemical Co. (Osaka, Japan), unless otherwise stated. Absorbances were measured using a Synergy™ MX microplate reader (BioTek Instruments, Inc., Winooski, VT, USA). NMR spectra were recorded using a Bruker AMX 500 instrument (Bruker BioSpin K.K., Bruker Instruments, Billerica, MA, USA), and residual solvents

were used as internal standards (CD₃OD: ¹H 3.31 ppm; acetone-*d*₆: ¹H 2.04 ppm, ¹³C 29.7 ppm). Mass spectra were recorded using an LCT-Premier MS (Waters Corp., Milford, MA, USA). A Waters Acquity UPLC system linked to an LCT-Premier MS was used for ultra-high-performance liquid chromatography (UPLC)-MS analysis. A Varian CP-3800 gas chromatograph equipped with a Varian 1200L MS (Agilent Technologies, Inc., Santa Clara, CA, USA) were used for GC-MS analysis. All experiments were repeated at least twice and representative data is shown in figures.

Plant material

A. concinna pods were purchased from a local store in Samut Prakan, Thailand in April 2018. The pods were stored at the Ministry of Public Health, Department of Thai Traditional and Alternative Medicine (Thailand) as voucher specimens (WR063, 10000621).

Purification of lipase inhibitory components from *A. concinna* pods

The dried pods of *A. concinna* (50.0 g) were soaked in 50% aq. methanol for 24 h. The procedure was repeated three times and the solutions were combined and dried to yield the extract (24.1 g). The extract was suspended in water and then partitioned with hexane, ethyl acetate and 1-butanol to obtain the hexane-soluble fraction (0.0208 g), ethyl acetate-soluble fraction (0.375 g), butanol-soluble fraction (5.01 g), water-soluble fraction (14.1 g) and insoluble fraction (3.75 g). The butanol-soluble fraction (2.46 g) was absorbed onto DIAION HP-20 (250 cm³, Mitsubishi Chemical Corp., Tokyo, Japan), washed with 750 mL of water and eluted with 750 mL of 50% aq. methanol followed by 750 mL of methanol to obtain the water-eluted fraction (0.90 g), 50% aq. methanol-eluted fraction (0.24 g) and methanol-eluted fraction (1.28 g). The methanol-eluted fraction (1.28 g) was subjected to reversed-phase column chromatography (3 × 18 cm) with 200 mL each of 20%, 40%, 60%, 80% aq. methanol as the eluent. The 80% aq. methanol eluted fraction (1.12 g) was subjected to Sephadex LH-20 column chromatography (3 × 16 cm) with 350 mL of 50% aq. methanol as the eluent. The eluent was determined as *A. concinna* saponin (AC saponin, 1.04 g) by NMR and MS analysis, and subsequent decomposition analysis (see Supplementary Figs. S1 and S2 for NMR and MS spectra).

Decomposition analysis of AC saponin

Alkaline hydrolysis

AC saponin (97.2 mg) was dissolved in a mixture of 0.5 M sodium hydroxide/methanol (7/1, 2 mL) and reacted for 24 h at room temperature (rt). After neutralizing with 1 M HCl, methanol was removed in vacuo and the material extracted with ethyl acetate and 1-butanol to give the ethyl acetate-soluble fraction (6.6 mg) and butanol-soluble fraction (101 mg). The butanol-soluble fraction was separated by HPLC [InertSustain C18 (20 × 250 mm, GL Science Co., Tokyo, Japan), 15%–35% aq. acetonitrile with 0.1% trifluoroacetic acid (0–30 min), 10.0 mL/min, 254 nm]. Four peaks (retention times (rT): 18, 20, 29, 35 min) were collected and analyzed by NMR. The obtained NMR spectra were compared with reported NMR data [7, 9] and the compounds were identified as monoterpenes connected with either xylose or quinovose [compound **1** (rT 18 min): (6*R*, 2*E*)-6-hydroxy-2-hydroxymethyl-6-methyl-2,7-octadienoic acid-6-O-β-D-quinovoside; compound **2** (rT 20 min): (6*S*, 2*E*)-6-hydroxy-2-hydroxymethyl-6-methyl-2,7-octadienoic acid-6-O-β-D-quinovoside; compound **3** (rT 29 min): (6*R*)-menthiafolic acid-6-O-β-D-xyloside; compound **4** (rT 35 min): (6*R*)-menthiafolic acid-6-O-β-D-quinovoside. See Supplementary Table S1 and Supplementary Figures S3–S6 for data].

Acidic methanolysis

AC saponin (94.2 mg) was dissolved in a methanol solution of 1.25 M HCl and refluxed for 1.5 h. The reaction mixture was dried and the residue suspended in water and extracted with ethyl acetate. The extract (20.1 mg) was separated by preparative TLC with chloroform-methanol (10:1) to yield two fractions (retardation factor (Rf): 0.52, 0.57). Each fraction was further separated by reversed-phase TLC with 70% aq. methanol and then by HPLC [InertSustain C18 (20 × 250 mm, GL Science Co., Tokyo, Japan), 80%–95% aq. acetonitrile with 0.1% trifluoroacetic acid (0–40 min), 10.0 mL/min, 254 nm] to yield compounds **5** (rT 22 min) and **6** (rT 21 min), which were both a conjugate of acacic acid and monoterpene, as revealed by NMR and HRMS analysis. See Tables 1 and 2 and Supplementary Figures S7–S16 for NMR data. Compound **5**: HR-ESI-MS (positive) m/z 668.4665 [M+H]⁺ (Calcd for C₄₁H₆₄O₆ 668.4652); Compound **6**: HR-ESI-MS (positive) m/z 684.4609 [M+H]⁺

(Calcd for C₄₁H₆₄O₇ 684.4601).

GC-MS analysis

GC-MS samples were prepared following a reported method with modifications [25]. Standard sugars (10 mg) were reduced with 0.25 M aq. NaBH₄ (0.5 mL) at room temperature overnight and then acetylated with 0.5 mL pyridine and 0.25 mL acetic anhydride overnight at room temperature. AC saponin was hydrolyzed initially with 1 M HCl at 100 °C for 6 h with reflux equipment to decompose oligosaccharides to monosaccharides, followed by reduction and acetylation.

Samples were analyzed under the following conditions: column, SHIMADZU SH-5 (30 m, 0.25 mm ID, 0.25 mm film); inlet temperature: 225 °C; flow: 1.0 ml/min; gas: He; oven temperature: 110 °C (0–1 min), 150 °C (20 °C/min, 1–3 min), 170 °C (1 °C/min, 3–23 min), 180 °C (10 °C/min, 23–24 min), 220 °C (20 °C/min, 24–26 min) and 320 °C (40 °C/min, >26 min).

LC-MS analysis

The butanol-soluble fraction from alkaline hydrolysis was analyzed. The sample was separated by using an InertSustain C18 column (100 mm, 2.0 mm ID, GL Science Co.) with gradient elution (5%–95% aq. methanol with 10 mM ammonium formate, 0–15 min, 0.2 mL/min) and detected by MS. The *m/z* corresponding to the prospective structure was used to identify the presence of each sugar unit.

Pancreatic lipase activity assay

Pancreatic lipase inhibitory activity was tested according to a method reported previously [6]. Briefly, glyceryl trioleate emulsion (200 µL) and the sample (100 µL in 50% DMSO aq.) were mixed and preincubated. A porcine pancreatic lipase solution (0.15 mg/mL, Sigma-Aldrich Co., 100 µL) was added and incubated for 15 min at 37 °C. To the mixture, 1 M aq. HCl (40 µL) and hexane (600 µL) were added and mixed. The hexane layer was dried and the residue was dissolved in DMSO (100 µL). Oleic acid in the solution was quantitated by LabAssay™ NEFA (Fujifilm Wako Pure Chem. Ind. Ltd.) to calculate the inhibitory activity. Cetilistat (5–10 µM) was used as a positive control.

Cell culture

3T3-L1 cells (JCRB9014) were supplied by the Japanese Collection of Research Bioresources Cell Bank. The cells were cultured in 10% fetal bovine serum (FBS)/Dulbecco's modified Eagles medium (DMEM) with antibiotics (100 units/mL penicillin G potassium salt, 50 µg/mL gentamycin sulfate, 100 µg/mL streptomycin sulfate) at 37 °C under a 10% CO₂ atmosphere.

Lipolysis activity assay

3T3-L1 cells were cultivated in 48 or 96 well plates to confluence. After reaching confluence (48–72 h; day 0), the medium was changed to differentiation medium (10% FBS/DMEM with 0.5 mM isobutylmethylxanthine, 0.25 µM dexamethazone and 10 µg/mL insulin). The medium was then replaced with insulin medium (10% FBS/DMEM with 5 µg/mL insulin) on day 2 and day 4, and replaced with 10% FBS/DMEM on day 6. On day 8, the cells were washed with PBS, and the medium was replaced with DMEM and the cells were serum starved overnight. On day 9, the sample was diluted with DMEM and incubated for 24 h. The medium was recovered, and liberated glycerol was quantitated by the Free Glycerol Reagent (Sigma-Aldrich Co.) to calculate the lipolysis enhancement activity. Viability of the remaining cells were evaluated with Cell Counting Kit-8 (Dojindo) to estimate the cytotoxicity of the samples. Isoproterenol (1 µM) was used as a positive control for the lipolysis test and Triton X-100 (0.1%) was used as a positive control for the viability test.

Lipid accumulation assay

3T3-L1 cells were cultivated and differentiated in 48 or 96 well plates until day 4. AC saponin samples diluted in insulin medium were added to the cells on day 4 and replaced with 10% FBS/DMEM containing the AC saponin sample on day 6. On day 9, the cells were stained with AdipoRed reagent (Lonza KK.) to evaluate the lipid content and calculate lipid accumulation. A *Brucea javanica* seed extract (50 µg/mL) was used as the positive control [26].

Acknowledgments

We thank Dr. Eri Fukushi and Mr. Yusuke Takata, GC-MS & NMR Laboratory, Research Faculty of Agriculture, Hokkaido University for NMR and GC-MS analysis. We thank Edanz Group (<https://en-author-services.edanz.com/ac>) for editing a draft of this manuscript.

Authors' contributions

WR and ZZ conducted the research. Design of study and structure analysis was performed by EK. All authors wrote and revised the manuscript.

Conflicts of interests

Authors declare no conflicts of interests.

References

1. Natarajan V, Natarajan S (2009) Antidermatophytic Activity of *Acacia concinna*. *Glob J Pharmacol* 3:6–7
2. Poomanee W, Chaiyana W, Intasai N, Leelapornpisid P (2015) Biological activities and characterization of the pod extracts from sompoi (*Acacia concinna* linn) grown in northern Thailand. *Int J Pharm Pharm Sci* 7:237–241
3. Khare CP (2007) *Acacia concinna* (Willd.) DC. In: *Indian Medicinal Plants*. Springer New York, New York, NY, pp 1–1
4. Wuthi-udomlert M, Vallisuta O (2011) In vitro effectiveness of *Acacia concinna* extract against dermatomycotic pathogens. *Pharmacogn J* 3:69–73. <https://doi.org/10.5530/pj.2011.19.13>
5. Liu Y, Sun M, Yao H, et al (2017) Herbal Medicine for the Treatment of Obesity: An Overview

- of Scientific Evidence from 2007 to 2017. Evidence-Based Complement Altern Med 2017:8943059. <https://doi.org/10.1155/2017/8943059>
6. Ruangaram W, Kato E (2020) Selection of Thai Medicinal Plants with Anti-Obesogenic Potential via In Vitro Methods. *Pharmaceuticals* 13:56. <https://doi.org/10.3390/ph13040056>
 7. Tezuka Y, Honda K, Banskota AH, et al (2000) Kinmoonosides A–C, Three New Cytotoxic Saponins from the Fruits of *Acacia concinna*, a Medicinal Plant Collected in Myanmar. *J Nat Prod* 63:1658–1664. <https://doi.org/10.1021/np000347f>
 8. Gafur MA, Obata T, Kiuchi F, Tsuda Y (1997) *Acacia concinna* Saponins. I. Structures of Prosapogenols, Concinnosides A-F, Isolated from the Alkaline Hydrolysate of the Highly Polar Saponin Fraction. *Chem Pharm Bull (Tokyo)* 45:620–625. <https://doi.org/10.1248/cpb.45.620>
 9. Kiuchi F, Gafur MA, Obata T, et al (1997) *Acacia concinna* Saponins. II. Structures of Monoterpenoid Glycosides in the Alkaline Hydrolysate of the Saponin Fraction. *Chem Pharm Bull (Tokyo)* 45:807–812. <https://doi.org/10.1248/cpb.45.807>
 10. Yoshizumi K, Hirano K, Ando H, et al (2006) Lupane-Type Saponins from Leaves of *Acanthopanax sessiliflorus* and Their Inhibitory Activity on Pancreatic Lipase. *J Agric Food Chem* 54:335–341. <https://doi.org/10.1021/jf052047f>
 11. Kimura H, Ogawa S, Jisaka M, et al (2006) Identification of novel saponins from edible seeds of Japanese horse chestnut (*Aesculus turbinata* Blume) after treatment with wooden ashes and their

- nutraceutical activity. *J Pharm Biomed Anal* 41:1657–1665.
- <https://doi.org/10.1016/j.jpba.2006.02.031>
12. Han L-K, Nose R, Li W, et al (2006) Reduction of fat storage in mice fed a high-fat diet long term by treatment with saponins prepared from *Kochia scoparia* fruit. *Phyther Res* 20:877–882.

<https://doi.org/10.1002/ptr.1981>

 13. Li F, Li W, Fu H, et al (2007) Pancreatic lipase-inhibiting triterpenoid saponins from fruits of *Acanthopanax senticosus*. *Chem Pharm Bull (Tokyo)* 55:1087–9.

<https://doi.org/10.1248/cpb.55.1087>

 14. Karu N, Reifen R, Kerem Z (2007) Weight Gain Reduction in Mice Fed *Panax ginseng* Saponin, a Pancreatic Lipase Inhibitor. *J Agric Food Chem* 55:2824–2828.

<https://doi.org/10.1021/jf0628025>

 15. Bai-Shen Sun (2012) Anti-obesity effects of mogrosides extracted from the fruits of *Siraitia grosvenorii* (Cucurbitaceae). *African J Pharm Pharmacol* 6:1492–1501.

<https://doi.org/10.5897/AJPP12.310>

 16. Mochizuki M, Hasegawa N (2006) Acceleration of lipid degradation by sericoside of *Terminalia sericea* roots in Fully differentiated 3T3-L1 cells. *Phyther Res* 20:1020–1021.

<https://doi.org/10.1002/ptr.1999>

 17. Sung J, Yu W, Lee J (2018) Effects of capsicoside G isolated from pepper seeds on lipolysis and

- fatty acid oxidation-related gene expression in 3T3-L1 adipocytes. *J Food Biochem* 42:e12640.
<https://doi.org/10.1111/jfbc.12640>
18. Lim S, Park J, Um J-Y (2019) Ginsenoside Rb1 Induces Beta 3 Adrenergic Receptor–Dependent Lipolysis and Thermogenesis in 3T3-L1 Adipocytes and db/db Mice. *Front Pharmacol* 10:1–13.
<https://doi.org/10.3389/fphar.2019.01154>
 19. Robidoux J, Kumar N, Daniel KW, et al (2006) Maximal β 3-Adrenergic Regulation of Lipolysis Involves Src and Epidermal Growth Factor Receptor-dependent ERK1/2 Activation. *J Biol Chem* 281:37794–37802. <https://doi.org/10.1074/jbc.M605572200>
 20. Bergan-Roller HE, Sheridan MA (2018) The growth hormone signaling system: Insights into coordinating the anabolic and catabolic actions of growth hormone. *Gen Comp Endocrinol* 258:119–133. <https://doi.org/10.1016/j.ygcen.2017.07.028>
 21. Liu LR, Lin SP, Chen CC, et al (2011) Serum Amyloid A Induces Lipolysis by Downregulating Perilipin Through ERK1/2 and PKA Signaling Pathways. *Obesity* 19:2301–2309.
<https://doi.org/10.1038/oby.2011.176>
 22. Pilitsi E, Farr OM, Polyzos SA, et al (2019) Pharmacotherapy of obesity: Available medications and drugs under investigation. *Metabolism* 92:170–192.
<https://doi.org/10.1016/j.metabol.2018.10.010>
 23. Poudel B, Lim S-W, Ki H-H, et al (2014) Dioscin inhibits adipogenesis through the

- AMPK/MAPK pathway in 3T3-L1 cells and modulates fat accumulation in obese mice. *Int J Mol Med* 34:1401–8. <https://doi.org/10.3892/ijmm.2014.1921>
24. Liu H, Wang J, Liu M, et al (2018) Antiobesity Effects of Ginsenoside Rg1 on 3T3-L1 Preadipocytes and High Fat Diet-Induced Obese Mice Mediated by AMPK. *Nutrients* 10:830. <https://doi.org/10.3390/nu10070830>
25. Sawardeker JS, Sloneker JH, Jeanes A (1965) Quantitative Determination of Monosaccharides as Their Alditol Acetates by Gas Liquid Chromatography. *Anal Chem* 37:1602–1604. <https://doi.org/10.1021/ac60231a048>
26. Lahrita L, Moriai K, Iwata R, et al (2019) Quassinoids in *Brucea javanica* are potent stimulators of lipolysis in adipocytes. *Fitoterapia* 137:104250. <https://doi.org/10.1016/j.fitote.2019.104250>

Tables

Table 1 ¹H-NMR spectral data (500 MHz, acetone-*d*₆) of compounds **5** and **6** (major isomer)

Compound 5				Compound 6				
No	ppm	Int.	Mult.	<i>J</i> (Hz)	ppm	Int.	Mult.	<i>J</i> (Hz)
Triterpene part								
1	1.02	1H	m		1.03	1H	m	
	1.61	1H	m		1.61	1H	m	
2	1.59	1H	m		1.56	1H	m	
	1.59	1H	m		1.61	1H	m	
3	3.15	1H	m		3.15	1H	m	
5	0.76	1H	m		0.79	1H	m	
6	1.39	1H	m		1.39	1H	m	
	1.56	1H	m		1.56	1H	m	
7	1.32	1H	m		1.32	1H	m	
	1.55	1H	m		1.55	1H	m	
9	1.68	1H	m		1.68	1H	m	
11	1.91	2H	m		1.91	2H	m	
12	5.36	1H	m		5.35	1H	m	
15	1.43	1H	m		1.43	1H	m	
	1.91	1H	m		1.91	1H	m	
16	4.57	1H	br s		4.58	1H	br s	
18	3.15	1H	m		3.15	1H	m	
19	1.2	1H	dd	4.5, 13.7	1.19	1H	dd	4.6, 13.7
	2.56	1H	t	13.7	2.56	1H	t	13.7
21	5.53	1H	dd	5.6, 11.3	5.56	1H	dd	5.4, 11.0
22	1.75	1H	m		1.76	1H	dd	11.0, 13.3
	2.14	1H	dd	5.6, 13.2	2.14	1H	dd	5.4, 13.3
23	0.78	3H	s		0.78	3H	s	
24	0.99	3H	s		0.99	3H	s	
25	0.95	3H	s		0.95	3H	s	
26	0.78	3H	s		0.78	3H	s	
27	1.45	3H	s		1.45	3H	s	
29	0.88	3H	s		0.89	3H	s	
30	1.09	3H	s		1.09	3H	s	
Monoterpene part								
3'	6.73	1H	t	7	6.82	1H	t	7.5
4'	2.33	2H	dt	7.0, 8.0	2.48	2H	dt	7.5, 7.4
5'	2.18	2H	t	8	2.19	2H	t	7.4
7'	5.36	1H	m		5.35	1H	m	
8'	3.90	2H	d	6	3.90	2H	d	6.3
9'	1.83	3H	s		4.31	2H	s	
10'	1.68	3H	s		1.68	3H	s	
11'	3.22	3H	s		3.22	3H	s	

Table 2 ^{13}C -NMR spectral data (500 MHz, acetone- d_6) of compounds **5** and **6** (major isomer)

Triterpene part			Monoterpene part		
No.	Compound 5	Compound 6	No.	Compound 5	Compound 6
1	39.19	39.36	1'	ND	167.25
2	27.88	27.94	2'	ND	133.78
3	78.32	78.38	3'	143.56	144.47
4	39.30	39.25	4'	27.25	27.04
5	56.11	56.17	5'	38.63	38.98
6	18.96	19.01	6'	ND	138.70
7	33.72	33.78	7'	123.07	123.20
8	40.08	40.14	8'	69.08	69.13
9	47.40	47.46	9'	12.44	56.55
10	37.65	37.71	10'	16.13	16.19
11	23.93	23.99	11'	57.36	57.46
12	123.44	123.50			
13	143.50	143.59			
14	41.95	42.00			
15	35.42	35.49			
16	74.03	74.10			
17	51.02	51.07			
18	40.76	40.80			
19	47.95	48.04			
20	35.42	35.59			
21	77.40	77.24			
22	37.13	37.10			
23	16.13	16.19			
24	28.52	28.58			
25	15.68	15.74			
26	17.42	17.48			
27	27.07	27.14			
28	176.70	177.06			
29	29.07	29.29			
30	19.11	19.15			

ND: not detected.

Figure captions

Fig. 1 Structures of isolated compounds from decomposed AC saponin

Fig. 2 Predicted structure of AC saponin. Sugar unit 1: bond type or connection between each sugar was not determined. Xylose may replace rhamnose. Sugar unit 2: example from the reference. ara: arabinose; glc: glucose; qui: quinovose; rha: rhamnose; xyl: xylose

Fig. 3 Anti-obesogenic activity of AC saponin. Top left: pancreatic lipase inhibitory activity. Top right: lipid accumulation reduction activity, PC: *B. javanica* seed extract. Bottom left: lipolysis enhancement activity, PC: isoproterenol. Bottom right: cell viability, TX: Triton-X100. Ctrl: control. Data are expressed as means \pm SEM. * $p < 0.05$, ** $p < 0.01$ (t -test to control)

Fig. 4 Effect of inhibitors on lipolysis enhancement activity of AC saponin. AC saponin: 6.3 $\mu\text{g/mL}$; IPT: isoproterenol (1 μM); β -estradiol: 20 μM ; H-89: 20 μM ; PD98059: 50 μM . Data are expressed as means \pm SEM. Different letters indicate significant difference ($p < 0.05$, Tukey's test)

Figures




Fig. 1 Structures of isolated compounds from decomposed AC saponin




Fig. 2 Predicted structure of AC saponin. Sugar unit 1: bond type or connection between each sugar was not determined. Xylose may replace rhamnose. Sugar unit 2: example from the reference. ara: arabinose; glc: glucose; qui: quinovose; rha: rhamnose; xyl: xylose




Fig. 3 Anti-obesogenic activity of AC saponin. Top left: pancreatic lipase inhibitory activity. Top right: lipid accumulation reduction activity, PC: *B. javanica* seed extract. Bottom left: lipolysis enhancement activity, PC: isoproterenol. Bottom right: cell viability, TX: Triton-X100. Ctrl: control. Data are expressed as means \pm SEM. * $p < 0.05$, ** $p < 0.01$ (t -test to control)




Fig. 4 Effect of inhibitors on lipolysis enhancement activity of AC saponin. AC saponin: 6.3 $\mu\text{g/mL}$; IPT: isoproterenol (1 μM); β -estradiol: 20 μM ; H-89: 20 μM ; PD98059: 50 μM . Data are expressed as means \pm SEM. Different letters indicate significant difference ($p < 0.05$, Tukey's test)