



HOKKAIDO UNIVERSITY

Title	Quasilinear elliptic equations with sub-natural growth terms in bounded domains
Author(s)	Hara, Takanobu
Citation	Nodea-nonlinear Differential Equations and Applications, 28(6), 62 https://doi.org/10.1007/s00030-021-00724-5
Issue Date	2021-11
Doc URL	https://hdl.handle.net/2115/87051
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature' s AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: http://dx.doi.org/10.1007/s00030-021-00724-5
Type	journal article
File Information	NoDea-Nonlinear Differ. Equ. Appl. 28-6_62.pdf



QUASILINEAR ELLIPTIC EQUATIONS WITH SUB-NATURAL GROWTH TERMS IN BOUNDED DOMAINS

TAKANOBU HARA

ABSTRACT. We consider the existence of positive solutions to weighted quasilinear elliptic differential equations of the type

$$\begin{cases} -\Delta_{p,w}u = \sigma u^q & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

in the sub-natural growth case $0 < q < p - 1$, where Ω is a bounded domain in \mathbb{R}^n , $\Delta_{p,w}$ is a weighted p -Laplacian, and σ is a nonnegative (locally finite) Radon measure on Ω . We give criteria for the existence problem. For the proof, we investigate various properties of p -superharmonic functions, especially the solvability of Dirichlet problems with infinite measure data.

1. INTRODUCTION

Let Ω be a bounded domain in \mathbb{R}^n and let $1 < p < \infty$. We consider the existence of positive solutions to quasilinear elliptic equations of the type

$$(1.1) \quad \begin{cases} -\Delta_{p,w}u = \sigma u^q & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

in the sub-natural growth case $0 < q < p - 1$, where $\Delta_{p,w}$ is a weighted (p, w) -Laplacian, w is a p -admissible weight on \mathbb{R}^n (see Section 2 below) and σ is a nonnegative (locally finite) Radon measure on Ω .

For the standard theory of sublinear equations, we refer to [32, 18, 11, 10] and the references therein. In the classical existence results of weak solutions, the boundedness of coefficients was assumed. Boccardo and Orsina [8] removed this assumption and pointed out that if the integrability of the coefficient σ is low, the corresponding solution u does not necessarily belong to the class of weak solutions. Therefore, we interpret this equation in the sense of p -superharmonic functions, or locally renormalized solutions. For details on such generalized solutions, especially on the relation between the two concepts, see [4, 28, 48] and references therein. Hereafter, we use the framework for p -superharmonic functions.

The measure-valued coefficient equation (1.1) is relevant to the following L^p - L^{1+q} trace inequality:

$$(1.2) \quad \|f\|_{L^{1+q}(\Omega; \sigma)} \leq C_T \|\nabla f\|_{L^p(\Omega; w)}, \quad \forall f \in C_c^\infty(\Omega).$$

Maz'ya and Netrusov [35] gave a capacity condition that characterizes (1.2). Cascante, Ortega and Verbitsky [15, 45] studied non-capacity characterizations

Date: August 13, 2021.

1991 Mathematics Subject Classification. 35J92, 35J20, 42B37.

Key words and phrases. Quasilinear elliptic equation, p -Laplacian, Wolff potential, Measure data, Trace inequality.

for inequalities of the type (1.2). For example, if $\Omega = \mathbb{R}^n$ and $w = 1$, then the best constant C_T in (1.2) satisfies

$$\frac{1}{c} C_T^{\frac{(1+q)p}{p-1-q}} \leq \int_{\mathbb{R}^n} (\mathbf{W}_{1,p}\sigma)^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma \leq c C_T^{\frac{(1+q)p}{p-1-q}},$$

where $c = c(n, p, q)$ and $\mathbf{W}_{1,p}\sigma$ is the *Wolff potential* of σ (see [33, 26] and Theorem 2.2 below).

Recently, Verbitsky and his colleagues studied the problem of existence of solutions to elliptic equations related to (1.1) and presented some criteria (see [12, 13, 14, 21, 22, 38, 39, 41, 42, 46, 47]). In their study, they treated the cases of $\Omega = \mathbb{R}^n$, or $p = 2$. For $\Omega = \mathbb{R}^n$, Wolff potentials are suitable potentials for the problem. Every p -superharmonic function is known to be locally estimated by Wolff potentials (see [29, 30]), and these estimates are some of the key pieces of proof. In contrast, we can directly use Green potentials for $p = 2$ or, more generally, for linear equations. However, Wolff potentials are not sufficient for estimating the boundary behavior of solutions. Furthermore, Radon measures satisfying (1.2) may not have compact support in Ω and are not even finite in general (see Section 7). Consequently, the complete criteria for the existence of solutions to quasilinear equations in bounded domains have not yet been obtained.

The purpose of this paper is to extend Verbitsky's theory to (weighted) quasilinear equations in bounded domains. Our basic idea is to replace Wolff or Green potentials with the minimal positive p -superharmonic solution u to $-\Delta_{p,w}u = \sigma$ (see Definition 3.7 for the precise meaning). The function $\mathcal{W}_{p,w}\sigma = u$ has no explicit integral representation; however, some required estimates can be obtained by directly using the properties of weak solutions. To realize this approach, we investigate various properties of p -superharmonic functions, especially the solvability of Dirichlet problems in the case of infinite measure data. Note that this existence problem has been stated as an open problem in [3, Problem 2].

Let $\mathcal{M}_0^+(\Omega)$ be the set of all nonnegative Radon measures on Ω that are absolutely continuous with respect to the (p, w) -capacity. Note that σ must belong to $\mathcal{M}_0^+(\Omega)$ if (1.2) holds. Our main result is as follows.

Theorem 1.1. *Let Ω be a bounded domain in \mathbb{R}^n . Let $1 < p < \infty$ and $0 < q < p-1$. Suppose that $\sigma \in \mathcal{M}_0^+(\Omega) \setminus \{0\}$. Fix $0 < \gamma < \infty$. Then the following statements are equivalent:*

- (1) *There exists a nontrivial nonnegative (p, w) -superharmonic supersolution v to $-\Delta_{p,w}v = \sigma v^q$ in Ω satisfying $\|v\|_{L^{\gamma+q}(\sigma)} \leq C_1 < \infty$.*
- (2) *The measure σ satisfies*

$$(1.3) \quad \left(\int_{\Omega} (\mathcal{W}_{p,w}\sigma)^{\frac{(\gamma+q)(p-1)}{p-1-q}} d\sigma \right)^{\frac{1}{\gamma+q}} \leq C_2 < \infty.$$

- (3) *The following weighted norm inequality holds:*

$$(1.4) \quad \|\mathcal{W}_{p,w}(|f|\sigma)\|_{L^{\gamma+q}(\sigma)} \leq C_3 \|f\|_{L^{\frac{\gamma+q}{q}}(\sigma)}, \quad \forall f \in L^{\frac{\gamma+q}{q}}(\sigma).$$

Moreover, if C_i ($i = 1, 2, 3$) are the best constants in the above statements, then

$$C_1 \leq C_3^{\frac{p-1}{p-1-q}} \leq c_E^{\frac{1}{p-1-q}} C_2 \leq \frac{c_E^{\frac{1}{p-1-q}}}{c_V} C_1,$$

where

$$(1.5) \quad c_E := \left(\frac{p-1+\gamma}{p} \right)^p \frac{1}{\gamma}, \quad c_V := \left(\frac{p-1-q}{p-1} \right)^{\frac{p-1}{p-1-q}}.$$

In addition, if one of the above statements holds, then there exists a minimal positive (p, w) -superharmonic solution u to $-\Delta_{p,w}u = \sigma u^q$ in Ω such that $\|u\|_{L^{\gamma+q}(\sigma)} \leq C_1$.

The boundary behavior of u is not discussed in Theorem 1.1; however, under appropriate assumptions on σ , we can prove that u vanishes on $\partial\Omega$ (see Proposition 4.2 and Corollary 4.4). One such condition is as follows.

Theorem 1.2. *Let Ω be a bounded domain in \mathbb{R}^n . Let $1 < p < \infty$ and $0 < q < p-1$. Suppose that $\sigma \in \mathcal{M}_0^+(\Omega) \setminus \{0\}$. Then there exists a unique positive weak solution $u \in H_0^{1,p}(\Omega; w)$ to (1.1) if and only if*

$$(1.6) \quad \int_{\Omega} (\mathcal{W}_{p,w}\sigma)^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma < \infty.$$

Moreover,

$$c_V^{1+q} \int_{\Omega} (\mathcal{W}_{p,w}\sigma)^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma \leq \|\nabla u\|_{L^p(w)}^p \leq \int_{\Omega} (\mathcal{W}_{p,w}\sigma)^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma.$$

We also give the Cascante-Ortega-Verbitsky type of theorem below.

Theorem 1.3. *Let Ω be a bounded domain in \mathbb{R}^n . Let $1 < p < \infty$ and let $-1 < q < p-1$. Suppose that C_T is the best constant of (1.2). Then*

$$C_T^{\frac{(1+q)p}{p-1-q}} \leq \int_{\Omega} (\mathcal{W}_{p,w}\sigma)^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma \leq \left(\frac{1}{1+q} \right)^{\frac{1+q}{p-1-q}} \frac{1}{c_V^{1+q}} C_T^{\frac{(1+q)p}{p-1-q}}.$$

In particular, if (1.2) holds with some $-1 < q < p-1$, then the equation $-\Delta_{p,w}u = \sigma$ in Ω has a minimal positive (p, w) -superharmonic solution.

Remark 1.4. The constants in Theorems 1.1-1.3 do not depend on n or the data of w . Due to the qualitative arguments in the proof, w must be p -admissible; however, its quantitative properties, especially the Sobolev-type inequalities are not used.

In particular, if (1.2) holds with $q > 0$, then there exists a unique positive weak solution u to (1.1) such that

$$\frac{1}{c(p, q)} C_T^{\frac{1+q}{p-1-q}} \leq \|\nabla u\|_{L^p(w)} \leq c(p, q) C_T^{\frac{1+q}{p-1-q}}.$$

Examples of concrete sufficient conditions for (1.3) will be discussed at the end of the paper. One is a Lorentz scale refinement of [8, Theorem 5.5], and the other is quasilinear ordinary differential equations with nonintegrable Hardy-type coefficients.

Organization of the paper. Section 2 presents various facts in nonlinear potential theory and introduces classes of smooth measures. Section 3 discusses minimal p -superharmonic solutions to Dirichlet problems. Section 4 defines the generalized energy of p -superharmonic functions and investigates its properties including Theorem 1.3. Section 5 provides the proof of Theorem 1.1 by using results in Sections 3 and 4. Sections 6 and 7 discuss two applications of Theorems 1.1-1.3.

Acknowledgments. The author would like to thank Professor Verbitsky for suggesting references to added to an earlier version of the manuscript. This work was supported by JSPS KAKENHI Grant Number JP18J00965 and JP17H01092. I would also like to thank my family for their support.

Notation. We use the following notation. Let Ω be a domain (connected open subset) in \mathbb{R}^n .

- $\mathbf{1}_E(x) :=$ the indicator function of a set E .
- $C_c^\infty(\Omega) :=$ the set of all infinitely differentiable functions with compact support in Ω .
- $\mathcal{M}^+(\Omega) :=$ the set of all nonnegative Radon measures on Ω .
- $L^p(\mu) :=$ the L^p space with respect to $\mu \in \mathcal{M}^+(\Omega)$.

For a ball $B = B(x, R)$ and $\lambda > 0$, $\lambda B := B(x, \lambda R)$. For measures μ and ν , we denote $\nu \leq \mu$ if $\mu - \nu$ is a nonnegative measure. For a sequence of extended real valued functions $\{f_j\}_{j=1}^\infty$, we denote $f_j \uparrow f$ if $f_{j+1} \geq f_j$ for all $j \geq 1$ and $\lim_{j \rightarrow \infty} f_j = f$. Moreover, c and C denote various constants with and without indices.

2. PRELIMINARIES

We first recall the basic properties of p -admissible weights from [5, Chapter A.2], [27, Chapter 20] and references therein. Throughout the paper, $1 < p < \infty$ is a fixed constant. A Lebesgue measurable function w on \mathbb{R}^n to be said as *weight* on \mathbb{R}^n if $w \in L_{\text{loc}}^1(\mathbb{R}^n; dx)$ and $w(x) > 0$ dx -a.e. We write $w(E) = \int_E w dx$ for a Lebesgue measurable set $E \subset \mathbb{R}^n$. We always assume that w is *p-admissible*, that is, positive constants C_D , C_P and $\lambda \geq 1$ exist, such that

$$w(2B) \leq C_D w(B)$$

and

$$\int_B |f - f_B| dw \leq C_P \text{diam}(B) \left(\int_{\lambda B} |\nabla f|^p dw \right)^{\frac{1}{p}}, \quad \forall f \in C_c^\infty(\mathbb{R}^n),$$

where B is an arbitrary ball in \mathbb{R}^n , $f_B = w(B)^{-1} \int_B f dw$ and $f_B = \int_B f dw$. One of the important properties of p -admissible weights is the Sobolev inequality ([23, 40]). In particular, the following form of the Poincaré inequality holds:

$$\int_B |f|^p dw \leq C \text{diam}(B)^p \int_B |\nabla f|^p dw, \quad \forall f \in C_c^\infty(B),$$

where C is a constant depending only on p , C_D , C_P and λ .

Next, we recall basics of nonlinear potential theory from [27, Chapters 1-10 and 21]. Let Ω be a bounded domain in \mathbb{R}^n . The weighted Sobolev space $H^{1,p}(\Omega; w)$ is the closure of $C^\infty(\Omega)$ with respect to the norm

$$\|u\|_{H^{1,p}(\Omega; w)} := \left(\int_\Omega |u|^p + |\nabla u|^p dw \right)^{\frac{1}{p}},$$

where ∇u is the gradient of u . The corresponding local space $H_{\text{loc}}^{1,p}(\Omega; w)$ is defined in the usual manner. We denote the closure of $C_c^\infty(\Omega)$ in $H^{1,p}(\Omega; w)$ by $H_0^{1,p}(\Omega; w)$. Since Ω is bounded, we can take $\|\nabla \cdot\|_{L^p(\Omega; w)}$ as the norm of $H_0^{1,p}(\Omega; w)$ by the Poincaré inequality.

For $u \in H_{\text{loc}}^{1,p}(\Omega; w)$, we define the weighted p -Laplace operator $\Delta_{p,w}$ by

$$\langle -\Delta_{p,w}u, \varphi \rangle = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \varphi \, dw, \quad \forall \varphi \in C_c^\infty(\Omega).$$

A function $u \in H_{\text{loc}}^{1,p}(\Omega; w)$ is called a *supersolution* to

$$(2.1) \quad -\Delta_{p,w}u = 0 \quad \text{in } \Omega$$

if $\langle -\Delta_{p,w}u, \varphi \rangle \geq 0$ for all nonnegative $\varphi \in C_c^\infty(\Omega)$. If μ is an element of the dual of $H_0^{1,p}(\Omega; w)$, then the Dirichlet problem

$$(2.2) \quad \begin{cases} \langle -\Delta_{p,w}u, \varphi \rangle = \langle \mu, \varphi \rangle & \forall \varphi \in H_0^{1,p}(\Omega; w) \\ u \in H_0^{1,p}(\Omega; w) \end{cases}$$

has a unique weak solution u .

Let $\Omega \subset \mathbb{R}^n$ be open, and let $K \subset \Omega$ be compact. The *variational (p, w) -capacity* $\text{cap}_{p,w}(K, \Omega)$ of the condenser (K, Ω) is defined by

$$\text{cap}_{p,w}(K, \Omega) := \inf \left\{ \|\nabla u\|_{L^p(\Omega; w)}^p : u \geq 1 \text{ on } K, u \in C_c^\infty(\Omega) \right\}.$$

Moreover, for $E \subset \Omega$, we define

$$\text{cap}_{p,w}(E, \Omega) := \inf_{\substack{E \subset U \subset \Omega \\ U: \text{open}}} \sup_{K \subset U: \text{compact}} \text{cap}_{p,w}(K, \Omega).$$

Since $\Omega \subset \mathbb{R}^n$ is bounded, $\text{cap}_{p,w}(E, \Omega) = 0$ if and only if $C_p(E) = 0$, where $C_p(\cdot)$ is the *Sobolev capacity* of E (see [5, Lemma 6.15]). We say that a property holds *quasieverywhere* (q.e.) if it holds except on a set of (p, w) -capacity zero. An extended real valued function u on Ω is called *quasicontinuous* if for every $\epsilon > 0$ there exists an open set G such that $C_p(G) < \epsilon$ and $u|_{\Omega \setminus G}$ is continuous. Every $u \in H_{\text{loc}}^{1,p}(\Omega; w)$ has a quasicontinuous representative \tilde{u} such that $u = \tilde{u}$ a.e.

A function $u: \Omega \rightarrow (-\infty, \infty]$ is called *(p, w) -superharmonic* if u is lower semicontinuous in Ω , is not identically infinite, and satisfies the comparison principle on each subdomain $D \Subset \Omega$: if $h \in C(\overline{D})$ is a continuous weak solution to $-\Delta_{p,w}u = 0$ in D , and if $u \geq h$ on ∂D , then $u \geq h$ in D . If u is a bounded (p, w) -superharmonic function, then u belongs to $H_{\text{loc}}^{1,p}(\Omega; w)$ and it is a supersolution to (2.1). Conversely, if u is a supersolution to (2.1), then its *lsc-regularization*

$$u^*(x) = \lim_{r \rightarrow 0} \text{ess inf}_{B(x,r)} u$$

is (p, w) -superharmonic in Ω . If u and v are (p, w) -superharmonic in Ω and $u(x) \leq v(x)$ for a.e. $x \in \Omega$, then $u(x) \leq v(x)$ for all $x \in \Omega$. Every (p, w) -superharmonic function is known to be quasicontinuous. In particular, the set $\{u = \infty\}$ has zero (p, w) -capacity whenever u is (p, w) -superharmonic.

Assume that u is a (p, w) -superharmonic function in Ω . Its truncation $u_k = \min\{u, k\}$ will then become a supersolution to (2.1) for all $k > 0$. As in [27, Theorem 7.46], the *very weak gradient* Du of u belongs to $L_{\text{loc}}^{Q(p-1)}(\Omega; w)$, where $Du = \lim_{k \rightarrow \infty} \nabla u_k$ and $Q > 1$ is a constant depending only on p, C_D, C_P and λ . Moreover, there exists a Radon measure $\mu[u]$ such that

$$\int_{\Omega} |Du|^{p-2} Du \cdot \nabla \varphi \, dw = \int_{\Omega} \varphi \, d\mu[u], \quad \forall \varphi \in C_c^\infty(\Omega).$$

The measure $\mu[u]$ is called the *Riesz measure* of u . By definition, if $u \in H_{\text{loc}}^{1,p}(\Omega; w)$, then $\mu[u] = -\Delta_{p,w}u$ in the sense of distribution.

As in Section 1, we denote by $\mathcal{M}_0^+(\Omega)$ the set of all Radon measures μ that are absolutely continuous with respect to the (p, w) -capacity, i.e., $\mu(E) = 0$ whenever E has zero (p, w) -capacity. If $u \in H_{\text{loc}}^{1,p}(\Omega; w)$ is (p, w) -superharmonic in Ω , then the Riesz measure of u belongs to $\mathcal{M}_0^+(\Omega)$. It is known that if $\mu \in \mathcal{M}_0^+(\Omega)$ is finite, then the integral $\int_{\Omega} f d\mu$ is well-defined for any quasicontinuous function $f \in H_0^{1,p}(\Omega; w)$.

If $u \in H^{1,p}(\Omega; w)$ is a supersolution to (2.1), then the Riesz measure of u^* satisfies

$$(2.3) \quad \left| \int_{\Omega} \varphi d\mu \right| \leq C \|\varphi\|_{H^{1,p}(\Omega; w)}, \quad \forall \varphi \in C_c^\infty(\Omega).$$

If μ is also finite, then we can replace $C_c^\infty(\Omega)$ with $H_0^{1,p}(\Omega; w)$ up to taking a quasicontinuous representative, i.e., the dual action of μ has the integral representative. Conversely, if a finite measure μ satisfies (2.3), then there exists a unique weak solution $u \in H_0^{1,p}(\Omega; w)$ to

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \varphi dw = \int_{\Omega} \tilde{\varphi} d\mu \quad \forall \varphi \in H_0^{1,p}(\Omega; w).$$

The following weak continuity result was given by Trudinger and Wang [44].

Theorem 2.1 ([44, Theorem 3.1]). *Suppose that $\{u_k\}_{k=1}^\infty$ is a sequence of nonnegative (p, w) -superharmonic functions in Ω . Assume that $u_k \rightarrow u$ a.e. in Ω and that u is (p, w) -superharmonic in Ω . Let $\mu[u_k]$ and $\mu[u]$ be the Riesz measures of u_k and u , respectively. Then $\mu[u_k]$ converges to $\mu[u]$ weakly, that is*

$$\int_{\Omega} \varphi d\mu[u_k] \rightarrow \int_{\Omega} \varphi d\mu[u], \quad \forall \varphi \in C_c^\infty(\Omega).$$

The following Harnack-type convergence theorem follows from combining Theorem 2.1 and [27, Lemma 7.3]: If $\{u_k\}_{k=1}^\infty$ is a nondecreasing sequence of (p, w) -superharmonic functions in Ω and if $u := \lim_{k \rightarrow \infty} u_k \not\equiv \infty$, then u is (p, w) -superharmonic in Ω and $\mu[u_k]$ converges to $\mu[u]$ weakly.

The following form of Wolff potential estimate was first established by Kilpeläinen and Malý [29, 30]. The extension to weighted equations is due to Mikkonen [36]. See also [44, 24] for other proofs.

Theorem 2.2 ([36, Theorem 3.1]). *Suppose that u is a nonnegative (p, w) -superharmonic function in $B(x, 2R)$. Let μ be the Riesz measure of u . Then*

$$\frac{1}{C} \mathbf{W}_{1,p,w}^R \mu(x) \leq u(x) \leq C \left(\inf_{B(x,R)} u + \mathbf{W}_{1,p,w}^{2R} \mu(x) \right),$$

where $C \geq 1$ is a constant depending only on p , C_D , C_P and λ , and $\mathbf{W}_{1,p,w}^R \mu$ is the truncated Wolff potential of μ , which is defined by

$$\mathbf{W}_{1,p,w}^R \mu(x) := \int_0^R \left(r^p \frac{\mu(B(x,r))}{w(B(x,r))} \right)^{\frac{1}{p-1}} \frac{dr}{r}.$$

3. MINIMAL p -SUPERHARMONIC SOLUTIONS

The following comparison principle improves [12, Lemma 5.2]. We do not assume the finiteness of the Riesz measures here.

Theorem 3.1. *Let Ω be a bounded domain in \mathbb{R}^n . Let u and v be nonnegative (p, w) -superharmonic functions in Ω with the Riesz measures μ and ν , respectively. Assume also that $\mu \leq \nu$ and $u \in H_0^{1,p}(\Omega; w)$. Then $u(x) \leq v(x)$ for all $x \in \Omega$.*

Proof. For each $k \in \mathbb{N}$, set $v_k = \min\{v, k\}$. Let ν_k be the Riesz measure of v_k . By the chain rule of Sobolev functions,

$$\Psi_\epsilon^k(v) := \frac{(k-v)_+}{(k-v)_+ + \epsilon} \in H_{\text{loc}}^{1,p}(\Omega; w) \cap L^\infty(\Omega), \quad \forall \epsilon > 0.$$

Fix $\varphi \in C_c^\infty(\Omega)$ such that $\varphi \geq 0$. For any $l \geq k$,

$$\int_\Omega |\nabla v_l|^{p-2} \nabla v_l \cdot \nabla (\varphi \Psi_\epsilon^k(v)) \, dw \leq \int_\Omega |\nabla v_l|^{p-2} \nabla v_l \cdot \nabla \varphi \Psi_\epsilon^k(v) \, dw.$$

Note that $\Psi_\epsilon^k(v) = 0$ on $\{v \geq k\}$ and $Dv = \nabla v_l = \nabla v_k$ on $\{v < k\}$. Thus, by [28, Theorem 3.15],

$$\begin{aligned} \int_\Omega \varphi \Psi_\epsilon^k(v) \, d\nu &= \int_\Omega |Dv|^{p-2} Dv \cdot \nabla (\varphi \Psi_\epsilon^k(v)) \, dw \\ &\leq \int_\Omega |\nabla v_k|^{p-2} \nabla v_k \cdot \nabla \varphi \Psi_\epsilon^k(v) \, dw. \end{aligned}$$

Since $\lim_{\epsilon \rightarrow 0} \Psi_\epsilon^k(v)(x) = \mathbf{1}_{\{v < k\}}(x)$ for all $x \in \Omega$, it follows from the dominated convergence theorem that $\mathbf{1}_{\{v < k\}} \nu \leq \nu_k$. Let $\mu_k = \mathbf{1}_{\{v < k\}} \mu$, and let $\{\Omega_k\}_{k=1}^\infty$ be a sequence of open sets such that $\bigcup_{k=1}^\infty \Omega_k = \Omega$ and $\Omega_k \Subset \Omega_{k+1}$ for all $k \geq 1$. Since $\mu_k \leq \mu$, there exists a weak solution $u_k \in H_0^{1,p}(\Omega_k; w)$ to $-\Delta_{p,w} u_k = \mu_k$ in Ω_k . Let us denote by u_k the zero extension of the lsc-regularization of u_k again. By the comparison principle for weak solutions, $u_k(x) \leq v_k(x) \leq v(x)$ for a.e. $x \in \Omega_k$. Accordingly, $u_k(x) \leq v(x)$ for all $x \in \Omega_k$ since u and v are (p, w) -superharmonic in Ω_k . Similarly, $u_k \leq u_{k+1} \leq u$ in Ω_k . Let $u' = \lim_{k \rightarrow \infty} u_k$, and let μ' be the Riesz measure of u' . Testing the equation of u_k with u_k , we have

$$\int_\Omega |\nabla u_k|^p \, dw = \int_{\Omega_k} u_k \, d\mu_k \leq \int_{\Omega_k} u_k \, d\mu \leq \|\nabla u\|_{L^p(\Omega; w)}^{p-1} \|\nabla u_k\|_{L^p(\Omega; w)}.$$

Therefore $u' \in H_0^{1,p}(\Omega; w)$. By Theorem 2.1, $\{\mu_k\}_{k=1}^\infty$ converges to μ' weakly. Meanwhile, since $\mu(\{v = \infty\}) = 0$, $\{\mu_k\}_{k=1}^\infty$ converges to μ weakly. Thus $\mu = \mu'$. By the uniqueness of weak solutions, $u = u' \leq v$ in Ω . \square

Following [19, Chapter 2], we introduce classes of smooth measures.

Definition 3.2. Let S_0 be the set of all Radon measures satisfying (2.3). For $\mu \in S_0$, we denote the lsc-regularization of the solution to (2.2) by $\mathcal{W}_{p,w}^0 \mu$. Furthermore, we define a subset S_{00} of S_0 as

$$S_{00} := \left\{ \mu \in S_0 : \sup_\Omega \mathcal{W}_{p,w}^0 \mu < \infty \text{ and } \mu(\Omega) < \infty \right\}.$$

Remark 3.3. Assume that $\mu \in S_0$ is finite. Set $u = \mathcal{W}_{p,w}^0 \mu$. Then $\sup_\Omega u = \|u\|_{L^\infty(\mu)}$. In fact, clearly $\sup_\Omega u \geq \|u\|_{L^\infty(\mu)}$. To prove the converse inequality, set $k = \|u\|_{L^\infty(\mu)}$. Then testing the equation of u with $(u-k)_+$, we find that

$$\int_{\{u > k\}} |\nabla u|^p \, dw = \int_\Omega (u-k)_+ \, d\mu = 0.$$

Therefore $u(x) \leq k$ for a.e. $x \in \Omega$. Since u is (p, w) -superharmonic in Ω , $u(x) \leq k$ for all $x \in \Omega$.

Remark 3.4. If $\mu \in S_{00}$ and $f \in L^\infty(\mu)$, then $|f|\mu \in S_{00}$.

Lemma 3.5. Let $\mu, \nu \in S_{00}$. Assume also that $\text{supp}(\mu + \nu) \Subset \Omega$. Then $\mu + \nu \in S_{00}$.

Proof. Clearly $\mu + \nu$ is finite and belongs to S_0 . Let $u = \mathcal{W}_{p,w}^0(\mu + \nu)$. By Remark 3.3, $\sup_\Omega u = \|u\|_{L^\infty(\mu + \nu)}$. Let $R = \text{dist}(\text{supp}(\mu + \nu), \partial\Omega)/4$ and fix $x \in \text{supp}(\mu + \nu)$. Then, by the latter inequality in Theorem 2.2, we have

$$u(x) \leq C \left(\inf_{B(x,R)} u + \mathbf{W}_{1,p,w}^{2R}(\mu + \nu)(x) \right).$$

Using a simple calculation and the former inequality in Theorem 2.2, we also obtain

$$\begin{aligned} \mathbf{W}_{1,p,w}^{2R}(\mu + \nu)(x) &\leq \max\{2^{\frac{2-R}{p-1}}, 1\} (\mathbf{W}_{1,p,w}^{2R}\mu(x) + \mathbf{W}_{1,p,w}^{2R}\nu(x)) \\ &\leq C (\mathcal{W}_{p,w}^0\mu(x) + \mathcal{W}_{p,w}^0\nu(x)). \end{aligned}$$

Furthermore, testing the equation of u with $\min\{u, \inf_{B(x,R)} u\}$ and using the Poincaré inequality, we find that

$$\inf_{B(x,R)} u \leq \left(\frac{(\mu + \nu)(\Omega)}{\text{cap}_{p,w}(\overline{B(x,R)}, \Omega)} \right)^{\frac{1}{p-1}} \leq C \left(\text{diam}(\Omega)^p \frac{(\mu + \nu)(\Omega)}{w(B(x,R))} \right)^{\frac{1}{p-1}}.$$

The right-hand side is continuous with respect to x . Hence it is bounded on $\text{supp}(\mu + \nu)$. Combining these estimates, we obtain the desired boundedness. \square

The following characterization for $\mathcal{M}_0^+(\Omega)$ is the nonlinear counterpart of [19, Theorem 2.2.4] (see also [43, Corollary 3.19]). For related characterizations for $\mathcal{M}_0^+(\Omega)$, see also [16] and [48, Proposition 1.2.7].

Theorem 3.6. Let $\mu \in \mathcal{M}^+(\Omega)$. Then, $\mu \in \mathcal{M}_0^+(\Omega)$ if and only if there exists an increasing sequence of compact sets $\{F_k\}_{k=1}^\infty$ such that $\mu_k := \mathbf{1}_{F_k}\mu \in S_{00}$ for all $k \geq 1$ and $\mu(\Omega \setminus \bigcup_{k=1}^\infty F_k) = 0$.

Proof. The ‘‘if’’ part is readily obtained from the definition of S_0 . Let us prove the ‘‘only if’’ part. Consider a sequence of open sets $\{\Omega_j\}_{j=1}^\infty$ such that $\bigcup_{j=1}^\infty \Omega_j = \Omega$ and $\Omega_j \Subset \Omega_{j+1}$ for all $j \geq 1$. For each $j \geq 1$, $\mathbf{1}_{\overline{\Omega_j}}\mu$ is finite. Thus, by [36, Theorem 6.6], there exists a (p, w) -superharmonic function u_j satisfying

$$\begin{cases} -\Delta_{p,w} u_j = \mathbf{1}_{\overline{\Omega_j}}\mu & \text{in } \Omega, \\ \min\{u_j, k\} \in H_0^{1,p}(\Omega; w) & \text{for all } k \geq 1. \end{cases}$$

Let $F_{k,j} = \{u_j \leq k\}$ and let $\mu_{j,k} = \mathbf{1}_{F_{k,j} \cap \overline{\Omega_j}}\mu$. Using the method in Theorem 3.1, we see that $\mu_{j,k} \leq \mathbf{1}_{\{u_j < k+1\}} \mathbf{1}_{\overline{\Omega_j}}\mu \leq \mu[\min\{u_j, k+1\}]$. Since $\min\{u_j, k+1\} \in H_0^{1,p}(\Omega; w)$, this implies that $\mu_{j,k} \in S_0$. By Remark 3.3,

$$\sup_\Omega \mathcal{W}_{p,w}^0 \mu_{j,k} = \|\mathcal{W}_{p,w}^0 \mu_{j,k}\|_{L^\infty(\mu_{j,k})} \leq \|\min\{u_j, k+1\}\|_{L^\infty(\mu_{j,k})} \leq k.$$

Hence $\mu_{j,k} \in S_{00}$. Let

$$F_k = \bigcup_{1 \leq j \leq k} (F_{k,j} \cap \overline{\Omega_j}).$$

Clearly, $F_k \subset F_{k+1}$ and $F_k \subset \overline{\Omega_k}$. Fix $j \geq 1$. Since $\mu \in \mathcal{M}_0^+(\Omega)$, we have $\mu(\{u_j = \infty\}) = 0$. Therefore,

$$\mathbf{1}_{\overline{\Omega_j}} = \lim_{k \rightarrow \infty} \mathbf{1}_{F_{k,j} \cap \overline{\Omega_j}} \leq \lim_{k \rightarrow \infty} \mathbf{1}_{F_k} \leq \mathbf{1}_\Omega \quad \mu\text{-a.e.}$$

Passing to the limit $j \rightarrow \infty$, we see that $\lim_{k \rightarrow \infty} \mathbf{1}_{F_k} = \mathbf{1}_\Omega$ μ -a.e. It remains to be shown that $\mathbf{1}_{F_k} \mu \in S_{00}$. By the comparison principle for weak solutions,

$$\sup_{\Omega} \mathcal{W}_{p,w}^0(\mathbf{1}_{F_k} \mu) \leq \sup_{\Omega} \mathcal{W}_{p,w}^0 \left(\sum_{1 \leq j \leq k} \mu_{j,k} \right).$$

Iterating Lemma 3.5 ($k-1$) times, we find that the right-hand side is finite. \square

Let us now consider the following Dirichlet problem:

$$(3.1) \quad \begin{cases} -\Delta_{p,w} u = \mu & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

We say that a function u is a (p, w) -superharmonic solution (supersolution) to $-\Delta_{p,w} u = \mu$ in Ω , if u is a (p, w) -superharmonic function in Ω and $\mu[u] = \mu$ ($\mu[u] \geq \mu$), where $\mu[u]$ is the Riesz measure of u . We say that a nontrivial nonnegative solution u is *minimal* if $v \geq u$ in Ω whenever v is a nontrivial nonnegative supersolution to the same equation.

Definition 3.7. For $\mu \in \mathcal{M}_0^+(\Omega)$, we define

$$\mathcal{W}_{p,w} \mu(x) := \sup \{ \mathcal{W}_{p,w}^0 \nu(x) : \nu \in S_{00} \text{ and } \nu \leq \mu \}.$$

Remark 3.8. Note that $u := \mathcal{W}_{p,w} \mu$ may be identically infinite. If μ is finite, then u is finite q.e. in Ω and satisfies (3.1) in the sense of entropy or renormalized solutions (see [29, 7, 31, 36, 17]). In general, even if $u \not\equiv \infty$ is a p -superharmonic solution to $-\Delta_{p,w} u = \mu$ in Ω , it does not satisfy the Dirichlet boundary condition in the sense of renormalized solutions. The author is not aware of the renowned name for this class of solutions. Sufficient conditions for $u \not\equiv \infty$ will be discussed in the next section.

Remark 3.9. Clearly, $\mathcal{W}_{p,w}(a\mu) = a^{\frac{1}{p-1}} \mathcal{W}_{p,w} \mu$ for any constant $a \geq 0$, and

$$\mu \leq \nu \Rightarrow \mathcal{W}_{p,w} \mu(x) \leq \mathcal{W}_{p,w} \nu(x), \quad \forall x \in \Omega.$$

Furthermore, Theorem 3.1 is still valid when $u = \mathcal{W}_{p,w} \mu$.

Proposition 3.10. Let $\mu \in \mathcal{M}_0^+(\Omega)$. The following statements hold.

- (i) Assume that $\mathcal{W}_{p,w} \mu \not\equiv \infty$. Then $u = \mathcal{W}_{p,w} \mu$ is the minimal nonnegative (p, w) -superharmonic solution to $-\Delta_{p,w} u = \mu$ in Ω .
- (ii) Let $\{f_j\}_{j=1}^\infty \subset L_{\text{loc}}^1(\mu)$ be a nondecreasing sequence of functions. Assume that $f_j \uparrow f$ μ -a.e. Let $u_j = \mathcal{W}_{p,w}(f_j \mu)$ and let $u = \lim_{j \rightarrow \infty} u_j$. Assume also that $u \not\equiv \infty$. Then $u = \mathcal{W}_{p,w}(f \mu)$.
- (iii) If $\mu \in S_0$, then $\mathcal{W}_{p,w} \mu = \mathcal{W}_{p,w}^0 \mu$.

Proof. (i) Let $\{\mu_k\}_{k=1}^\infty$ be a sequence of Radon measures in Theorem 3.6. By the comparison principle for weak solutions and the definition of u , for each $k \geq 1$,

$$0 \leq \mathcal{W}_{p,w}^0 \mu_k(x) \leq \mathcal{W}_{p,w}^0 \mu_{k+1}(x) \leq u(x), \quad \forall x \in \Omega.$$

Set

$$u'(x) = \lim_{k \rightarrow \infty} \mathcal{W}_{p,w}^0 \mu_k(x).$$

By [27, Lemma 7.3] and Theorem 2.1, u' is a nonnegative (p, w) -superharmonic solution to $-\Delta_{p,w}u = \mu$ in Ω . On the other hand, by Theorem 3.1, if $\nu \leq \mu$ and $\nu \in S_{00}$, then $\mathcal{W}_{p,w}^0\nu \leq u'$ in Ω . Thus, $u = u'$. Using the same argument again, we see that u is minimal. (ii) Let ω be the Riesz measure of u . By Theorem 2.1, $f_k\mu$ converges to ω weakly. By the monotone convergence theorem, $\omega = f\mu$ and $f\mu \in \mathcal{M}_0^+(\Omega)$. For each $j \geq 1$, $u_j \leq \mathcal{W}_{p,w}(f\mu)$, and hence $u \leq \mathcal{W}_{p,w}(f\mu)$. By the minimality of $\mathcal{W}_{p,w}(f\mu)$, $u = \mathcal{W}_{p,w}(f\mu)$. (iii) Set $u_k = \mathcal{W}_{p,w}^0\mu_k$. Since $\mu_k \in S_{00}$ and $\mu \in S_0$, we have

$$\int_{\Omega} |\nabla u_k|^p dw = \int_{\Omega} u_k d\mu_k \leq \int_{\Omega} u_k d\mu \leq C \|\nabla u_k\|_{L^p(\Omega; w)}.$$

Passing to the limit $k \rightarrow \infty$, we see that $u = \lim_{k \rightarrow \infty} u$ belongs to $H_0^{1,p}(\Omega; w)$. From Theorem 2.1 and the uniqueness of weak solutions, the assertion follows. \square

Lemma 3.11. *Let $\mu \in \mathcal{M}_0^+(\Omega)$, and let $u = \mathcal{W}_{p,w}\mu$. Then, $\sup_{\Omega} u = \|u\|_{L^\infty(\mu)}$.*

Proof. Take $\{\mu_k\}_{k=1}^\infty$ by using Theorem 3.6. Set $u_k = \mathcal{W}_{p,w}^0\mu_k$. By Remark 3.3, for any $x \in \Omega$,

$$u(x) = \lim_{k \rightarrow \infty} u_k(x) \leq \lim_{k \rightarrow \infty} \sup_{\Omega} u_k = \lim_{k \rightarrow \infty} \|u_k\|_{L^\infty(\mu_k)} \leq \|u\|_{L^\infty(\mu)}.$$

The converse inequality is clear. \square

4. GENERALIZED ENERGY

Let us define generalized p -energy by using minimal p -superharmonic solutions. For $p = 2$ or $\Omega = \mathbb{R}^n$, we refer to [42, 25].

Definition 4.1. For $0 \leq \gamma < \infty$, set $S^\gamma := \{\mu \in \mathcal{M}_0^+(\Omega) : \mathcal{E}_\gamma(\mu) < \infty\}$, where

$$\mathcal{E}_\gamma(\mu) := \int_{\Omega} (\mathcal{W}_{p,w}\mu)^\gamma d\mu.$$

Furthermore, let $S^\infty := \{\mu \in \mathcal{M}_0^+(\Omega) : \|\mathcal{W}_{p,w}\mu\|_{L^\infty(\mu)} < \infty\}$ for $\gamma = \infty$.

By definition, if $\mu \in S^\gamma$, then the Dirichlet problem (3.1) has a minimal non-negative (p, w) -superharmonic solution $u = \mathcal{W}_{p,w}\mu$. We can verify $S^1 = S_0$ by modifying the proof of Theorem 1.3 below. For any $0 \leq \gamma \leq \infty$,

$$S_{00} = S^0 \cap S^\infty \subset S^\gamma \subset \mathcal{M}_0^+(\Omega).$$

Assume that $0 < \gamma < \infty$ and $\mu \in S_{00}$. Let $u = \mathcal{W}_{p,w}^0\mu$ and let $v = u^{\frac{p-1+\gamma}{p}}$. Then, for any $\epsilon > 0$,

$$\int_{\Omega} (u^\gamma - \epsilon)_+ d\mu = \gamma \int_{\{u^\gamma > \epsilon\}} |\nabla u|^p u^{\gamma-1} dw = \frac{1}{c_E} \int_{\Omega} |\nabla v_\epsilon|^p dw,$$

where $v_\epsilon = (v - \epsilon^{\frac{p-1+\gamma}{\gamma p}})_+$ and c_E is the constant in (1.5). Thus, by the monotone convergence theorem,

$$(4.1) \quad \int_{\Omega} (\mathcal{W}_{p,w}^0\mu)^\gamma d\mu = \gamma \int_{\Omega} |\nabla u|^p u^{\gamma-1} dw = \frac{1}{c_E} \int_{\Omega} |\nabla v|^p dw.$$

Moreover, since $\{v_\epsilon\}$ is a bounded sequence in $H_0^{1,p}(\Omega; w)$, $v \in H_0^{1,p}(\Omega; w)$.

Proposition 4.2. *Let $\mu \in \mathcal{M}_0^+(\Omega)$, and let $u = \mathcal{W}_{p,w}\mu$.*

(i) *If $\mu \in S^\gamma$ with $0 \leq \gamma \leq 1$, then $\min\{u, l\} \in H_0^{1,p}(\Omega; w)$ for all $l > 0$.*

- (ii) If $\mu \in S^\gamma$ with $1 \leq \gamma \leq \infty$, then u belongs to $H_{\text{loc}}^{1,p}(\Omega; w)$ and satisfies $-\Delta_{p,w}u = \mu$ in the sense of weak solutions.
- (iii) Assume that $\mu \in S^{\gamma_0} \cap S^{\gamma_1}$ with $0 \leq \gamma_0 \leq 1$ and $1 \leq \gamma_1 \leq \infty$. Then $u = \mathcal{W}_{p,w}^0\mu \in H_0^{1,p}(\Omega; w)$. In particular, u satisfies (3.1) in the sense of finite energy weak solutions.

Proof. Let $\{\mu_k\}_{k=1}^\infty$ be a sequence of Radon measures in Theorem 3.6, and let $u_k = \mathcal{W}_{p,w}^0\mu_k$. (i) Testing the equation of u_k with $\min\{u_k, l\} \in H_0^{1,p}(\Omega; w)$, we get

$$(4.2) \quad \int_{\Omega} |\nabla \min\{u_k, l\}|^p dw = \int_{\Omega} \min\{u_k, l\} d\mu_k \leq l^{1-\gamma} \int_{\Omega} (\mathcal{W}_{p,w}\mu)^\gamma d\mu.$$

Letting $k \rightarrow \infty$ gives the desired boundary condition. (ii) Assume that $1 \leq \gamma < \infty$. Take a ball B such that $B \Subset \Omega$. Without loss of generality, we may assume that $u_{k_0} \neq 0$ for some $k_0 \geq 1$. By the strong minimum principle, $\inf_{\bar{B}} u_{k_0} > 0$. Let $k \geq k_0$. Subsequently, (4.1) gives

$$\begin{aligned} \int_B |\nabla u_k|^p dw &\leq \frac{1}{\gamma} \left(\inf_{\bar{B}} u_k \right)^{1-\gamma} \int_{\Omega} (\mathcal{W}_{p,w}\mu_k)^\gamma d\mu_k \\ &\leq \frac{1}{\gamma} \left(\inf_{\bar{B}} u_{k_0} \right)^{1-\gamma} \int_{\Omega} (\mathcal{W}_{p,w}\mu)^\gamma d\mu. \end{aligned}$$

On the other hand, by the Poincaré inequality,

$$\begin{aligned} \left(\int_B u_k^p dw \right)^{\frac{p-1+\gamma}{p}} &\leq w(B)^{\frac{\gamma-1}{p}} \int_{\Omega} u_k^{p-1+\gamma} dw \leq C \int_{\Omega} |\nabla u_k|^{\frac{p-1+\gamma}{p}} dw \\ &= c_E C \int_{\Omega} (\mathcal{W}_{p,w}\mu_k)^\gamma d\mu_k \leq c_E C \int_{\Omega} (\mathcal{W}_{p,w}\mu)^\gamma d\mu. \end{aligned}$$

Hence, $\|u_k\|_{H^{1,p}(B;w)}$ is bounded. Passing to the limit $k \rightarrow \infty$, we see that $u \in H^{1,p}(B;w)$. Thus, $u \in H_{\text{loc}}^{1,p}(\Omega; w)$. If $\mu \in S^\infty$, then u is a bounded (p, w) -superharmonic function. Therefore, $u \in H_{\text{loc}}^{1,p}(\Omega; w)$. (iii) By Hölder's inequality, $S^{\gamma_0} \cap S^{\gamma_1} \subset S^1 = S_0$. Hence, $u = \mathcal{W}_{p,w}\mu \in H_0^{1,p}(\Omega; w)$. \square

As in the proof of [25, Lemma 3.3], the Picone-type inequality in [1, 9] yields the following estimate.

Lemma 4.3. *Let $1 < p < \infty$, $0 < \gamma < \infty$ and $-\gamma < q < p - 1$. Suppose that $\nu \in S_{00}$ and $v = \mathcal{W}_{p,w}\nu$. Assume that $u \in H_0^{1,p}(\Omega; w) \cap L^\infty(\Omega)$ and $u \geq 0$ in Ω . Then,*

$$\begin{aligned} \int_{\Omega} \tilde{u}^{\gamma+q} d\nu &\leq \left(\left(\frac{p-1+\gamma}{p} \right)^p \int_{\Omega} |\nabla u|^p u^{\gamma-1} dw \right)^{\frac{\gamma+q}{p-1+\gamma}} \\ &\quad \times \left(\int_{\Omega} v^{\frac{(\gamma+q)(p-1)}{p-1-q}} d\nu \right)^{\frac{p-1-q}{p-1+\gamma}}. \end{aligned}$$

Proof of Theorem 1.3. We first prove the upper bound of C_T . Take a sequence of measures $\{\sigma_k\}_{k=1}^\infty \subset S_{00}$ by using Theorem 3.6. Apply Lemma 4.3 to $u_k = \min\{|u|, k\}$ and σ_k ; we get

$$\int_{\Omega} u_k^{1+q} d\sigma_k \leq \left(\int_{\Omega} |\nabla u_k|^p dw \right)^{\frac{1+q}{p}} \left(\int_{\Omega} (\mathcal{W}_{p,w}\sigma_k)^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma_k \right)^{\frac{p-1-q}{p}}.$$

The desired estimate then follows from the monotone convergence theorem. Let us prove the lower bound. Take $\{\sigma_k\}_{k=1}^\infty \subset S_{00}$. Let $u = \mathcal{W}_{p,w}^0 \sigma_k$ and let $v = u^{\frac{p-1}{p-1-q}}$. Note that

$$\nabla v = \frac{p-1}{p-1-q} \nabla u u^{\frac{q}{p-1-q}} \quad \text{a.e. in } \Omega.$$

Thus, using (4.1) with $\gamma = \frac{(1+q)(p-1)}{p-1-q}$, we get

$$\begin{aligned} \int_{\Omega} |\nabla v|^p dw &= \left(\frac{p-1}{p-1-q} \right)^p \int_{\Omega} |\nabla u|^p u^{\frac{pq}{p-1-q}} dw \\ &= \frac{1}{1+q} \left(\frac{p-1}{p-1-q} \right)^{p-1} \int_{\Omega} u^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma_k. \end{aligned}$$

By density, (1.2) gives

$$\begin{aligned} \left(\int_{\Omega} u^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma_k \right)^{\frac{1}{1+q}} &\leq \|v\|_{L^{1+q}(\sigma)} \leq C_T \|\nabla v\|_{L^p(w)} \\ &= \left(\frac{1}{1+q} \right)^{\frac{1}{p}} \left(\frac{p-1}{p-1-q} \right)^{\frac{p-1}{p}} C_T \left(\int_{\Omega} u^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma_k \right)^{\frac{1}{p}}. \end{aligned}$$

Therefore,

$$\int_{\Omega} (\mathcal{W}_{p,w} \sigma_k)^{\frac{(1+q)(p-1)}{p-1-q}} d\sigma_k \leq \left(\frac{1}{1+q} \right)^{\frac{1+q}{p-1-q}} \frac{1}{c_V^{1+q}} C_T^{\frac{(1+q)p}{p-1-q}}.$$

Passing to the limit $k \rightarrow \infty$, we arrive at the desired lower bound. \square

Corollary 4.4. *Let $\mu \in \mathcal{M}_0^+(\Omega)$ and let $0 < \gamma < \infty$. Then, $\mu \in S^\gamma$ if and only if $v := (\mathcal{W}_{p,w} \mu)^{\frac{p-1+\gamma}{p}} \in H_0^{1,p}(\Omega; w)$. Moreover,*

$$\int_{\Omega} (\mathcal{W}_{p,w} \mu)^\gamma d\mu \leq \int_{\Omega} |\nabla v|^p dw \leq c_E \int_{\Omega} (\mathcal{W}_{p,w} \mu)^\gamma d\mu.$$

Proof. Let $\{\mu_k\}_{k=1}^\infty$ be a sequence of Radon measures in Theorem 3.6. Set $u_k = \mathcal{W}_{p,w}^0 \mu_k$ and $v_k = (u_k)^{\frac{p-1+\gamma}{p}}$. Assume that $\mu \in S^\gamma$. Then, by (4.1), $\{v_k\}_{k=1}^\infty$ is a bounded sequence in $H_0^{1,p}(\Omega; w)$. Since $v_k \uparrow v$, v belongs to $H_0^{1,p}(\Omega; w)$ and satisfies the latter inequality. Conversely, assume that $v \in H_0^{1,p}(\Omega; w)$. By Theorem 1.3,

$$\int_{\Omega} |\varphi|^{1+q} d\mu_k \leq \left(\int_{\Omega} u_k^\gamma d\mu_k \right)^{\frac{p-1-q}{p}} \left(\int_{\Omega} |\nabla \varphi|^p dw \right)^{\frac{1+q}{p}}, \quad \forall \varphi \in C_c^\infty(\Omega),$$

where $q = \frac{\gamma p}{p-1+\gamma} - 1$. By density, the same inequality holds for any $\varphi \in H_0^{1,p}(\Omega; w)$. Therefore,

$$\int_{\Omega} u_k^\gamma d\mu_k \leq \int_{\Omega} v^{1+q} d\mu_k \leq \left(\int_{\Omega} u_k^\gamma d\mu_k \right)^{\frac{p-1-q}{p}} \left(\int_{\Omega} |\nabla v|^p dw \right)^{\frac{1+q}{p}}.$$

Then the monotone convergence theorem yields

$$\int_{\Omega} (\mathcal{W}_{p,w} \mu)^\gamma d\mu = \lim_{k \rightarrow \infty} \int_{\Omega} u_k^\gamma d\mu_k \leq \int_{\Omega} |\nabla v|^p dw.$$

This completes the proof. \square

We also obtain the following estimate using a similar approximation argument.

Theorem 4.5. *Let $1 < p < \infty$, $0 < \gamma < \infty$ and $-\gamma < q < p - 1$. Then, for any $\mu, \nu \in \mathcal{M}_0^+(\Omega)$,*

$$(4.3) \quad \int_{\Omega} (\mathcal{W}_{p,w}\mu)^{\gamma+q} d\nu \leq \left(c_E \int_{\Omega} (\mathcal{W}_{p,w}\mu)^{\gamma} d\mu \right)^{\frac{\gamma+q}{p-1+\gamma}} \mathcal{E}_{\frac{(\gamma+q)(p-1)}{p-1-q}}(\nu)^{\frac{p-1-q}{p-1+\gamma}}.$$

In particular, if $\mu \in S^{\gamma}$ and $\nu \in S^{\frac{(\gamma+q)(p-1)}{p-1-q}}$, then $\mathcal{W}_{p,w}\mu \in L^{\gamma+q}(\nu)$.

Remark 4.6. Note that $c_E = 1$ if $\gamma = 1$. This sharp constant is achieved when $\mu = \nu$ is an equilibrium measure.

Remark 4.7. For $0 < \gamma < \infty$, set $\|\mu\|_{\gamma} = \mathcal{E}_{\gamma}(\mu)^{\frac{p-1}{p-1+\gamma}}$. Then by Theorem 4.5,

$$\begin{aligned} \int_{\Omega} \mathcal{W}_{p,w}(\mu + \nu)^{\gamma} d(\mu + \nu) &= \int_{\Omega} \mathcal{W}_{p,w}(\mu + \nu)^{\gamma} d\mu + \int_{\Omega} \mathcal{W}_{p,w}(\mu + \nu)^{\gamma} d\nu \\ &\leq c_E^{\frac{\gamma}{p-1+\gamma}} \left(\int_{\Omega} \mathcal{W}_{p,w}(\mu + \nu)^{\gamma} d(\mu + \nu) \right)^{\frac{\gamma}{p-1+\gamma}} \left(\|\mu\|_{\gamma} + \|\nu\|_{\gamma} \right), \end{aligned}$$

and hence

$$\|\mu + \nu\|_{\gamma} \leq c_E^{\frac{\gamma}{p-1+\gamma}} \left(\|\mu\|_{\gamma} + \|\nu\|_{\gamma} \right).$$

In particular, S^{γ} is a convex cone.

Finally, we prove weighted norm inequalities.

Theorem 4.8. *Let $1 < p < \infty$, $0 < q < p - 1$ and $0 < \gamma < \infty$. Assume that $\sigma \in S^{\frac{(\gamma+q)(p-1)}{p-1-q}}$. Then, for any $f \in L^{\frac{\gamma+q}{q}}(\sigma)$, $|f|\sigma \in S^{\gamma}$. Moreover,*

$$\mathcal{E}_{\gamma}(|f|\sigma)^{\frac{p-1}{p-1+\gamma}} \leq \left(c_E \mathcal{E}_{\frac{(\gamma+q)(p-1)}{p-1-q}}(\sigma)^{\frac{p-1-q}{\gamma+q}} \right)^{\frac{\gamma}{p-1+\gamma}} \|f\|_{L^{\frac{\gamma+q}{q}}(\sigma)}$$

and

$$\|\mathcal{W}_{p,w}(|f|\sigma)\|_{L^{\gamma+q}(\sigma)} \leq \left(c_E \mathcal{E}_{\frac{(\gamma+q)(p-1)}{p-1-q}}(\sigma)^{\frac{p-1-q}{\gamma+q}} \right)^{\frac{1}{p-1}} \|f\|_{L^{\frac{\gamma+q}{q}}(\sigma)}^{\frac{1}{p-1}}.$$

Proof. We may assume that $f \geq 0$ without loss of generality. By Theorem 4.5,

$$\int_{\Omega} (\mathcal{W}_{p,w}(f\sigma))^{\gamma+q} d\sigma \leq \left(c_E \int_{\Omega} (\mathcal{W}_{p,w}(f\sigma))^{\gamma} f d\sigma \right)^{\frac{\gamma+q}{p-1+\gamma}} \mathcal{E}_{\frac{(\gamma+q)(p-1)}{p-1-q}}(\sigma)^{\frac{p-1-q}{p-1+\gamma}}.$$

Meanwhile, by Hölder's inequality,

$$\int_{\Omega} (\mathcal{W}_{p,w}(f\sigma))^{\gamma} f d\sigma \leq \left(\int_{\Omega} (\mathcal{W}_{p,w}(f\sigma))^{\gamma+q} d\sigma \right)^{\frac{\gamma}{\gamma+q}} \|f\|_{L^{\frac{\gamma+q}{q}}(\sigma)}.$$

Combining the two inequalities, we obtain the desired estimates. \square

Remark 4.9. Under the same assumptions, suppose also that $\nu \in S^{\frac{(\gamma+Q)(p-1)}{p-1-Q}}$ with $-\gamma < Q < p - 1$. Then, the following two weight norm inequality holds:

$$\|\mathcal{W}_{p,w}(|f|\sigma)\|_{L^{\gamma+Q}(\nu)} \leq C \|f\|_{L^{\frac{\gamma+q}{q}}(\sigma)}^{\frac{1}{p-1}}, \quad \forall f \in L^{\frac{\gamma+q}{q}}(\sigma).$$

In fact, by Theorem 4.5,

$$\int_{\Omega} (\mathcal{W}_{p,w}(f\sigma))^{\gamma+Q} d\nu \leq C \left(\int_{\Omega} (\mathcal{W}_{p,w}(f\sigma))^{\gamma} f d\sigma \right)^{\frac{\gamma+Q}{p-1+\gamma}} \mathcal{E}_{\frac{(\gamma+Q)(p-1)}{p-1-Q}}(\nu)^{\frac{p-1-Q}{p-1+\gamma}}.$$

The right-hand side is estimated by Theorem 4.8.

5. PROPERTIES OF SOLUTIONS TO (1.1)

First, we give the counterpart of [12, Lemma 3.5] or [22, Remark 2.6].

Lemma 5.1. *Let $\sigma \in \mathcal{M}_0^+(\Omega)$, and let $\beta \geq 1$. Assume that $(\mathcal{W}_{p,w}\sigma)^{(\beta-1)(p-1)} \sigma \in \mathcal{M}_0^+(\Omega)$. Then,*

$$(\mathcal{W}_{p,w}\sigma)^\beta(x) \leq \beta \mathcal{W}_{p,w} \left((\mathcal{W}_{p,w}\sigma)^{(\beta-1)(p-1)} \sigma \right) (x), \quad \forall x \in \Omega.$$

Proof. We use the argument in [25, Lemma 4.4]. We first assume that $\sigma \in S_{00}$. Let $u = \mathcal{W}_{p,w}\sigma$. Since u is bounded on Ω , $u^\beta \in H_0^{1,p}(\Omega; w)$ and $u^{(\beta-1)(p-1)}\sigma \in S_{00}$. Fix a nonnegative function $\varphi \in C_c^\infty(\Omega)$. Testing the equation of u with $\varphi(u^{(\beta-1)(p-1)} - \epsilon)_+$, we find that

$$\begin{aligned} \int_{\Omega} \varphi(u^{(\beta-1)(p-1)} - \epsilon)_+ d\sigma &= \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \left(\varphi(u^{(\beta-1)(p-1)} - \epsilon)_+ \right) dw \\ &\geq \beta^{1-p} \int_{\{u^{(\beta-1)(p-1)} > \epsilon\}} |\nabla u^\beta|^{p-2} \nabla u^\beta \cdot \nabla \varphi dw. \end{aligned}$$

Applying the dominated convergence theorem to the right-hand side, we obtain

$$\int_{\Omega} \varphi u^{(\beta-1)(p-1)} d\sigma \geq \beta^{1-p} \int_{\Omega} |\nabla u^\beta|^{p-2} \nabla u^\beta \cdot \nabla \varphi dw.$$

By the comparison principle for weak solutions, this implies that

$$u^\beta \leq \mathcal{W}_{p,w}^0(\beta^{p-1} u^{(\beta-1)(p-1)} \sigma) \quad \text{a.e. in } \Omega.$$

Since u is (p, w) -superharmonic in Ω , the desired inequality holds.

For $\sigma \in \mathcal{M}_0^+(\Omega)$, we take a sequence of measures $\{\sigma_k\}_{k=1}^\infty \subset S_{00}$ by using Theorem 3.6. By the above result,

$$\begin{aligned} (\mathcal{W}_{p,w}\sigma_k)^\beta(x) &\leq \beta \mathcal{W}_{p,w} \left((\mathcal{W}_{p,w}\sigma_k)^{(\beta-1)(p-1)} \sigma_k \right) (x) \\ &\leq \beta \mathcal{W}_{p,w} \left((\mathcal{W}_{p,w}\sigma)^{(\beta-1)(p-1)} \sigma \right) (x), \quad \forall x \in \Omega. \end{aligned}$$

Taking the supremum over $k \geq 1$, we arrive at the desired estimate. \square

Next, we give the counterpart of [12, Theorem 3.4] or [22, Theorem 1.3].

Theorem 5.2. *Let $1 < p < \infty$ and $0 < q < p-1$. Let $\sigma \in \mathcal{M}_0^+(\Omega)$. Let $v \in L_{\text{loc}}^q(\sigma)$ be a nontrivial nonnegative (p, w) -superharmonic supersolution to $-\Delta_{p,w}v = \sigma v^q$ in Ω . Then,*

$$v(x) \geq c_V (\mathcal{W}_{p,w}\sigma)^{\frac{p-1}{p-1-q}}(x), \quad \forall x \in \Omega,$$

where c_V is the constant in (1.5).

Proof. For simplicity, we write $\mathcal{W}_{p,w}\mu$ as $\mathcal{W}\mu$. Let $u = \mathcal{W}(v^q\sigma)$. By Theorem 3.1, $v(x) \geq u(x)$ for all $x \in \Omega$. Fix $a > 0$, and set $\sigma_a = \mathbf{1}_{\{x \in \Omega: u(x) > a\}}\sigma$. Using Theorem 3.1 again, we get

$$u \geq \mathcal{W}(u^q\sigma) \geq \mathcal{W}(a^q\sigma_a) = a^{\frac{q}{p-1}} \mathcal{W}\sigma_a.$$

Continuing this argument k -times, we obtain

$$\begin{aligned} (5.1) \quad u &\geq \mathcal{W}(\mathcal{W}(\cdots (\mathcal{W}(u^q\sigma))^q \cdots \sigma)^q\sigma) \\ &\geq \mathcal{W}(\mathcal{W}(\cdots (\mathcal{W}(a^q\sigma_a))^q \cdots \sigma)^q\sigma) \\ &\geq a^{\left(\frac{q}{p-1}\right)^k} \mathcal{W}(\mathcal{W}(\cdots (\mathcal{W}\sigma_a)^q \cdots \sigma_a)^q\sigma_a). \end{aligned}$$

Meanwhile, by Lemma 5.1,

$$(\mathcal{W}\sigma_a)^{\beta_{i+1}} \leq \beta_{i+1} \mathcal{W}\left((\mathcal{W}\sigma_a)^{\beta_i q} \sigma_a\right)$$

for each $i \geq 0$, where $\beta_0 = 1$ and $\beta_{i+1} = \beta_i \frac{q}{p-1} + 1$. Iterating this estimate k times, we get

$$(5.2) \quad (\mathcal{W}\sigma_a)^{\beta_k} \leq \prod_{i=1}^k \beta_i \left(\frac{q}{p-1}\right)^{k-i} \mathcal{W}(\mathcal{W}(\cdots (\mathcal{W}\sigma_a)^q \cdots \sigma_a)^q \sigma_a).$$

By definition, $\beta_k = \sum_{i=0}^k \left(\frac{q}{p-1}\right)^i$. Therefore $\beta_k \uparrow \frac{p-1}{p-1-q}$ as $k \rightarrow \infty$ and

$$(5.3) \quad \prod_{i=1}^k \beta_i \left(\frac{q}{p-1}\right)^{k-i} \leq \left(\frac{p-1}{p-1-q}\right)^{\sum_{i=1}^k \left(\frac{q}{p-1}\right)^{k-i}} \leq \left(\frac{p-1}{p-1-q}\right)^{\frac{p-1}{p-1-q}}.$$

Combining (5.1), (5.2) and (5.3) and letting $k \rightarrow \infty$, we obtain

$$u \geq \left(\frac{p-1-q}{p-1}\right)^{\frac{p-1}{p-1-q}} (\mathcal{W}\sigma_a)^{\frac{p-1}{p-1-q}}.$$

Without loss of generality, we may assume that $\sigma_a \neq 0$ for small $a > 0$. Then $u \geq a^{\frac{q}{p-1}} \mathcal{W}\sigma_a > 0$ in Ω by the strong minimum principle. Thus, taking the limit $a \rightarrow 0$, we arrive at the desired estimate. \square

Finally, we prove Theorem 1.1 and its variants.

Proof of Theorem 1.1. (1) \Rightarrow (2): Using Theorem 5.2, we find that

$$\left(\int_{\Omega} (\mathcal{W}_{p,w}\sigma)^{\frac{(\gamma+q)(p-1)}{p-1-q}} d\sigma\right)^{\frac{1}{\gamma+q}} \leq \frac{1}{c_V} \|v\|_{L^{\gamma+q}(\sigma)} \leq \frac{C_1}{c_V}.$$

(2) \Rightarrow (3): By Theorem 4.8, (1.4) holds with

$$C_3^{\frac{p-1}{p-1-q}} \leq \left(c_E \mathcal{E}_{\frac{(\gamma+q)(p-1)}{p-1-q}}(\sigma)^{\frac{p-1-q}{\gamma+q}}\right)^{\frac{1}{p-1-q}} \leq c_E^{\frac{1}{p-1-q}} C_2.$$

(3) \Rightarrow (1): Take $\{\mathbf{1}_{F_k}\sigma\}_{k=1}^{\infty} \subset S_{00}$ by using Theorem 3.6. Applying (1.4) to $f = (\mathcal{W}\sigma_k)^{\frac{q(p-1)}{p-1-q}} \mathbf{1}_{F_k} \in L^{\frac{\gamma+q}{q}}(\sigma)$ and using Lemma 5.1, we obtain

$$\left(\int_{\Omega} (\mathcal{W}_{p,w}\sigma_k)^{\frac{(\gamma+q)(p-1)}{p-1-q}} \mathbf{1}_{F_k} d\sigma\right)^{\frac{1}{\gamma+q}} \leq C.$$

Therefore, by the monotone convergence theorem,

$$u_0 := c_V (\mathcal{W}_{p,w}\sigma)^{\frac{p-1}{p-1-q}} \in L^{\gamma+q}(\sigma),$$

where c_V is the constant in (1.5). Define a sequence of (p, w) -superharmonic functions $\{u_i\}_{i=1}^{\infty}$ by

$$u_{i+1} := \mathcal{W}_{p,w}(u_i^q \sigma), \quad i = 1, 2, \dots$$

By (1.4),

$$\|u_{i+1}\|_{L^{\gamma+q}(\sigma)} = \|\mathcal{W}_{p,w}(u_i^q \sigma)\|_{L^{\gamma+q}(\sigma)} \leq C_3 \|u_i\|_{L^{\gamma+q}(\sigma)}^{\frac{q}{p-1}},$$

and hence $\{u_i\}_{i=1}^\infty \subset L^{\gamma+q}(\sigma) \subset L^q_{\text{loc}}(\sigma)$. By Lemma 5.1, $u_0 \leq u_1$. Hence, by induction, $u_i \leq u_{i+1}$ for all $i \geq 1$. Let $u = \lim_{i \rightarrow \infty} u_i$. By the monotone convergence theorem,

$$\|u\|_{L^{\gamma+q}(\sigma)} = \lim_{i \rightarrow \infty} \|u_{i+1}\|_{L^{\gamma+q}(\sigma)} \leq C_3^{\frac{p-1}{p-1-q}}.$$

By Proposition 3.10, u is (p, w) -superharmonic in Ω and $u = \mathcal{W}_{p,w}(u^q \sigma)$.

Assume that v is a nontrivial nonnegative (p, w) -superharmonic solution to $-\Delta_{p,w} v = \sigma v^q$ in Ω . Then $u_0 \leq v$ by Theorem 5.2, and hence, $u_i \leq v$ for all $i \geq 1$ by induction. Therefore $u \leq v$. \square

Remark 5.3. In Theorem 1.1, the equivalence (1) \Leftrightarrow (2) still holds even if $q = 0$.

Theorem 5.4. *Let Ω be a bounded domain in \mathbb{R}^n . Let $1 < p < \infty$ and $0 < q < p-1$. Suppose that $\sigma \in \mathcal{M}_0^+(\Omega) \setminus \{0\}$. Then, the following statements are equivalent:*

- (1) *There exists a bounded positive weak supersolution $v \in H_{\text{loc}}^{1,p}(\Omega; w)$ to $-\Delta_{p,w} v = \sigma v^q$ in Ω satisfying $\|v\|_{L^\infty(\sigma)} \leq C_1 < \infty$.*
- (2) *$\|\mathcal{W}_{p,w} \sigma\|_{L^\infty(\sigma)}^{\frac{p-1}{p-1-q}} \leq C_2 < \infty$.*
- (3) *The following weighted norm inequality holds:*

$$\|\mathcal{W}_{p,w}(|f|\sigma)\|_{L^\infty(\sigma)} \leq C_3 \|f\|_{L^\infty(\sigma)}^{\frac{1}{p-1}}, \quad \forall f \in L^\infty(\sigma).$$

Moreover, if C_i ($i = 1, 2, 3$) are the best constants in the above statements, then

$$C_1 \leq C_3^{\frac{p-1}{p-1-q}} \leq C_2 \leq \frac{C_1}{c_V}.$$

In addition, if one of the above statements holds, then there exists a minimal positive (p, w) -superharmonic solution u to $-\Delta_{p,w} u = \sigma u^q$ in Ω such that $\sup_\Omega u \leq C_1$.

Proof. We modify the Proof of Theorem 1.1. Clearly, (1) \Rightarrow (2) \Rightarrow (3). Assume that (3) holds. Then, for all $i \geq 1$,

$$\|u_{i+1}\|_{L^\infty(\sigma)} = \|\mathcal{W}_{p,w}(u_i^q \sigma)\|_{L^\infty(\sigma)} \leq C_3 \|u_i\|_{L^\infty(\sigma)}^{\frac{q}{p-1}}.$$

Thus,

$$\|u\|_{L^\infty(u^q \sigma)} \leq \|u\|_{L^\infty(\sigma)} \leq C_3^{\frac{p-1}{p-1-q}}.$$

By Lemma 3.11, $\sup_\Omega u \leq C_3^{\frac{p-1}{p-1-q}}$ and (1) holds. \square

Proof of Theorem 1.2. The case of $q = 0$ is [27, Corollary 21.18], so we consider $q > 0$. Assume that a positive finite energy solution u exists. We may assume that u is quasicontinuous without loss of generality. Then, we find that

$$\|\nabla u\|_{L^p(w)}^p = \int_\Omega |\nabla u|^p dw = \int_\Omega u^{1+q} d\sigma = \|u\|_{L^{1+q}(\sigma)}^{1+q}.$$

Therefore, (1.6) follows from Theorem 1.1. Conversely, assume that (1.6) holds. Then, Theorem 1.1 and Proposition 4.2 give a minimal positive finite energy weak solution $u \in H_0^{1,p}(\Omega; w)$ to $-\Delta_{p,w} u = \sigma u^q$ in Ω . The uniqueness of such a solution follows from a convexity argument as in [13, Theorem 5.1]. \square

Remark 5.5. As the proof of [25, Corollary 6.3], using Remark 4.7, we can construct weak solutions to quasilinear equations of the type

$$\begin{cases} -\Delta_{p,w} u = \sum_{j=1}^J \sigma_j u^{q_j} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $0 \leq q_j < p - 1$ and $\sigma_j \in S^{\frac{(1+q_j)(p-1)}{p-1-q_j}}$ for $j = 1, 2, \dots, J$.

6. QUASILINEAR PDE WITH $L^{s,t}$ COEFFICIENTS

Let us assume that $\mu \in \mathcal{M}_0^+(\Omega)$ is finite. Then as in [37, Theorem 2.1],

$$(6.1) \quad \mathcal{W}_{p,w}\mu(x) \leq C \mathbf{W}_{1,p,w}^{2\text{diam}(\Omega)} \mu(x), \quad \forall x \in \Omega.$$

Using this, we can estimate the generalized p -energy of μ .

We now consider unweighted equations. As the usual notation, we write $H_0^{1,p}(\Omega; 1)$ as $W_0^{1,p}(\Omega)$. For a Lebesgue measurable function f on Ω , we define

$$\|f\|_{L^{r,\rho}(\Omega)} = \begin{cases} \left(\int_0^\infty \left(t^{\frac{1}{r}} f^*(t) \right)^\rho \frac{dt}{t} \right)^{\frac{1}{\rho}} & \text{if } \rho < \infty, \\ \sup_{t>0} t^{\frac{1}{r}} f^*(t) & \text{if } \rho = \infty, \end{cases}$$

where $0 < r, \rho \leq \infty$ and $f^*(t) = \inf\{\alpha > 0 : |\{x \in \Omega : |f(x)| > \alpha\}| \leq t\}$. The space of all f with $\|f\|_{L^{r,\rho}(\Omega)} < \infty$ is called the *Lorentz space*. For the basics of Lorentz spaces, we refer to [20, Chapter 1].

Corollary 6.1. *Let $1 < p < n$, $0 \leq q < p - 1$ and $0 < \gamma \leq \infty$. Set*

$$s = \frac{n(p-1+\gamma)}{n(p-1-q) + p(\gamma+q)} \quad \text{and} \quad t = \frac{p-1+\gamma}{p-1-q}.$$

Let $\sigma = \theta dx$, where $\theta \neq 0$ is a nonnegative function in $L^{s,t}(\Omega)$. Then there exists a minimal positive (p, w) -superharmonic solution u to (1.1) such that $\min\{u, l\} \in W_0^{1,p}(\Omega)$ for all $l > 0$. Moreover, $u \in L^{r,\rho}(\Omega)$ and

$$(6.2) \quad \|u\|_{L^{r,\rho}(\Omega)} \leq C \|\theta\|_{L^{s,t}(\Omega)}^{\frac{1}{p-1-q}},$$

where

$$r = \frac{n(p-1+\gamma)}{n-p}, \quad \rho = p-1+\gamma$$

and C is a positive constant depending only on n, p, q and γ . Assume also that $\gamma \geq 1$. Then u belongs to $W_0^{1,p}(\Omega)$ and satisfies (1.1) in the sense of weak solutions.

Proof. As in the proof of [25, Corollary 3.6], using (6.1) and the Havin-Maz'ya potential estimate (see [26]), we find that

$$\int_{\Omega} (\mathcal{W}_{p,1}(\theta dx))^{\frac{(\gamma+q)(p-1)}{p-1-q}} \theta dx \leq C \|\theta\|_{L^{s,t}(\Omega)}^{\frac{p-1+\gamma}{p-1-q}}.$$

By Theorem 1.1, there exists a minimal p -superharmonic solution u such that

$$(6.3) \quad \mathcal{E}_\gamma(u^q \sigma) = \|u\|_{L^{\gamma+q}(\sigma)}^{\gamma+q} \leq C \|\theta\|_{L^{s,t}(\Omega)}^{\frac{p-1+\gamma}{p-1-q}}.$$

Furthermore, by Corollary 4.4 and a sharp form of Sobolev inequality (see, e.g., [34, p.234]),

$$(6.4) \quad \|u\|_{L^{r,\rho}(\Omega)}^{p-1+\gamma} = \|v\|_{L^{p^*,p}(\Omega)}^p \leq C \int_{\Omega} |\nabla v|^p dx \leq C \mathcal{E}_{\gamma}(u^q \sigma),$$

where $v = u^{\frac{p-1+\gamma}{p}}$. Combining (6.3) and (6.4), we obtain (6.2). Since Ω is bounded, we have

$$\begin{aligned} \int_{\Omega} u^q \theta dx &\leq \left(\int_{\Omega} u^{\gamma+q} \theta dx \right)^{\frac{q}{\gamma+q}} \left(\int_{\Omega} \theta dx \right)^{\frac{\gamma}{\gamma+q}} \\ &\leq \left(\int_{\Omega} u^{\gamma+q} \theta dx \right)^{\frac{q}{\gamma+q}} \left(C \|\theta\|_{L^{s,t}(\Omega)} |\Omega|^{\frac{s-1}{s}} \right)^{\frac{\gamma}{\gamma+q}} < \infty. \end{aligned}$$

Thus, $u^q \sigma = u^q \theta dx \in S^0$. By Proposition 4.2, this implies that $\min\{u, l\} \in W_0^{1,p}(\Omega)$ for all $l > 0$. If $\gamma \geq 1$, then $u^q \sigma \in S^0 \cap S^{\gamma} \subset S^1$, and hence $u \in W_0^{1,p}(\Omega)$. \square

Remark 6.2. For $0 < \gamma < 1$, using an interpolation argument (see, e.g., [2, Lemma 4.2]), from (6.4) and (4.2), we can deduce a gradient estimate of u .

7. QUASILINEAR ODE WITH HARDY-TYPE COEFFICIENTS

Let us now consider the model ordinary differential equation

$$(7.1) \quad \begin{cases} -(w|u'|^{p-2}u')' = \theta u^q & \text{in } (-1, 1), \\ u(-1) = u(1) = 0, \end{cases}$$

where

$$\begin{aligned} w(x) &= (1 - |x|)^{\beta}, \quad \beta \in (-1, p-1), \\ \theta(x) &= (1 - |x|)^{-\alpha}, \quad \alpha \in \mathbb{R}, \end{aligned}$$

and $' = \frac{d}{dx}$. The function w can be regarded as a Muckenhoupt A_p -weight in \mathbb{R} . Therefore it is also p -admissible (see [27, Chapter 15] [6, Theorem 2]). Note that θ is not integrable on $(-1, 1)$ if $\alpha \geq 1$.

Corollary 7.1. *Let $1 < p < \infty$ and $0 \leq q < p-1$. Assume that $\alpha < p-\beta$. Then there exists a bounded minimal positive weak solution $u \in H_{\text{loc}}^{1,p}((-1, 1); w) \cap C([-1, 1])$ to (7.1). Moreover, there exists a positive finite energy weak solution $u \in H_0^{1,p}((-1, 1); w)$ to (7.1) if and only if*

$$(7.2) \quad \alpha < 1 + (1+q) \left(1 - \frac{1}{p}\right) \left(1 - \frac{\beta}{p-1}\right).$$

Proof. Let $v = (1 - |x|)^A$. Then

$$-(w|v'|^{p-2}v')' = c(p, \beta, A)(1 - |x|)^{(A-1)(p-1)+\beta-1} + 2A^{p-1}\delta_0$$

in the sense of distribution, where $c(p, \beta, A) = -A^{p-1}\{(A-1)(p-1) + \beta\}$ and δ_0 is the Dirac mass concentrated at 0. Hence taking $A \in (0, 1 - \frac{\beta}{p-1})$ such that $A \leq \frac{p-\alpha-\beta}{p-1-q}$ and choosing a large C , we can make $V(x) = C(1 - |x|)^A$ a bounded supersolution to (7.1). Then Theorem 5.4 gives a positive bounded

weak solution $u \in H_{\text{loc}}^{1,p}((-1,1); w)$ to (7.1). The Sobolev embedding theorem provides $u \in C(-1,1)$. Furthermore, u is continuous up to the boundary because $0 \leq u(x) \leq V(x)$ for all $x \in (-1,1)$.

According to the Hardy-type inequality in [34, Theorem 1.3.3], (7.2) is necessary and sufficient for the embedding

$$\left(\int_{-1}^1 |u|^{1+q} \theta \, dx \right)^{\frac{1}{1+q}} \leq C \left(\int_{-1}^1 |u'|^p w \, dx \right)^{\frac{1}{p}}, \quad \forall u \in C_c^\infty(-1,1).$$

Thus Theorems 1.2 and 1.3 give the desired assertion. \square

REFERENCES

- [1] W. Allegretto and Y. X. Huang. A Picone's identity for the p -Laplacian and applications. *Nonlinear Anal.*, 32(7):819–830, 1998.
- [2] P. Bénilan, L. Boccardo, T. Gallouët, R. Gariepy, M. Pierre, and J. L. Vázquez. An L^1 -theory of existence and uniqueness of solutions of nonlinear elliptic equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 22(2):241–273, 1995.
- [3] M.-F. Bidaut-Véron. Necessary conditions of existence for an elliptic equation with source term and measure data involving p -Laplacian. In *Proceedings of the 2001 Luminy Conference on Quasilinear Elliptic and Parabolic Equations and System*, volume 8 of *Electron. J. Differ. Equ. Conf.*, pages 23–34. Southwest Texas State Univ., San Marcos, TX, 2002.
- [4] M. F. Bidaut-Véron. Removable singularities and existence for a quasilinear equation with absorption or source term and measure data. *Adv. Nonlinear Stud.*, 3(1):25–63, 2003.
- [5] A. Björn and J. Björn. *Nonlinear potential theory on metric spaces*, volume 17 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2011.
- [6] J. Björn, S. Buckley, and S. Keith. Admissible measures in one dimension. *Proc. Amer. Math. Soc.*, 134(3):703–705, 2006.
- [7] L. Boccardo, T. Gallouët, and L. Orsina. Existence and uniqueness of entropy solutions for nonlinear elliptic equations with measure data. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 13(5):539–551, 1996.
- [8] L. Boccardo and L. Orsina. Sublinear equations in L^s . *Houston J. Math.*, 20(1):99–114, 1994.
- [9] L. Brasco and G. Franzina. Convexity properties of Dirichlet integrals and Picone-type inequalities. *Kodai Math. J.*, 37(3):769–799, 2014.
- [10] H. Brezis and S. Kamin. Sublinear elliptic equations in \mathbf{R}^n . *Manuscripta Math.*, 74(1):87–106, 1992.
- [11] H. Brezis and L. Oswald. Remarks on sublinear elliptic equations. *Nonlinear Anal.*, 10(1):55–64, 1986.
- [12] D. Cao and I. Verbitsky. Nonlinear elliptic equations and intrinsic potentials of Wolff type. *J. Funct. Anal.*, 272(1):112–165, 2017.
- [13] D. T. Cao and I. E. Verbitsky. Finite energy solutions of quasilinear elliptic equations with sub-natural growth terms. *Calc. Var. Partial Differential Equations*, 52(3-4):529–546, 2015.
- [14] D. T. Cao and I. E. Verbitsky. Pointwise estimates of Brezis-Kamin type for solutions of sublinear elliptic equations. *Nonlinear Anal.*, 146:1–19, 2016.
- [15] C. Cascante, J. M. Ortega, and I. E. Verbitsky. Trace inequalities of Sobolev type in the upper triangle case. *Proc. London Math. Soc. (3)*, 80(2):391–414, 2000.
- [16] G. Dal Maso. On the integral representation of certain local functionals. *Ricerche Mat.*, 32(1):85–113, 1983.
- [17] G. Dal Maso, F. Murat, L. Orsina, and A. Prignet. Renormalized solutions of elliptic equations with general measure data. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 28(4):741–808, 1999.
- [18] D. G. de Figueiredo. Positive solutions of semilinear elliptic problems. In *Differential equations (São Paulo, 1981)*, volume 957 of *Lecture Notes in Math.*, pages 34–87. Springer, Berlin-New York, 1982.
- [19] M. Fukushima, Y. Oshima, and M. Takeda. *Dirichlet forms and symmetric Markov processes*, volume 19 of *De Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, extended edition, 2011.
- [20] L. Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2008.

- [21] A. Grigor'yan and I. Verbitsky. Pointwise estimates of solutions to semilinear elliptic equations and inequalities. *J. Anal. Math.*, 137(2):559–601, 2019.
- [22] A. Grigor'yan and I. Verbitsky. Pointwise estimates of solutions to nonlinear equations for nonlocal operators. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 20(2):721–750, 2020.
- [23] A. A. Grigor'yan. The heat equation on noncompact Riemannian manifolds. *Mat. Sb.*, 182(1):55–87, 1991.
- [24] T. Hara. Wolff potential estimates for Cheeger p -harmonic functions. *Collect. Math.*, 69(3):407–426, 2018.
- [25] T. Hara and A. Seesanea. Existence of minimal solutions to quasilinear elliptic equations with several sub-natural growth terms. *Nonlinear Analysis*, 197, 2020. <https://doi.org/10.1016/j.na.2020.111847>.
- [26] L. I. Hedberg and T. H. Wolff. Thin sets in nonlinear potential theory. *Ann. Inst. Fourier (Grenoble)*, 33(4):161–187, 1983.
- [27] J. Heinonen, T. Kilpeläinen, and O. Martio. *Nonlinear potential theory of degenerate elliptic equations*. Dover Publications, Inc., Mineola, NY, 2006. Unabridged republication of the 1993 original.
- [28] T. Kilpeläinen, T. Kuusi, and A. Tuhola-Kujanpää. Superharmonic functions are locally renormalized solutions. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 28(6):775–795, 2011.
- [29] T. Kilpeläinen and J. Malý. Degenerate elliptic equations with measure data and nonlinear potentials. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 19(4):591–613, 1992.
- [30] T. Kilpeläinen and J. Malý. The Wiener test and potential estimates for quasilinear elliptic equations. *Acta Math.*, 172(1):137–161, 1994.
- [31] T. Kilpeläinen and X. Xu. On the uniqueness problem for quasilinear elliptic equations involving measures. *Rev. Mat. Iberoamericana*, 12(2):461–475, 1996.
- [32] M. A. Krasnosel'skiĭ. *Positive solutions of operator equations*. Translated from the Russian by Richard E. Flaherty; edited by Leo F. Boron. P. Noordhoff Ltd. Groningen, 1964.
- [33] V. G. Maz'ya and V. P. Havin. A nonlinear potential theory. *Uspehi Mat. Nauk*, 27(6):67–138, 1972.
- [34] V. Maz'ya. *Sobolev spaces with applications to elliptic partial differential equations*, volume 342 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, augmented edition, 2011.
- [35] V. Maz'ya and Y. Netrusov. Some counterexamples for the theory of Sobolev spaces on bad domains. *Potential Anal.*, 4(1):47–65, 1995.
- [36] P. Mikkonen. On the Wolff potential and quasilinear elliptic equations involving measures. *Ann. Acad. Sci. Fenn. Math. Diss.*, (104):71, 1996.
- [37] N. C. Phuc and I. E. Verbitsky. Quasilinear and Hessian equations of Lane-Emden type. *Ann. of Math. (2)*, 168(3):859–914, 2008.
- [38] S. Quinn and I. E. Verbitsky. Weighted norm inequalities of $(1, q)$ -type for integral and fractional maximal operators. In *Harmonic analysis, partial differential equations and applications*, Appl. Numer. Harmon. Anal., pages 217–238. Birkhäuser/Springer, Cham, 2017.
- [39] S. Quinn and I. E. Verbitsky. A sublinear version of Schur's lemma and elliptic PDE. *Anal. PDE*, 11(2):439–466, 2018.
- [40] L. Saloff-Coste. A note on Poincaré, Sobolev, and Harnack inequalities. *Internat. Math. Res. Notices*, (2):27–38, 1992.
- [41] A. Seesanea and I. E. Verbitsky. Solutions in Lebesgue spaces to nonlinear elliptic equations with subnatural growth terms. *Algebra i Analiz*, 31(3):216–238, 2019.
- [42] A. Seesanea and I. E. Verbitsky. Solutions to sublinear elliptic equations with finite generalized energy. *Calc. Var. Partial Differential Equations*, 58(1):Art. 6, 21, 2019.
- [43] M. L. Silverstein. *Symmetric Markov processes*. Lecture Notes in Mathematics, Vol. 426. Springer-Verlag, Berlin-New York, 1974.
- [44] N. S. Trudinger and X.-J. Wang. On the weak continuity of elliptic operators and applications to potential theory. *Amer. J. Math.*, 124(2):369–410, 2002.
- [45] I. E. Verbitsky. Nonlinear potentials and trace inequalities. In *The Maz'ya anniversary collection, Vol. 2 (Rostock, 1998)*, volume 110 of *Oper. Theory Adv. Appl.*, pages 323–343. Birkhäuser, Basel, 1999.
- [46] I. E. Verbitsky. Sublinear equations and Schur's test for integral operators. In *50 years with Hardy spaces*, volume 261 of *Oper. Theory Adv. Appl.*, pages 467–484. Birkhäuser/Springer, Cham, 2018.

- [47] I. E. Verbitsky. Wolf's inequality for intrinsic nonlinear potentials and quasilinear elliptic equations. *Nonlinear Analysis*, 194, 2020. <https://doi.org/10.1016/j.na.2019.04.015>.
- [48] L. Véron. *Local and global aspects of quasilinear degenerate elliptic equations*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2017. Quasilinear elliptic singular problems.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, KITA 8 NISHI 10 SAPPORO, HOKKAIDO
060-0810, JAPAN

Email address: `takanobu.hara.math@gmail.com`