



HOKKAIDO UNIVERSITY

Title	Embryonic development and effect of water temperature on hatching of <i>Lophius litulon</i>
Author(s)	Ishikawa, Tomoya; Nakaya, Mitsuhiro; Takatsu, Tetsuya
Citation	Environmental biology of fishes, 105(1), 77-86 https://doi.org/10.1007/s10641-021-01195-4
Issue Date	2022-01
Doc URL	https://hdl.handle.net/2115/87848
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: http://dx.doi.org/10.1007/s10641-021-01195-4
Type	journal article
File Information	210918 Ishikawa et al Egg dev.pdf



Embryonic development and effect of water temperature on hatching of *Lophius litulon*

Tomoya Ishikawa^{1,*}, Mitsuhiro Nakaya², Tetsuya Takatsu²

¹ Graduate School of Fisheries Science, Hokkaido University, 3-1-1, Minato-cho, Hakodate, Hokkaido, 041-8611, Japan.

² Faculty of Fisheries Sciences, Hokkaido University, 3-1-1, Minato-cho, Hakodate, Hokkaido, 041-8611, Japan.

* Corresponding author: Tel, (+81) 138-40-8821;

E-mail: glyptoperichthys@eis.hokudai.ac.jp ([T. Ishikawa](mailto:glyptoperichthys@eis.hokudai.ac.jp)).

Declarations

Funding: No funding was received for conducting this study.

Conflicts of interest/Competing interests: The authors have no relevant financial or non-financial interests to disclose.

Availability of data: The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Ethics approval: All procedures and experiments were carried out in accordance with the experimental guidelines set by Hokkaido University.

Code availability: Not applicable

Consent to participate: All of the authors have approved the contents of this paper and have agreed to the Environmental Biology of Fishes 's submission policies.

Consent for publication: All of the authors have agreed to the Environmental Biology of Fishes's article publishing agreement.

Abstract

Most Lophiiformes including *Lophius* species are considered to spawn egg masses called an “egg veil” or “egg ribbon” composed of numerous mucoid chambers. However, *Lophius* eggs have been rarely collected in the field and thus little is known about their biological features. Here we examined the embryonic development of *Lophius litulon* and also the effects of water temperature on the embryonic development of eggs from egg veils collected in the field. Embryonic development was observed from the late cleavage stage to hatching. Low rates of mortality and deformity of embryos and early larvae were observed at 13–21°C. The estimated appropriate water temperature range closely approximated the range of temperatures (12–20°C) at which egg veils of *L. litulon* have been found in the field. Embryonic development accelerated with increased water temperatures. These results will enable estimation of the origin of and passive movement of egg veils and subsequently larvae, and will help to reveal the spatiotemporal distribution of the spawning and nursery grounds of *L. litulon*.

Key words: early development; goosfish; hatching process; larva; incubation temperature

Introduction

Pietsch and Arnold (2020) summarized information about Lophiiformes eggs. Most Lophiiformes are considered to spawn eggs encapsulated within a balloon-shaped or ribbon-shaped mucoid mass. Egg masses are formed by numerous mucoid chambers including eggs with seawater (Rasquin 1958) (Fig. 1a). There are few descriptions of the embryonic development of lophiiform fishes. Only *Histrio histrio* (Mosher 1954; Rasquin 1958; Fujita and Uchida 1959) and *Antennarius striatus* (Rasquin 1958) within the suborder Antennarioidei are recorded from fertilized ova to hatching. Descriptions of the embryonic development of species in Lophiidei are fewer than Antennarioidei. Unfertilized ova and the development from the gastrula stage of *Lophius americanus* was described by Everly (2002). However, other studies have reported only snippets of information like collection records or records over a short period of development (e.g., *L. piscatorius*: Laurenson (2006), *L. litulon*: Oh and Kim (2015)).

Lophiiformes species particularly in the genus *Lophius* are important fishery resources in regions throughout the world. However, their recruitment as well as landings often fluctuate significantly year by year (*L. americanus*: FAO (2020a), *L. budegassa*: ICES (2020), *L. litulon*: Takeya (2017), *L. piscatorius*: Solmundsson et al. (2010); FAO (2020b); ICES (2020), *L. vomerinus*: FAO (2020c)). Slight differences in survival rates during the early life stages have a large effect on fish stock fluctuations (Houde 1987). Thus, the mechanism of survival during the early life stages needs to be detailed to clarify the factors determining their population dynamics. However, information on the biological features of their early life stage is not sufficient to understand the mechanism of survival.

Lophius litulon is a commercially important demersal fish that is distributed in eastern Asia (Yoneda et al. 1997; Yoneda et al. 2001). Some of the reproductive ecology of *L. litulon* has been described. For example, spawning migrations have been shown in several regions (Yoneda et al. 2002; Iwasaki

et al. 2010; Takeya et al. 2013). At SST (sea surface temperature) of 9.0–16.3°C, large numbers of pre-spawning individuals were caught in areas shallower than 60 m (Takeya 2017). They seem to have either a single (Yoneda et al. 2001; Takeya 2017) or double (Kim et al. 2020) period of spawning in a year. They spawn buoyant egg masses which are called an “egg veil” (Hoshino et al. 2006), adults have been suggested to rise to the surface layer to spawn. Egg veils have been reported around Japan from February to August (Mito 1963; Takeya et al. 2013). Oh and Kim (2015) described the morphological and molecular characterization of eggs preserved in ethanol solution using pre-hatching specimens. Although information about its reproduction is increasingly available in the literature, details of the embryonic development, embryonic period and appropriate water temperature range for eggs of *L. litulon* remain unclear.

Using *L. litulon* egg veils collected in the field, this study investigated (1) the embryonic development from early development stage eggs (2) the relationship between incubation temperature and rate of development (3) the appropriate temperature range determined from mortality and deformity rates of early larvae, to describe the morphology of eggs and determine the relationship between reproduction and water temperature in *L. litulon*. We then considered the embryonic features of the genus *Lophius* and the effect of temperature on their growth and survival during the early life stages in relation to recruitment success.

Materials and methods

Field samplings (egg collection)

Three egg veils were collected with surrounding water using a scoop net (50 cm diameter with 2 cm mesh and 60 cm long) and bucket (ca. 5 l) from below the sea surface in the coastal area (depth: 5–30 m) off Kazamaura, Aomori Prefecture (Fig. 1b, Fig. 2), on 17 June and 21 July 2020 (SST: 15–16°C on 17 June, 18.7–20.7°C on 21 July). A part of the egg veils (ca. 1 l volume) was kept in a plastic

bag (30 l volume) in seawater (ca. 10 l) with oxygen and then transported to Hokkaido University, Hakodate by truck and ship taking approximately for four hours.

Laboratory observation

Egg diameters from these egg veils were measured using a stereomicroscope (Table 1).

Two egg veils collected on 21 July were experimentally incubated at different temperatures for observation of embryonic development as follows: When we took out and observed these egg veils on 21 July, one egg veil was in the late cleavage stage (this egg veil was termed 1B), and the other egg veil was in the blastula stage (termed as 2B). The egg veils were cut with scissors into ca. 2–3 cm square sizes, containing approximately 100 eggs in each piece. Then, four pieces from each egg veil were kept in small-sized plastic tanks one each with 500 ml of the artificial seawater at 18–20°C (Tetra Marine Salt Pro, Spectrum Brands Japan Inc; Japan, salinity: 32–33) (total 8 tanks). The tanks were put in incubators set at 13, 15, 18 and 21°C to examine the difference in rate of development within the approximate water temperature range that egg veils of *L. litulon* have been collected in the field (12–14°C: Oh and Kim (2015); 14.3°C: Nakaya et al. (2017b); 15.9°C: Mito (1963); 18.2°C: Kim (1976); 19–20°C: Hoshino et al. (2006) and this study (16–21°C (Table 1)). Although, some eggs became detached from the pieces from the cut edge, they remained incubated in the tanks. A piece of the egg veils was carefully transferred to a petri dish, observations and photographs were taken of 20 to 30 individuals twice a day (7:00–8:00 and 15:00–16:00) using a stereomicroscope. Then, the piece of egg veils was returned to the experimental tanks and continued to be incubated. The experimental observations were conducted until 50% hatching (24–29 July). The process of embryonic development was divided roughly into 4 stages following Oozeki and Hirano (1985) and Kawabe (2005), and more precisely classified into 9 stages following the stages described for the closely related *Lophius americanus* as defined by Everly (2002). Everly (2002) characterized these 9 stages as follows.

111 Stage 8: Blastodisc comprised of cells 4–5 layers deep

112 Stage 9: Formation of germ ring and embryonic shield

113 Stage 10: Blastoderm advances to cover 33% of yolk

114 Stage 11: Blastoderm advances to cover 34% and 66% of yolk

115 Stage 12: Blastoderm advances to cover 67% and 90% of yolk

116 Stage 13: Blastoderm advances to cover 91% and 100% of yolk; blastopore closes

117 Stage 14: Caudal fin begins to develop, torsion of the caudal fin occurs at end of stage 14

118 Stage 15: Appearance of pectoral and pelvic fin buds

119 Stage 16: Flexure of brain begins at hatching

120 The appropriate temperature range for early development

121 Appropriate temperature range of early development of *L. litulon* was examined as follows: The

122 three egg veils collected on 17 June and 21 July were cut with scissors into ca. 2–3 cm squares

123 containing approximately 100 eggs in each piece. 10 or 12 pieces of egg veils containing only 1 or 0

124 dead eggs were selected from each egg veil. The selected pieces were maintained in a bucket with 1 l

125 of the artificial seawater (18–20°C) for 1 hour to allow peripheral eggs along the cut edge to detach

126 from the egg veils. One piece from each egg veil was incubated in a small-sized plastic tank with 500

127 ml of the artificial seawater. Two replicate tanks were set in temperature incubators for each of the

128 three egg veils (total 6 tanks for each temperature but 4 tanks at 9°C by absence of 1B). To cover the

129 range of water temperature that egg veils were collected in previous study ((12–14°C: Oh and Kim

130 (2015); 14.3°C: Nakaya et al. (2017b); 15.9°C: Mito (1963); 18.2°C: Kim (1976); 19–20°C: Hoshino

131 et al. (2006)), incubation temperatures were maintained at 9, 13, 15, 18, 21 and 24°C. The pieces of

132 egg veils were acclimated from 18–20°C to the adjacent temperature incubator every 1–2 hours so

133 that the experiment could be started within 12 hours. After all temperatures were adjusted gradually

(0.1–2.0°C per hour), the experiment started. Incubation water was renewed 50% every two days and with gentle aeration (10–80 ml/min). Twice a day (8:30–9:30 and 14:30–15:30), emerged larvae and dead individuals were collected and counted the number of them and then emerged larvae were preserved in 90% ethanol solution. Deformity rate was determined by observing specimens, using a stereomicroscope. Deformed individuals were characterized from the bending of the notochord against the abdominal cavity and the stunting of the notochord as well as the stunting of the jaw. Percentage of NDL (normal developing larvae i.e., non-deformed live larvae) to the total collected individuals was calculated.

Results

Morphology of eggs

The eggs were spherical or slightly oval spherical, the diameter range was 1.51 to 1.98 mm (Table 1). Each egg had a single colorless and transparent yolk. The perivitelline space was extremely narrow. Color of the oil globule is orange, and becomes nearly colorless by the time to hatch. Eggs mainly had a single oil globule, some had small double or triple globules (1B: 24.1% at Stage 9–11, 2B: 2.1% at Stage 9–10) that had fused during development. Melanophores appeared sparsely on the blastoderm from three quarters epiboly.

Embryonic development

Embryonic development of *L. litulon* was recorded as Cleavage stage, Blastula stage, Gastrula stage and Embryo stage, and from Stage 8 through to Stage 16 as defined by Everly (2002) (Fig. 3). Each stage was characterized as follows.

Cleavage Stage (Egg veil 1B was at this stage just before the experiment)

Stage 8

Small blastomeres, and marginal periblast between the blastoderm and the yolk were observed.

157 Blastula Stage (Egg veil 2B was at this stage just before the experiment)

158 Stage 9

159 The central part of the blastoderm became transparent. The blastoderm started to spread over the
160 yolk.

161 Gastrula stage (Period, 13°C: 28h; 21°C: 16h)

162 Stage 10

163 The blastoderm became mostly transparent. The embryonic shield was observed. Epiboly
164 advanced, and the blastoderm covered approximately 30% of the yolk.

165 Stage 11

166 The blastoderm became transparent completely. The length of the embryonic shield was 20% of
167 the circumference of the yolk. The blastoderm covered approximately 50% of the yolk.

168 Embryo stage (Period, 13°C: 100 h; 21°C: 40 h)

169 Stage 12

170 Optic vesicles and the notochord were observed. Melanophores appeared on the blastoderm. The
171 embryo extended to 30% of the circumference of the yolk. The blastoderm covered approximately
172 70% of the yolk.

173 Stage 13

174 The lens, somite and Kupffer's vesicle appeared. Oil globule was located at the tip of the tail. The
175 melanophores increased in size by aggregating. The embryo extended to 40% of the circumference
176 of the yolk. The blastoderm completely covered the yolk.

177 Stage 14

178 Heart was visible clearly and started beating. The yolk around the tip of the tail somewhat became
179 hollow. The embryo extended to 50% of the circumference of the yolk.

Stage 15

The coloration of the embryo became mostly black. Melanophores covered the head, abdominal region, a part of yolk behind the abdominal and the tip of the tail. The embryo greatly increased in size and extended to 60% of the circumference of the yolk.

Hatching

Stage 16

The oil globule located posterior of the yolk. The vent located behind the yolk. The mouth is still closed.

Effect of water temperature on incubation period and survival

Percentage of NDL at 9, 13, 15, 18, 21 and 24°C were $17 \pm 14.8\%$, $64 \pm 12.2\%$, $71 \pm 14.5\%$, $71 \pm 13.5\%$, $81 \pm 7.0\%$ and $33 \pm 14.8\%$ (mean $\pm$ standard deviation), respectively (Fig. 4). The relationship between incubation temperature and rate of NDL was explained by a convex upward quadratic curve approximation

$$y = -0.91x^2 + 31.74x - 196.19 \quad (r^2 = 0.660, p < 0.001, n = 34)$$

where the vertex located at 17.4°C, and straight lines connecting the mean rate of NDL between adjacent incubation temperatures. The curve and polyline showed that over 50% rate of NDL were between 11.6°C and 23.2°C, and 11.8°C and 23.0°C, respectively.

The higher water temperature, the shorter time to hatch at 13–21°C (Fig. 5). Incubation period ranged 2.8–7.1 days for Stage 8 to Stage 16 depending on water temperature. Time from stage 10 to 50% hatch at 13, 15, 18 and 21°C were 128 ± 11.3 hours, 100 ± 5.7 hours, $72 \pm 0.0\%$ and 56 ± 0.0 (mean $\pm$ standard deviation) hours, respectively.

Discussion

Embryonic development

The eggs from the three egg veils were confirmed as *L. litulon* from the descriptions of morphological features of eggs (Oh and Kim 2015). Embryonic development from late cleavage stage through to hatch in *L. litulon* were observed. Information on egg or embryonic development of two species of *Lophius* has been reported (*L. americanus*: Fahay (1983); Everly (2002), *L. piscatorius*: Laurenson (2006)). Observations in this study have shown that eggs of *L. litulon* resembled that of *L. americanus* more than that of *L. piscatorius*. The most remarkable differences between *L. piscatorius* and the other two species were the egg diameter and rate of development. Egg diameter of *L. litulon* in this study was 1.51–1.98 mm. In comparison, the egg diameter of *L. americanus* was described as 1.61–1.84 mm (Fahay 1983), whereas egg diameter of *L. piscatorius* as approximately 3 mm (Laurenson 2006), twice larger than that of the others. In this study, *L. litulon* took 56 hours from stage 10 to hatching at 21°C. In comparison, *L. americanus* took approximately 75 hours from stage 10 to hatching at 20°C (Everly 2002), whereas *L. piscatorius* took approximately 120 hours from stage 12 to stage 14 at 7°C (Laurenson 2006). Rate of embryonic development of *L. litulon* is as rapid as that of *L. americanus* at almost the same temperatures, but that of *L. piscatorius* is slower at lower temperature. The eggs of *L. piscatorius* seemed to be collected under 10°C (Laurenson 2006), and subsequently the embryonic period of *L. litulon* and *L. americanus* would be much shorter than that of *L. piscatorius* in the field. The embryonic development process of eggs in *Lophius* seem to be common with that observed in this study because the appearances of the eggs in the three species are quite similar, however, eggs in other *Lophius* species remain poorly known.

Period of egg from fertilization to late cleavage stage is still unknown in *L. litulon* because eggs just after being spawned are difficult to collect. The embryonic period of related species such as *Histrionicus* took 16.5 hours from fertilization to 30% epiboly, and 48.3 hours from fertilization to hatching at 26.8–27.4°C (Fujita and Uchida 1959). In other words, time from fertilization to 30% epiboly

accounted for 34% of the whole embryonic period at stable water temperature in *Histrio histrio*. Eggs in *L. litulon* could be estimated to take 3.5–8.1 days from fertilization to hatch at estimated appropriate temperature range 13–21°C extrapolating from rate of development in *Histrio histrio*.

Effect of temperature on embryo

Incubation temperatures affected the mortality and the deformity rates for embryos because the rate of NDL was higher than 50% between 11.6°C and 23.2°C or 11.8°C and 23.0°C. Deformity in larvae can worsen their growth and survival (Andrades et al. 1996; Cobcroft and Battaglene 2009) and thus appropriate water temperature range was demonstrated at 12 to 23°C. This result is supported by the suggestion that areas under 12°C may be inappropriate as nursery grounds (Nakaya et al. 2017b). The appropriate temperature range almost covered the water temperature range that egg veils have been collected in previous studies (12–14°C: Oh and Kim (2015); 14.3°C: Nakaya et al. (2017b); 15.9°C: Mito (1963); 18.2°C: Kim (1976); 19–20°C: Hoshino et al. (2006) and this study (16–21°C (Table 1)). *L. litulon* may spawn within the appropriate water temperature range for eggs and early larval stage (at 12 to 23°C).

In this study, the rate of embryonic development in *L. litulon* was accelerated with the increase in incubation temperature between 13 and 21°C. Such a result has been typically shown and generally recognized in teleosts (e.g., Kamler 2002; Morehead and Hart 2003; Kawabe 2005; Yang and Chen 2005; Peña et al. 2014; Hu et al. 2017; Nakaya et al. 2017a). The origin and destination of egg veil in the field could be estimated by the relationship between stage of egg and water temperature, passive transport of egg veil by ocean currents or waves are being conducted to this end. Variation in water temperature coming with the changes of years and seasons will change period to hatch and then affect transportation of eggs and larvae in *L. litulon*.

This study found that water temperature affects growth and survival of eggs and early larval stage

of *L. litulon*. Takeya (2017) described a positive correlation between PDO (Pacific Decadal Oscillation) and reproductive success of *L. litulon* off Aomori Prefecture including this study area, and suggested that a cold regime can enhance their abundance. Solmundsson et al. (2010) described that the abundance and distribution of *L. piscatorius* off Iceland increased rapidly in the decade concurrent with increasing seawater temperature. These tendencies imply that spatiotemporal changes of water temperature by climatic conditions could alter the survival rate in any of the life stages until recruitment. For example, eggs can be exposed to inappropriate temperature if steep SST gradient is formed around spawning area. Survival rate in each life stage of *Lophius* is unknown. However, variation in recruitment of fish is known to be determined by survival during the early life stage (Hjört 1914). Thus, variation in water temperature during the egg and early larval stage is considered to effect their passive transport or growth and then recruitment success in *Lophius* e.g., “Stage duration hypothesis” (Houde 1987), “Transport hypothesis” (North and Houde 2003; Suzuki et al. 2019) and “Growth-predation hypothesis” (Litvak and Legget 1992; Takasuka et al. 2003). Information that we gained in this study is important to clarify the spatiotemporal distributions of spawning and nursery ground of *L. litulon* and the effect of temperature on recruitment success in the field. However, further research is needed to examine the passive transport of egg veils and larvae like the previous study in *L. piscatorius* conducted by Hislop et al. (2001).

Acknowledgments

We are grateful to Dr. Y. Takeya (Aomori Prefectural government), Dr. K. Noro and Mr. R. Suzuki (Aomori Prefectural Industrial Technology Research Center Fisheries Institute) for their support and valuable suggestions. We thank to Mr. Y. Kindaichi (Kazamaura fisheries cooperative association), Mr. G. Komamine (Komamine Corporation) the staff of Kazamaura Village Government, Laboratory of

Marine Bioresources Ecology of Hokkaido University Faculty of Fisheries and crew of the T/S Ushio-
maru of Hokkaido University, Faculty of Fisheries for their kind assistance in collection of specimens.
This work was conducted under permits from Hokkaido University Manual for Implementing Animal
Experimentation.

References

- Andrades JA, Becerra J, Fernández-Llebrez P (1996) Skeletal deformities in larval, juvenile and adult
stages of cultured gilthead sea bream (*Sparus aurata* L.). Aquaculture 141: 1–11
[https://doi.org/10.1016/0044-8486\(95\)01226-5](https://doi.org/10.1016/0044-8486(95)01226-5)
- Cobcroft JM, Battaglene SC (2009) Jaw malformation in striped trumpeter *Latris lineata* larvae linked
to walling behavior and tank color. Aquaculture 289: 274–282
<https://doi.org/10.1016/j.aquaculture.2008.12.018>
- Everly AW (2002) Stage of development of the goosfish, *Lophius americanus*, and comments on the
phylogenetic significance of the development of the luring apparatus in Lophiiformes. Environ
Biol Fishes 64: 393–417 <https://doi.org/10.1023/A:1016104612917>
- Fahay MP (1983) Guide to the early stages of marine fishes occurring in the western North Atlantic
Ocean, Cape Hatteras to the southern Scotian shelf. J Northwest Atl Fish Sci 4: 1–423
<https://doi.org/10.2960/J.v4.a1>
- FAO Fisheries and Aquaculture Department (2020a) Species Fact Sheets - *Lophius americanus*
(Valenciennes, 1837). <http://www.fao.org/fishery/species/3380/en>. Accessed 3 December 2020
- FAO Fisheries and Aquaculture Department (2020b) Species Fact Sheets - *Lophius piscatorius*
(Linnaeus, 1758). <http://www.fao.org/fishery/species/3379/en>. Accessed 3 December 2020
- FAO Fisheries and Aquaculture Department (2020c) Species Fact Sheets - *Lophius vomerinus*

(Valenciennes, 1837). <http://www.fao.org/fishery/species/2572/en>. Accessed 3 December 2020

Fujita S, Uchida K (1959) Spawning habits and early development of a sargassum fish, *Pterophryne histrio* (Linné). Sci Bull Fac Agr, Kyushu Univ 17: 277–282 (in Japanese with English abstract) <https://doi.org/10.15017/21499>

Hislop JRG, Gallego A, Heath MR, Kennedy FM, Reeves SA, Wright PJ (2001) A synthesis of the early life history of the anglerfish, *Lophius piscatorius* (Linnaeus, 1758) in northern British waters. ICES J Mar Sci 58: 70–86 <https://doi.org/10.1006/jmsc.2000.0991>

Hjørt J (1914) Fluctuations in the great fisheries of northern Europe. Rapp P.-v Réun Cons perm.int Explor Mer 20: 1–228

Hoshino N, Ichige S, Suzuki M, Yamaguchi Y, Toyoshima S (2006) Breeding experiment of anglerfish, *Lophius litulon* and examination as an object species farming. Ibaraki Pref Fish Exp Rep 40: 11–28 (in Japanese with English abstract)

Houde ED (1987) Fish early life dynamics and recruitment variability. Am Fish Soc Symp 2: 17–29

Hu F, Pan L, Gao F, Jian Y, Wang X, Li L, Zhang S, Guo W (2017) Effect of temperature on incubation period and hatching success of fat greenling (*Hexagrammos otakii* Jordan & Starks) eggs. Aquacult Res 48: 361–365 <https://doi.org/10.1111/are.12853>

ICES (2020) Working Group for the Bay of Biscay and the Iberian Waters Ecoregion (WGBIE). ICES Scientific Reports: 2 pp 36–85

Iwasaki T, Yoshida T, Chiyokubo M, Sato M (2010) Anglerfish, *Lophius litulon* landed at Fukushima Prefecture. Bull Fukushima Pref Fish Exp Stat 15: 11–25 (in Japanese)

Kamler E (2002) Ontogeny of yolk-feeding fish: an ecological perspective. Rev Fish Biol Fish 12: 79–103 <https://doi.org/10.1023/A:1022603204337>

Kawabe K (2005) Embryonic development and effect of water temperature on hatching of the blacktip

318 grouper, *Epinephelus fasciatus*. Aquacult Sci 53: 333–342 (in Japanese with English abstract)
 319 <https://doi.org/10.11233/aquaculturesci.1953.53.333>
 320 Kim DG, Im YJ, Kim JN, Lee HW, Jin S, Choi YJ, Baek GW (2020) Maturity and spawning of the
 321 yellow goosfish *Lophius litulon* in the coastal waters of Korea. Korean J Fish Aquat Sci 53: 67–
 322 73 (in Korean with English abstract) <https://doi.org/10.5657/KFAS.2020.0067>
 323 Kim UY (1976) On the morphology of larval stage of *Lophius litulon*. Bull Korean Fish Soc 9: 273–
 324 280 (in Korean with English abstract)
 325 Laurenson CH (2006) A note on the development of the embryos of anglerfish *Lophius piscatorius*. J
 326 Fish Biol 68: 1287–1290 <https://doi.org/10.1111/j.0022-1112.2006.01001.x>
 327 Litvak MK, Leggett WC (1992) Age and size-selective predation on larval fishes: the bigger-is-better
 328 hypothesis revisited. Mar Ecol Prog Ser 81: 13–24 <https://doi.org/10.3354/meps081013>
 329 Morehead DT, Hart PR (2003) Effect of temperature on hatching success and size of striped trumpeter
 330 (*Latris lineata*) larvae. Aquaculture 220: 595–606 [https://doi.org/10.1016/S0044-8486\(02\)00636-](https://doi.org/10.1016/S0044-8486(02)00636-1)
 331 [1](https://doi.org/10.1016/S0044-8486(02)00636-1)
 332 Mosher C (1954) Observation on the spawning behavior and the early larval development of the
 333 sargassum fish, *Histrio histrio* (Linnaeus). Zoologica 39: 141–152 3pls
 334 Mito S (1963) Pelagic fish eggs from Japanese waters-X Gadida and Lophiida. Jpn J Ichthyol 11: 103–
 335 113 (in Japanese with English abstract) <https://doi.org/10.11369/jji1950.11.103>
 336 Nakaya M, Marannu S, Inagaki Y, Kajiwara K, Sato Y, Takatsu T (2017a) Relationship between
 337 temperature and embryonic period of Arabesque greenling *Pleurogrammus azonus*. Aquacult Sci
 338 65: 247–250 <https://doi.org/10.11233/aquaculturesci.65.247>
 339 Nakaya M, Takeya Y, Suzuki R, Noro K, Gao W, Takatsu T (2017b) The effect of temperature on the
 340 early development and starvation tolerance of yellow goosfish *Lophius litulon*. Aquacult Sci

341 65: 251–254 <https://doi.org/10.11233/aquaculturesci.65.251>
 342 North EW, Houde ED (2003) Linking ETM physics, zooplankton prey, and fish early-life histories to
 343 striped bass *Morone saxatilis* and white perch *M. americana* recruitment. Mar Ecol Prog Ser 260:
 344 219–236 <https://doi.org/10.3354/meps260219>
 345 Oh J, Kim S (2015) Morphological and molecular characterization of separated pelagic eggs from
 346 *Lophius litulon* (Lophiiformes; Lophiidae). J Fish Biol 86: 1887–1891
 347 <https://doi.org/10.1111/jfb.12701>
 348 Oozeki Y, Hirano R (1985) Effect of temperature changes on the development of eggs of the Japanese
 349 whiting *Sillago japonica* Temminck et Schlegel. Bull Jpn Soc Sci Fish 51: 557–572
 350 <https://doi.org/10.2331/suisan.51.557>
 351 Peña R, Dumas S, Zavala-Leal I, Contreras-Olguín M (2014) Effect of incubation temperature on the
 352 embryonic development and yolk-sac larvae of the Pacific red snapper *Lutjanus peru* (Nichols &
 353 Murphy, 1922). Aquacult Res 45: 519–527 <https://doi.org/10.1111/j.1365-2109.2012.03255.x>
 354 Pietsch TW, Arnold RJ (2020) Frogfishes biodiversity zoogeography and behavioral ecology. Johns
 355 Hopkins University Press, Baltimore
 356 Rasquin P (1958) Ovarian morphology and early embryology of the pediculate fishes *Antennarius* and
 357 *Histrio*. Bull Am Mus Nat Hist 114: 327–372
 358 Solmundsson J, Jonsson E, Bjornsson H (2010) Phase transition in recruitment and distribution of
 359 monkfish (*Lophius piscatorius*) in Icelandic waters. Mar Biol 157: 295–305
 360 <https://doi.org/10.1007/s00227-009-1317-8>
 361 Suzuki KW, Fuji T, Kasai A, Itoh S, Kimura S, Yamashita Y (2019) Winter monsoon promotes the
 362 transport of Japanese temperate bass *Lateolabrax japonicus* eggs and larvae toward the innermost
 363 part of Tango Bay, the Sea of Japan. Fish Oceanogr 29: 66–83 <https://doi.org/10.1111/fog.12450>

364 Takasuka A, Aoki I, Mitani I (2003) Evidence of growth-selective predation on larval Japanese
 365 anchovy *Engraulis japonicus* in Sagami Bay. Mar Ecol Prog Ser 252: 223–238
 366 <https://doi.org/10.3354/meps252223>

367 Takeya Y, Nara K, Kosaka Y (2013) Behavioral analysis of the yellow goosfish *Lophius litulon* using
 368 bio-logging techniques. J Fish Tech 6: 1–15 (in Japanese with English abstract)

369 Takeya Y (2017) Ecological study and resource assessment of yellow goosfish *Lophius litulon* off
 370 Aomori Prefecture, northern Japan. Dissertation, Hokkaido University (in Japanese)

371 Yang Z, Chen Y (2005) Effect of temperature on incubation period and hatching success of obscure
 372 puffer *Takifugu obscurus* (Abe) eggs. Aquaculture 246: 173–179
 373 <https://doi.org/10.1016/j.aquaculture.2004.12.030>

374 Yoneda M, Tokimura M, Fujita H, Takeshita N, Takeshita K, Matsuyama M, Matsuura S (1997) Age
 375 and growth of the anglerfish *Lophius litulon* in the East China and the Yellow seas. Fish Sci 63:
 376 887–892 <https://doi.org/10.2331/fishsci.63.887>

377 Yoneda M, Tokimura M, Fujita H, Takeshita N, Takeshita K, Matsuyama M, Matsuura S (2001)
 378 Reproductive cycle, fecundity, and seasonal distribution of the anglerfish *Lophius litulon* in the
 379 East China and Yellow seas. Fish Bull 99: 356–370

380 Yoneda M, Tokimura M, Horikawa H, Yamamoto K, Matsuyama M, Matsuura S (2002) Spawning
 381 migration of the anglerfish *Lophius litulon* in the East China and Yellow Seas. Fish Sci 68: 310–
 382 313 https://doi.org/10.2331/fishsci.68.sup1_310

Fig. 1 Yellow goosfish *Lophius litulon*. a: egg veil chambers and ova, b: egg veil floating in the sea

Fig. 2 Location of the sampling site off Kazamaura

Fig. 3 Embryonic development of yellow goosfish, from late cleavage stage to newly hatched larva. Stages used are as defined by Everly (2002). V or L with stage number indicates ventral or lateral view

Fig. 4 Relationship between incubation temperature and rate of normal developing larvae (NDL) in yellow goosfish. The relationship is explained by a quadratic curve approximation (solid line) and a line graph (broken line). The dashed line indicates the 50% rate of normal developing larvae

Fig. 5 Relationship between mean water temperature and mean time from Stage 10 (Everly 2002) to 50% hatch in *Lophius litulon* eggs. Closed circle and cross mark indicate different egg veils, respectively

Table. 1. Summary of egg diameter in yellow goosfish collected around Tsugaru Strait

	Date collected	SST ^{*1} (°C)	Stage ^{*2}	<i>n</i>	Egg diameter (mm)	
					Mean ± s.d. ^{*3}	Range
Egg veil A	17 June 2020	16–18	14	30	1.8 ± 0.06	1.73–1.98
Egg veil B	21 July 2020	19–21	14	30	1.6 ± 0.07	1.51–1.80
Egg veil C	21 July 2020	19–21	15	30	1.8 ± 0.07	1.62–1.88

*1 SST is sea surface temperature at the sampling.

*2 Stages used are as defined Everly (2002). Stage is at the measurement.

*3 s.d. indicates standard deviation.




Fig. 1 Ishikawa et al.




Fig. 2 Ishikawa et al.




Fig. 3 Ishikawa et al.




Fig. 4 Ishikawa et al.




Fig. 5 Ishikawa et al.