



Title	High infection rate of tick-borne protozoan and rickettsial pathogens of cattle in Malawi and the development of a multiplex PCR for Babesia and Theileria species identification
Author(s)	Chatanga, Elisha; Maganga, Emmanuel; Mohamed, Wessam Mohamed Ahmed et al.
Citation	Acta tropica, 231, 106413 https://doi.org/10.1016/j.actatropica.2022.106413
Issue Date	2022-07
Doc URL	https://hdl.handle.net/2115/90204
Rights	© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	Chatanga_et al_HUSCAP.pdf



High infection rate of tick-borne protozoan and rickettsial pathogens of cattle in Malawi and the development of a multiplex PCR for *Babesia* and *Theileria* species identification

Elisha Chatanga^{a,b}, Emmanuel Maganga^{c,†}, Wessam Mohamed Ahmed Mohamed^a, Shohei Ogata^a, Pandey Sadaula Gita^a, Abdelbaset Eweda Abdelbaset^{a,d}, Kyoko Hayashida^e, Chihiro Sugimoto^e, Ken Katakura^a, Nariaki Nonaka^a, Ryo Nakao^{a,*}

^a. Laboratory of Parasitology, Graduate School of Infectious Diseases, Faculty of Veterinary Medicine, Hokkaido University, Kita-18, Nishi-9, Sapporo, Hokkaido 060-0818, Japan

^b. Department of Veterinary Pathobiology, Lilongwe University of Agriculture and Natural Resources, P.O. Box 219, Lilongwe, Malawi

^c. Mikolongwe College of Veterinary Sciences, P.O. Box 5193, Limbe, Blantyre, Malawi

^d. Clinical Laboratory Diagnosis, Department of Animal Medicine, Faculty of Veterinary Medicine, Assiut University, Assiut 71515, Egypt

^e. Division of Collaboration and Education, International Institute for Zoonosis Control, Hokkaido University, Kita-20, Nishi-10, Sapporo, Hokkaido 001-0020, Japan

EC: chatanga@vetmed.hokudai.ac.jp

EM: magangamanuel@gmail.com

SO: s.ogata@vetmed.hokudai.ac.jp

WMAM: wessam@czc.hokudai.ac.jp

PSG: gitapandeyk@gmail.com

AEA: abdelbaset2006@hotmail.com

KH: kyouko-h@czc.hokudai.ac.jp

CS: sugimoto@czc.hokudai.ac.jp

KK: kenkata@vetmed.hokudai.ac.jp

NN: nnonaka@vetmed.hokudai.ac.jp

RN: ryo.nakao@vetmed.hokudai.ac.jp

† Deceased

* Correspondence: ryo.nakao@vetmed.hokudai.ac.jp

Abstract

Malawi has an estimated cattle population of 1,884,803 heads, the indigenous Malawi zebu breed accounts for 91.2%, while the exotic and crossbred accounts for the remaining 8.8%. Although ticks and tick-borne diseases are widespread in Malawi, no molecular study has been conducted to investigate the tick-borne Anaplasmataceae and piroplasms infecting cattle. To provide an insight into the current status of tick-borne pathogens (TBPs) of cattle, a molecular survey was conducted in the central and southern regions of Malawi. A total of 191 cattle of which 132 were Malawi zebu, 44 were Holstein Friesian and 15 were Holstein-Friesian/ Malawi zebu crosses were screened for Anaplasmataceae and piroplasms using the heat shock protein *groEL* and 18S rDNA, respectively. A new 18S rDNA multiplex PCR assay was designed for *Babesia* and *Theileria* species identification without sequencing. Overall, 92.3% ($n = 177$) of the examined animals were infected with at least one TBP. Anaplasmataceae-positive rate was 57.6% ($n = 110$) while for piroplasms it was 80.1% ($n = 153$). The detected Anaplasmataceae were *Anaplasma bovis* 2.6% ($n = 5$), *Anaplasma marginale* 24.6% ($n = 47$), *Anaplasma platys*-like 13.6% ($n = 26$), uncharacterized *Anaplasma* sp. 14.1% ($n = 27$), and uncharacterized *Ehrlichia* sp. 16.2% ($n = 31$). The detected piroplasms were *Babesia bigemina* 2.6% ($n = 5$), *Theileria mutans* 73.8% ($n = 141$), *Theileria parva* 33.0% ($n = 63$), *Theileria taurotragi* 12.6% ($n = 24$), and *Theileria velifera*

53.4% ($n = 102$). Mixed infection rate was 79.6% ($n = 152$) of the samples analyzed. This study has shown a high burden of TBPs among cattle in Malawi which highlights the need to conceive new methods to control ticks and TBPs in order to improve animal health and productivity. The newly developed multiplex PCR assay would be a useful tool especially in resource limited settings where sequencing is not available and when mixed infections are expected.

Keywords: Anaplasmataceae, cattle, Malawi, molecular epidemiology, piroplasms.

1. Introduction

Tick-borne pathogens (TBPs) that belong to the genera *Anaplasma*, *Babesia*, *Ehrlichia* and *Theileria* are widespread in sub-Saharan Africa where the vector ticks are present. In Malawi, 15 tick species namely: *Amblyomma variegatum*, *Hyalomma marginatum rufipes*, *Hyalomma truncatum*, *Rhipicephalus appendiculatus*, *Rhipicephalus compositus*, *Rhipicephalus decoloratus*, *Rhipicephalus kochi*, *Rhipicephalus masseyi*, *Rhipicephalus microplus*, *Rhipicephalus planus*, *Rhipicephalus pravus*, *Rhipicephalus punctatus*, *Rhipicephalus sanguineus sensu lato*, *Rhipicephalus simus* and *Rhipicephalus tricuspis*, have been reported to infest cattle (Walker et al., 2003; Berggren, 1978).

Bovine anaplasmosis (gall sickness) is caused by *Anaplasma marginale* infection, and the clinical outcome is characterized by anemia, icterus, fever, weight loss, abortion, and lethargy and in severe cases, it may be fatal (Kocan et al., 2003). The occurrence of clinical disease is dependent on the age of the animal, as susceptibility increases with age. Calves under 6 months of age are generally less susceptible (Kocan et al., 2003; Richey, 1991). In Malawi, cases of bovine anaplasmosis have been reported but its causative agents have not yet been characterized. *Anaplasma marginale* is transmitted by the vector ticks such as *A.*

74 *variegatum*, *R. decoloratus*, *R. evertsi evertsi* and *R. microplus* (Walker et al., 2003). Cattle
75 may also be infected with *Anaplasma centrale*, and *Anaplasma bovis* formerly known as
76 *Ehrlichia bovis* (Dumler et al., 2001), which cause benign anaplasmosis. *Anaplasma platys*-
77 like infection in cattle has also been reported in Bolivia (Ogata et al., 2021), Kenya (Peters et
78 al., 2020), and Tunisia (Ben Said et al., 2017) however, its clinical impact on cattle has not
79 yet been fully explored.

80 Bovine ehrlichiosis (heartwater) is caused by *Ehrlichia ruminantium* formerly known
81 as *Cowdria ruminantium*. The clinical disease is commonly associated with endocarditis in
82 cattle and other non-specific signs such as fever, anorexia, aggressiveness and sudden death
83 (Allsopp, 2015). The vector ticks of *E. ruminantium* in southern Africa are *Amblyomma*
84 *hebraeum* and *A. variegatum* (Walker et al., 2003). Other uncharacterized *Ehrlichia* species
85 closely related to *Ehrlichia minasensis* but separated from the pathogenic *E. ruminantium*
86 have been reported to infect cattle in Kenya, in apparently healthy dairy cattle (Peters et al.,
87 2019, 2020). In Malawi, *E. ruminantium* infection in goats has been reported using molecular
88 techniques (Chatanga et al., 2021a).

89 Bovine babesiosis (red water) which is caused by *Babesia bovis* and *Babesia*
90 *bigemina* has also been documented in Malawi (Lawrence et al., 1996). The newly described
91 pathogenic *Babesia* sp. Mymegnsingh has also been documented to cause clinical disease in
92 cattle in Argentina, Philippines, Sri Lanka, Uganda and Vietnam (Sivakumar et al., 2018,
93 2020). The clinical disease is characterized by high fever, hemoglobinuria, dark colored
94 urine, anorexia, lethargy and neurological signs which usually result in fatal outcome
95 (Uilenberg et al., 1995). The vector ticks for *B. bigemina* and *B. bovis* in southern Africa are
96 *R. microplus*, *R. decoloratus*, and *R. evertsi evertsi* (Walker et al., 2003).

East Coast fever (ECF) (one of bovine theilerioses) is caused by *Theileria parva* and is considered the most economically important tick-borne diseases in Africa (Lubembe et al., 2020). Calves of indigenous zebu cattle below the age of 6 months are highly susceptible to the infection (Moll et al., 1984, 1986). The disease is severe in exotic and crossbred cattle with morbidity and mortality rates ranging from 80-100% (Oura et al., 2007). The clinical disease is characterized by anemia, icterus, tachypnea, tachycardia, lethargy, froth coming out of the mouth, anorexia, fever, abortion, and enlarged superficial lymph nodes (Nene et al., 2015). *Theileria parva* is transmitted by the brown ear tick, *R. appendiculatus* (Walker et al., 2003). Other *Theileria* species that infect cattle but cause benign theileriosis in southern Africa are *Theileria mutans*, *Theileria taurotragi*, and *Theileria velifera*.

Malawi cattle population is estimated at 1,884,803 heads of which the indigenous Malawi zebu accounts for 91.2% (1,719,641) while the exotic and crossbred cattle account for the remaining 8.8% (165,862) as of 2020 (DAHLD, 2020). With an estimated human population of 17,563,749 as of 2018 (NSO, 2018), Malawi has one of the lowest cattle population per capita in Africa. Approximately 71% of the total pure and crossbred dairy cattle in Malawi are found in the southern region which is considered to be non-endemic to ECF (DAHLD, 2006). The endemic status of ECF has hampered the establishment of the dairy industry in the central and northern regions of the country (Lawrence et al., 1996; Chinombo et al., 1988).

This study aimed to provide molecular epidemiological data on the current prevalence and characterization of TBPs infecting cattle in Malawi. Cattle blood samples were screened for Anaplasmatidae and piroplasms using conventional PCR and a newly developed multiplex PCR assay. This epidemiological data is a pre-requisite for the development of novel control measures against tick and TBPs in Malawi to achieve the national goal of Malawi being self-sufficient in safe locally produced livestock and livestock products.

122

123 **2. Material and methods**

124 **2.1 Ethical consideration.**

125 This was a parallel study of our previous study (Chatanga et al., 2020) which was
126 approved by Ministry of Agriculture, Irrigation and Water Development (MoAIWD) in
127 Malawi through the Department of Animal Health and Livestock Development (DAHLD)
128 reference number 10/15/32/D and permission for sampling was obtained from the host
129 institutions and farmers.

130 **2.2 Study site and sample collection**

131 Cattle blood samples ($n = 191$) were collected from five districts in the central region
132 and southern regions of Malawi (Figure 1) from February 2018 to March 2019 during the rainy
133 seasons which runs from November to April from apparently healthy animals. Specifically, the
134 samples were collected from Kasungu ($n = 26$), Nkhotakota ($n = 37$), Katete ($n = 33$), Likasi
135 ($n = 34$), Lilongwe University of Agriculture and Natural Resources (LUANAR) ($n = 33$) and
136 Mikolongwe ($n = 28$). Overall, the sampled animals were Malawi zebu ($n = 132$), Holstein
137 Friesian ($n = 44$) and Holstein Friesian/ Malawi zebu crosses ($n = 15$).

138 The animals investigated in Kasungu and Nkhotakota were Malawi Zebu cattle that
139 were managed under extensive grazing in communal grazing lands. In this management system,
140 animals from different smallholder farms were mixed freely and there was no ECF vaccination
141 or dipping history to control ticks. The animals investigated at Katete farm in Lilongwe were
142 Holstein-Friesian which were kept under a semi-intensive management with no contact with
143 other herds. They were dipped weekly during rainy season and fortnightly during dry season
144 to control ticks and Muguga cocktail (MC) vaccine was also used to control ECF in some
145 animals. A total of 20 of the sampled animals at Katete farm were vaccinated with Muguga
146 cocktail vaccine 2 years and 3 months prior to sampling period while the remainder ($n = 13$)

were unvaccinated but these animals had been co-grazing for more than 2 years. At Likasi, LUANAR and Mikolongwe farms both exotic dairy breed Holstein Friesian and local Malawi zebu were kept. The animals were dipped fortnightly and MC vaccine was used to immunize the exotic breeds but not the local Malawi zebu. Approximately 5 ml of whole blood was collected by venipuncture of the external jugular vein after disinfection of the puncture site with methylated spirit cotton swab into ethylene diamine tetraacetic acid (EDTA) vacutainer tube.



Figure 1. Map of Malawi showing the sample collection sites.

2.3 DNA extraction

DNA was extracted from 200 µl of whole blood using the Quick Gene DNA whole blood kit S (DB-S) (Kurabo Industries Ltd., Osaka, Japan) according to the manufacturer's recommendations. The extracted DNA was stored at -20°C until required for use.

2.4 Polymerase chain reaction

All the 191 samples from six sampling sites were screened for *Anaplasma*, *Ehrlichia*, *Babesia* and *Theileria* species. The screening of Anaplasmataceae was done targeting the heat shock protein gene (*groEL*) nested PCR assays as previously described (Rar et al., 2011; Liz et al., 2000; Gofton et al., 2016). *Anaplasma ovis* DNA extracted from positive sheep blood was used as positive control while molecular grade water instead of genomic DNA was used as negative control for quality control.

The *groEL* gene secondary PCR assays make use of two primer sets: one that is specific for the genus *Anaplasma* and another that is specific for the genus *Ehrlichia*. This allows detection of mixed infection of *Anaplasma* and *Ehrlichia* spp. in a sample. Further, characterization was done using the citrate synthase gene (*gltA*) nested PCR assays as previously described (Inokuma et al., 2005). The PCR conditions for both 1st and 2nd reactions were set at initial denaturation at 94°C for 1min, followed by 35 cycles of denaturation at 98°C for 10 sec, annealing at 50/54/55°C for 15sec and extension at 68°C for 1 min and final extension at 68°C for 5min.

The screening for piroplasms was done targeting almost the full length of the *18S ribosomal RNA gene* (rDNA) nested PCR assays targeting *Babesia*, *Theileria* and *Hepatozoon* (BTH) species as described by Masatani et al. (2017), and Qiu et al. (2018). All PCR reactions were conducted in a 25 µl reaction mixture containing 0.5 µl of Tks Gflex DNA Polymerase

(1.25 units/ μ l) (TaKaRa Bio Inc., Shiga, Japan), 12.5 μ l of 2 $\times$ Gflex PCR Buffer (Mg^{2+} , dNTP plus), 200 nM of each primer, 1.0 μ l of template DNA and molecular grade water. *Theileria orientalis* Ikeda strain DNA extracted from cattle blood was used as positive control while molecular grade water instead of genomic DNA was used as negative control for quality control. Due to mixed signals observed in selected sequenced 2nd BTH PCR products a new item multiplex PCR assay for *B. bigemina*, *B. bovis*, *T. mutans*, *T. parva*, *T. taurotragi*, and *T. velifera* was designed in this study to identify the detected pathogens to species level (Figure 2A).

To design the new multiplex PCR primers, sequences of the 18S rDNA of *Babesia* species (*Babesia bovis*, AY150059, HQ264127; and *Babesia bigemina*, KM046917, JQ437264), *Theileria* species (*Theileria mutans*, AF078815; *Theileria parva*, MG952923, AF013418; *Theileria taurotragi*, L19082 and *Theileria velifera*, AF097993) deposited in the GenBank were downloaded. The sequences were aligned using ClustalW in Molecular Evolutionary Genetics Analysis (MEGA) version 7 (Kumar et al., 2016) and were visually checked for regions that have homologies within each species. Primers were designed by using Primer 3 plus (<https://primer3plus.com/cgi-bin/dev/primer3plus.cgi>). The specificity of the designed primers was confirmed by additional BLAST searches (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The 100-fold diluted first BTH PCR product was used as a template for the *Babesia/ Theileria* multiplex PCR assay. The multiplex PCR was used to identify the detected pathogens based on the 2nd BTH PCR results to species level. The multiplex PCR primers were used for sequencing of the detected pathogens to generate sequences for phylogenetic tree analysis. The primers, their annealing temperature and the expected amplicon size are listed in Table 1.




Figure 2. Representative gel electrophoresis images of the multiplex PCR products.

A: M = 50 bp marker; B.bi = *Babesia bigemina*, B.bo = *Babesia bovis*, T.mu = *Theileria mutans*, T.pa = *Theileria parva*, T.ta = *Theileria tautotragi*, T.ve = *Theileria velifera*, NC = negative control.

B: M = 50 bp marker; lanes 1-13 samples, NC = negative control.

210 Table 1. List of primers used for the detection and characterization of tick-borne Anaplasmataceae and piroplasms of cattle in Malawi.

Primer name	Primer sequence (5' to 3')	Target gene/ organism	PCR type	Amplicon size (bp)	Annealing temperature (°C)	Reference
BTH 1st F	GTGAAACTGCGAATGGCTCATTAC	18S rDNA/ <i>Babesia</i> , <i>Theileria</i> and <i>Hepatozoon</i>	1st PCR	1400 - 1600	55	Masatani et al., 2017
BTH 1st R	AAGTGATAAGGTTACAAAACTTCCC					
BTH 2nd F	GGCTCATTACAACAGTTATAGTTTATTTG		2nd PCR	1400 - 1600		
BTH 2nd R	CGGTCCGAATAATTCACCGGAT					
BTH_Bbig_F	AGCCTTGGTAATGGTTAATAG	18S rDNA/ <i>Babesia</i> , and <i>Theileria</i>	Multiplex PCR	848	57	This study
BTH_Bbov_F3	TGTCCTTTCTTGATTCTCTGGGTAG			469		
BTH_Tmut_F	GGCCCTTGCCTTGAATACTTTAG			915		
BTH_Tpar_F	CTCTGCATGTGGCTTATTTTCGG			995		
BTH_Ttau_F	TGCTAAATAGGGTACGGGAGC			330		
BTH_Tveli_F	AGGCTTTTGCCGTCCCGTGATCGC			310		
BTH_NEW_R	TCCGAATAATTCACCGGATCAC					
HS1-F	CGYCAGTGGGCTGGTAATGAA	<i>groEL</i> gene/ Anaplasmataceae	1st PCR	1300	54	Rar et al., 2011
HS6-R	CCWCCWGGTACWACACCTTC					
HS3-F	ATAGTYATGAAGGAGAGTGAT	<i>groEL</i> gene/ <i>Anaplasma</i>	2nd PCR	1256	50	Liz et al., 2000
HSV-R	TCAACAGCAGCTCTAGTWG					
groEL_fwd3	TGGCAAATGTAGTTGTAACAGG	<i>groEL</i> gene/ <i>Ehrlichia</i>	2nd PCR	1100	50	Gofton et al., 2016
groEL_rev2	GCCGACTTTTAGTACAGCAA					
F4b	CCAGGCTTTATGTCAACTGC	<i>gltA</i> gene/ Anaplasmataceae	1st PCR	800	55	Inokuma et al., 2005
R1b	CGATGACCAAAAACCCAT					
EHR-CS136F	TTYATGTCYACTGCTGCKTG	<i>gltA</i> gene/ <i>Anaplasma</i> and <i>Ehrlichia</i>	2nd PCR	650	55	Inokuma et al., 2005
EHR-CS778R	GCNCCMCCATGMGCTGG					

211 PCR = Polymerase chain reaction; F = Forward; R = Reverse

2.5 Sequencing

All samples that were positive on 2nd *groEL* gene PCR ($n = 110$) were purified using ExoSAP-IT™ PCR Product Clean-up Reagent (Applied Biosystems, CA, USA) and sequenced. Thereafter, based on the species identified from the obtained *groEL* genes sequences, i.e. 4, 3, 1, 1 and 7 for *A. platys*, *A. marginale*, *A. bovis*, uncharacterised *Anaplasma* sp., and uncharacterised *Ehrlichia* sp., respectively, we selected 5 samples per *Anaplasma* or *Ehrlichia* species of the 2nd *gltA* gene PCR for sequencing. For *Babesia* and *Theileria* species, five randomly selected samples per species from the designed new item (multiplex PCR) products were cut from the gel and purified using NucleoSpin Gel and PCR Clean-Up Kit (Takara Bio Inc.). Sequencing was done in both directions using BigDye™ Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, CA, USA) and ABI genetic analyzer 3500xl (Applied Biosystems, CA, USA). Sequence data editing was conducted using ATGC software (GENETYX Corporation, Tokyo, JP) by trimming the primers annealing sites and the consensus sequence was extracted for phylogenetic analysis. The sequences generated in this study were submitted to the DNA Data Bank of Japan (DDBJ) (<http://www.ddbj.nig.ac.jp/>) under the accession numbers LC664051 to LC664059 for piroplasms 18S rDNA, LC664060 to LC664073 and LC671361 for Anaplasmataceae *gltA* gene and LC664074 to LC664088 and LC671362 for Anaplasmataceae *groEL* gene.

2.6 Statistical analysis

The significance of co-infections was examined by statistical modelling using generalized linear model (Glm) function in R version 4.1.2. Thereafter, the odds ratio were calculated based on modelling results using odds ration function in R which were eventually used for calculation of the correlation of co-infection using kendall method cor function in stats package in R software (<http://www.rstudio.com/>). The “Holm” method was used for

adjusting the p-value over multiple testing (<https://www.jstor.org/stable/2346101>). The $p < 0.05$ was considered significant.

2.7 Phylogenetic analysis

Alignments of the consensus nucleotide sequences generated from the amplified DNA fragments were done using ClustalW in MEGA version 7. To understand the relationship of the sequences obtained in this study and those deposited in the GenBank, maximum likelihood (ML) phylogenetic trees were constructed using MEGA version 7 software using the Tamura 2 parameter model.

3. Results

3.1 PCR screening and sequence analysis

The screening for Anaplasmataceae using *groEL* gene nested PCR assay showed that 110/191 (57.6%) were positive. To identify the detected *Anaplasma* and *Ehrlichia* species sequencing of all 2nd *groEL* gene PCR positive samples ($n = 110$) was done. The Anaplasmataceae sequence data showed that the examined cattle in Malawi were infected with *Anaplasma bovis* 5/191 (2.6%), *Anaplasma marginale* 47/191 (24.6%), *Anaplasma platys*-like 26/191 (13.6%), uncharacterized *Anaplasma* sp. 27/191 (14.1%) and uncharacterized *Ehrlichia* sp. 31/191 (16.2%) as shown in Table 2.

The sequences of all the uncharacterised *Anaplasma* sp. ($n = 27$) were identical and closely related to *A. bovis* in both the *groEL* and *gltA* genes. However, the sequences of the uncharacterised *Ehrlichia* sp. ($n = 31$) produced 7 and 10 sequences for the *groEL* and *gltA* genes, respectively. Since, we did not expect any wider variation between the results of *groEL* gene and *gltA* gene, five (5) for each of the above detected pathogens were also sequenced for *gltA* gene for further characterization.

260 The screening for *Babesia* and *Theileria* species using 18S rDNA gene BTH nested
261 PCR assay showed that 153/191 (80.1%) were positive. Further screening of the 2nd BTH
262 PCR amplicons using the new multiplex PCR assay showed that the examined animals were
263 infected with *B. bigemina* 5/191 (2.6%), *T. mutans* 141/ 191 (73.8%), *T. parva* 63/191
264 (33.0%), *T. taurotragi* 24/191 (12.6%), and *T. velifera* 102/191 (53.4%), as shown in Table 2.
265 The samples from Katete showed that among the 20 samples from vaccinated cattle 14 were
266 positive for *T. parva* representing 70% positive rate, while among the 13 unvaccinated
267 samples 3 were positive for *T. parva* representing 23% positive rate.

268 Table 2. Positive detection rates of tick-borne pathogens (TBPs) in the studied cattle in Malawi.

Pathogen	Study site						Overall <i>n</i> = 191
	Kasungu	Nkhotakota	Katete	Likasi	LUANAR	Mikolongwe	
	<i>n</i> = 26	<i>n</i> = 37	<i>n</i> = 33	<i>n</i> = 34	<i>n</i> = 33	<i>n</i> = 28	
<i>A. bovis</i>	0	0	0	0	2 (6%)	3 (11%)	5 (2.9%)
<i>A. marginale</i>	5 (19%)	6 (16%)	7 (21%)	11 (32%)	6 (18%)	12 (43%)	47 (24.6%)
<i>A. platys</i>	2 (8%)	13 (35%)	4 (12%)	6 (18%)	0	1 (4%)	26 (13.6%)
<i>Anaplasma</i> sp.	7 (27%)	4 (11%)	5 (15%)	4 (11%)	3 (9%)	4 (14%)	27 (14.1%)
<i>Ehrlichia</i> sp.	12 (46%)	5 (14%)	2 (6%)	6 (17%)	1 (3%)	5 (18%)	31 (16.2%)
<i>B. bigemina</i>	1 (4%)	2 (5%)	1 (3%)	2 (6%)	0	0	6 (3.1%)
<i>B. bovis</i>	0	0	0	0	0	0	0
<i>T. mutans</i>	19 (73%)	28 (77%)	22 (67%)	26 (76%)	28 (85%)	18 (64%)	141 (73.8%)
<i>T. parva</i>	9 (35%)	10 (27%)	17 (52%)	9 (26%)	11 (33%)	7 (25%)	63 (33.0%)
<i>T. taurotragi</i>	4 (15%)	2 (5%)	6 (18%)	3 (9%)	3 (9%)	6 (21%)	24 (12.6%)
<i>T. velifera</i>	14 (54%)	19 (51%)	13 (39%)	19 (56%)	25 (76%)	12 (43%)	102 (53.4%)

269 *n* = number of samples, number of samples positive and percentage is provided in parenthesis.

270

271

3.2 Infection status of TBPs in cattle

Overall, 177 cattle were infected with at least one TBP, representing 92.7% positive rate. Co-infections were observed in 152 cattle (79.6%), and the number of pathogens co-infecting a sample ranged from two to six (Table 3). Triple co-infections had the highest frequency at 39.3% ($n = 75$), followed by double co-infections at 28.3% ($n = 54$), as shown in Table 3. Sextuple co-infection had the least frequency at 0.5% ($n = 1$) (Table 3). LUANAR farm had the highest co-infection rate at 91%, followed by Likasi, Mikolongwe, Nkhotakota, Kasungu and Katete, at 85%, 79%, 78%, 77% and 67%, respectively (Table 3). The double infections combination with the highest frequency was *T. mutans* + *T. velifera* at 12.0% ($n = 23$). In triple infections, the highest frequency was observed in the combination *T. mutans* + *T. parva* + *T. velifera* at 7.9% ($n = 15$). For quadruple infections, the highest frequency was observed in the combination *A. marginale* + *T. mutans* + *T. parva* + *T. velifera* at 2.6% ($n = 5$).

285 Table 3. Number of TBPs co-infections observed in the studied cattle in Malawi.

Number of pathogens in co-infection	Study site						Overall <i>n</i> = 191
	Kasungu <i>n</i> = 26	Nkhotakota <i>n</i> = 37	Katete <i>n</i> = 33	Likasi <i>n</i> = 34	LUANAR <i>n</i> =33	Mikolongwe <i>n</i> = 28	
2	7 (27%)	14 (38%)	10 (30%)	8 (24%)	13 (39%)	2 (7%)	54 (28.3%)
3	12 (46%)	13 (35%)	7(21%)	17 (50%)	13 (39%)	13 (46%)	75 (39.3%)
4	1 (4%)	1 (3%)	5 (15%)	4 (12%)	1 (3%)	6 (21%)	18 (9.4%)
5	0	1 (3%)	0	0	2 (6%)	1 (4%)	4 (2.1%)
6	0	0	0	0	1 (3%)	0	1 (0.5%)
Total	20 (77%)	29 (78%)	22 (67%)	29 (85%)	30 (91%)	22 (79%)	152 (79.6%)

286 *n* = number of samples, number of samples positive and percentage is provided parenthesis.

287

288

289

290

291

292

293

The summary of the pathogen combinations per sampling site has been provided in Table S1. Statistical analysis showed both positive and negative correlations between the co-infecting pathogens examined in this study. Statistically significant correlation that was statistically significant was observed between *Anaplasma* species infection with *A. bovis* ($p = < 0.001$) and *A. platys*-like ($p = 0.026$). This study also showed statistically significant correlation between co-infection with *T. mutans* and *T. taurotragi* ($p = 0.01$) and *T. mutans* with *T. velifera* ($p = < 0.001$) Table 4.

	<i>A. marginale</i>	<i>A. platys</i> -like	<i>Anaplasma</i> sp.	<i>Ehrlichia</i> sp.	<i>B. bigemina</i>	<i>T. mutans</i>	<i>T. parva</i>	<i>T. taurotragi</i>	<i>T. velifera</i>
<i>A. bovis</i>	C: 1, (0.5) r = -0.008 95% CI: -0.15-0.134 p = 0.914	C: 2, (1.0) r = 0.106 95% CI: -0.037-0.244 p = 0.146	C: 3, (1.6) r = 0.294 95% CI: 0.159-0.419 p = < 0.001*	NCO	NCO	C: 5, (2.6) r = 0.079 95% CI: -0.064-0.219 p = 0.277	C: 1, (0.5) r = -0.047 95% CI: -0.188-0.096 p = 0.519	C: 1, (0.5) r = 0.034 95% CI: -0.109-0.175 p = 0.645	C: 3, (1.6) r = 0.025 95% CI: -0.117-0.166 p = 0.73
<i>A. marginale</i>		NCO	C: 1, (0.5) r = -0.122 95% CI: -0.259-0.021 p = 0.094	NCO	C: 1, (0.5) r = 0.011 95% CI: -0.132-0.152 p = 0.884	C: 34, (17.8) r = -0.035 95% CI: -0.176-0.107 p = 0.63	C: 12, (6.3) r = -0.029 95% CI: -0.17-0.114 p = 0.693	C: 7, (3.7) r = 0.056 95% CI: -0.086-0.197 p = 0.439	C: 21, (11.0) r = -0.025 95% CI: -0.166-0.117 p = 0.731
<i>A. platys</i> -like			C: 6, (3.1) r = 0.162 95% CI: 0.02-0.297 p = 0.026*	NCO	C: 1, (0.5) r = 0.035 95% CI: -0.108-0.176 p = 0.633	C: 26, (13.6) r = 0.031 95% CI: -0.112-0.172 p = 0.674	C: 12, (6.3) r = 0.049 95% CI: -0.094-0.189 p = 0.505	C: 4, (2.1) r = -0.002 95% CI: -0.144-0.14 p = 0.973	C: 20, (10.5) r = 0.136 95% CI: -0.007-0.272 p = 0.061
<i>Anaplasma</i> sp.				1, (0.5) r = -0.074 95% CI: -0.214-0.069 p = 0.309	NCO	C: 16, (8.4) r = 0.104 95% CI: -0.039-0.242 p = 0.154	C: 4, (2.1) r = -0.066 95% CI: -0.266-0.077 p = 0.364	C: 4, (2.1) r = 0.097 95% CI: -0.046-0.235 p = 0.183	C: 9, (4.7) r = 0.004 95% CI: -0.138-0.146 p = 0.96

<i>Ehrlichia</i> sp.	NCO	C: 19, (9.9) r = -0.073 95% CI: -0.213-0.069 p = 0.313	C: 6, (3.1) r = -0.129 95% CI: -0.266-0.013 p = 0.076	C: 3, (1.6) r = -0.024 95% CI: -0.165-0.119 p = 0.744	C: 14, (7.3) r = -0.004 95% CI: -0.146-0.138 p = 0.955
<i>B. bigemina</i>		C: 3, (1.6) r = -0.023 95% CI: -0.164-0.119 p = 0.752	C: 1, (0.5) r = -0.026 95% CI: -0.168-0.116 p = 0.717	NCO	C: 3, (1.6) r = 0.066 95% CI: -0.076-0.206 p = 0.362
<i>T. mutans</i>			C: 57, (29.8) r = 0.087 95% CI: 0.056-0.0226 p = 0.232	C: 25, (13.1) r = 0.187 95% CI: 0.046-0.32 p = 0.01*	C: 96, (50.2) r = 0.344 95% CI: 0.213-0.464 p = <0.001*
<i>T. parva</i>				9, (4.7) r = 0.02 95% CI: -0.122-0.162 p = 0.778	31, (16.2) r = -0.056 95% CI: -0.196-0.087 p = 0.444
<i>T. taurotragi</i>					NCO

317 C = Number of co-infections and percent related to the total cattle population examined (191 specimens), NCO = no co-infection observed, R =
318 correlation coefficient of infections between pathogens in cattle in Malawi, 95% CI = 95% confidence intervals for R, * A p-value of ≤ 0.05 was
319 considered to be significant.

3.3 Phylogenetic analysis

To determine the phylogenetic relationship of the detected pathogens with those reported elsewhere phylogenetic trees were constructed as shown in figures 3, 4, and 5 for *Babesia* and *Theileria* 18S rDNA, *Anaplasma* and *Ehrlichia* species *gltA* and *groEL* genes, respectively. The *Babesia* and *Theileria* sequences obtained in this study clustered with homologous sequences available in the GenBank (Figure 3). We observed divergence in the clustering of *T. mutans* sequences obtained in this study. This finding suggests that there are at least two strains of *T. mutans* circulating in cattle in Malawi. The *T. parva* sequences obtained in this study clustered together with sequences of *T. parva* Muguga cocktail vaccine strains (MG952923, MG952924 and MG952925) even from non-vaccinated cattle.




Figure 3. The maximum likelihood phylogenetic tree of the *Babesia* and *Theileria* species detected in cattle in Malawi. The analysis was based on partial sequences of 18S rDNA which was constructed using the Kimura 2 parameter model. *Plasmodium falciparum* was used as outgroup. All bootstrap values > 75% from 1000 replications are shown on the interior branch nodes and the sequences obtained in this study are indicated in red. The host and country of origin are indicated in parenthesis, NP means not provided. The bar 0.10 is the genetic distance of each sequence from the root of the tree.

338 *Anaplasma marginale* sequences obtained in this study clustered with other
339 homologous sequences obtained from the GenBank. The *A. bovis* sequences obtained from all
340 the five samples were identical and shared 100% identity with homologous sequence of *A.*
341 *bovis* (MH255914) reported from cattle in China. The phylogenetic trees of both the *gltA* and
342 *groEL* genes phylogenetic trees showed that the *A. platys*-like sequences obtained in this
343 study clustered with other *A. platys*-like sequences from ruminants and associated ticks but
344 separated from those reported from dogs or dog ticks (Figures 4 and 5). The obtained
345 sequences of *Ehrlichia* species showed high degree of divergence as evidenced by the
346 branching of their clade. Some sequences had 98-100% identity with homologous sequence
347 of *E. minasensis* (JX629807) isolated from *R. microplus* tick in Brazil. However, the
348 sequences of *Ehrlichia* species clustered separately from those of *E. ruminantium* (Figures 4
349 and 5).




Figure 4. The maximum likelihood phylogenetic tree of *Anaplasma* and *Ehrlichia* species detected in cattle in Malawi. The analysis was based on partial sequences of *gltA* gene which was constructed using the Tamura 2 parameter model. *Rickettsia rickettsii* was used as outgroup. All bootstrap values > 75% from 1000 replications are shown on the interior branch nodes and the sequences obtained in this study are indicated in red. The host and country of origin are indicated in parenthesis, NP means not provided. The bar 0.10 is the genetic distance of each sequence from the root of the tree.




Figure 5. The maximum likelihood phylogenetic tree of the *Anaplasma* and *Ehrlichia* species detected in cattle in Malawi. The analysis was based on partial sequences of *groEL* gene which was constructed using the Tamura 2 parameter model. *Rickettsia rickettsii* was used as outgroup. All bootstrap values > 75% from 1000 replications are shown on the interior branch nodes and the sequences obtained in this study are indicated in bold. The host and country of origin are indicated in parenthesis, NP means not provided. The bar 0.050 is the genetic distance of each sequence from the root of the tree.

4. Discussion

Tick-borne Anaplasmataceae and piroplasms in cattle have been investigated using molecular techniques in the neighboring countries of Malawi, such as Mozambique (Martins et al., 2010, 2008; Alfredo et al. 2005), Tanzania (Ringo et al., 2020, 2018; Swai et al., 2009), and Zambia (Tembo et al., 2018; Simuunza et al., 2011; Makala et al., 2003). However, in Malawi, the epidemiological data on TBPs of cattle is mainly based on basic blood smear and serological examinations. This is the first study that has investigated tick-borne Anaplasmataceae and piroplasms in cattle in Malawi using molecular techniques. DNA nucleotide sequence analysis has shown that cattle in Malawi were infected with *A. bovis*, *A. marginale*, *A. platys*-like, uncharacterized *Anaplasma* species, uncharacterized *Ehrlichia* species, *B. bigemina*, *T. mutans*, *T. parva*, *T. taurotragi*, *T. velifera* and *Theileria* sp. MSD like strain.

The 18S rDNA multiplex PCR assay designed in this study for the identification of *Babesia* and *Theileria* species would be a useful resource in the resource limited settings without the use of sequencing technology in the diagnosis of tick-borne piroplasms especially when co-infections are highly prevalent. This assay may also be used to screen for the targeted *Babesia* and *Theileria* species before animals are introduced in a new area. Multiplex PCR assays have the advantage of being relatively cost effective compared to the sequencing technologies such as next generation sequencing (NGS) (Qiu et al., 2018). However, the major limitation of multiplex PCR assay is that only targeted pathogens can be detected. This does not allow the detection of novel or non-characterized pathogens. On the other hand, although NGS technology is expensive, it has the advantage of being able to detect non-target pathogens (Squarre et al., 2020).

Anaplasma marginale which causes bovine anaplasmosis was detected in samples from all the six sampling sites. The observed prevalence of 24.6% is comparable to 25.7% reported

in Zambia (Tembo et al., 2018). However, it is higher than 10.2%, 7.4% and 18.0% reported in Tanzania (Ringo et al., 2020, 2018; Swai et al., 2007) and 9.4% reported in Mozambique (Martins et al., 2010). The *A. bovis* was detected at a prevalence of 2.9%. Although it is considered non-pathogenic, it has been reported to cause mild disease in calves (Park et al., 2018), and could be fatal when there is co-infection with other tick-borne pathogens (Jurković et al., 2020).

Anaplasma platys-like infection in cattle with a prevalence of 13.6% is being reported for the first time in southern Africa. *Anaplasma platys* is known to be pathogenic in dogs where it cause infectious cyclic thrombocytopenia. The clinical disease in dogs is characterized by lethargy, anorexia, fever, weight loss and bleeding disorder (Skyes & Foley, 2014). The prevalence of 13.6% reported herein is higher than 3.5% reported in Tunisia in North Africa (Ben Said et al., 2017). However, it is lower than 44.7% reported in Kenya in East Africa (Peters et al., 2020). This finding suggests that either the pathogen is expanding its geographical distribution or that it was previously neglected. The phylogenetic trees of the *gltA* and *groEL* genes showed that *A. platys*-like isolates from cattle, water buffalo and associated ticks clustered separately from the isolates from dogs and associated ticks as previously reported (Ogata et al., 2021). This is evident because dogs are specifically infected with the pathogenic *A. platys* while ruminants are infected with genetically related strains (*A. platys*-like).

In this study, *Ehrlichia* species have been detected at a prevalence of 16.2%. All sequences obtained were closely related to sequence of *E. minasensis* (JX629807) isolated from *R. microplus* tick which was previously reported to be closely related to *Ehrlichia canis* (Cabezas-Cruz et al., 2012). This finding is also in accordance with the study conducted in Kenya by Peters et al. (2020), where all sequences of *Ehrlichia* species detected were closely related to *E. minasensis*.

However, *E. ruminantium* which causes bovine ehrlichiosis (heartwater) was not detected in this study, although it has been reported with prevalence ranging from 0.33% to 45.3% in neighboring Zambia (Tembo et al., 2018; Simmunza et al., 2011). The failure to detect the pathogen in this study may be either due to the predisposition of *E. ruminantium* to infect vascular endothelial cells but not circulating cells which limits rickettsemia and chances of being detected in peripheral blood (Deem, 2008).

The prevalence of *B. bigemina* of 2.6% reported in this study is within the same range as 3.3% (Tembo et al., 2018) and 5.5% (Simuunza et al., 2011) reported in Zambia and 5.1% (Ringo et al., 2020) reported in Tanzania. However, in this study, *B. bovis* which is the most pathogenic *Babesia* species was not detected in the examined animals. This is in contrast to 7.7% (Tembo et al., 2018) and 15.5% (Simuunza et al., 2011) reported in Zambia and 4.5% reported in Tanzania (Ringo et al., 2020). The negative results of *B. bovis* may be due to low prevalence of the pathogen in the sampled areas or due to small sample size used in this study.

Theileria parva which causes East Coast fever (ECF) was one of the pathogens reported with the highest frequency in this study. The detection of *T. parva* at all the six sampling sites including the previously non-endemic region indicates that the pathogen has expanded its geographical distribution in Malawi. The prevalence in this study (33.0 %) is comparable to 34.4% and 41.2% reported in Tanzania (Ringo et al., 2020; 2018) but higher than 24.4% and 0.3% reported in Zambia (Simuunza et al., 2011; Tembo et al., 2018). However, because in some sampling sites in the central region the Muguga cocktail vaccine is used to control ECF some positive samples may be due to the vaccinations. The phylogenetic analysis also showed that the sequences of *T. parva* from Malawi clustered together with sequences obtained from the components of the Muguga cocktail vaccine (Muguga, Kiambu-5 and Serengeti transformed) deposited in GenBank even those from animals without vaccination history. Furthermore, the obtained sequences of *T. parva* clustered with the sequence of Katete vaccine

strain from Zambia. Since the stocks of the Muguga cocktail vaccine are originally from Kenya and Tanzania, the findings of this study may suggest that the strains of *T. parva* circulating in cattle in eastern and southern Africa are closely related to one another and the vaccine strains. However, more detailed studies are required to support this hypothesis as the marker used (18S rDNA) is too conserved amongst the *T. parva* strains.

There was high infection rates of *T. mutans* and *T. velifera* at 73.8% and 53.4%, respectively. These rates are higher than 38.1% and 3.4% reported in Tanzania (Ringo et al., 2020) and 54.5% and 51.5% in Zambia (Tembo et al., 2018), respectively. *Theileria mutans* has been reported to be the most prevalent tick-borne pathogen in both Tanzania (Ringo et al., 2020, 2018) and Zambia (Tembo et al., 2018; Simuunza et al., 2011). The *T. mutans* sequences obtained in this study clustered in two clades, one with *T. mutans* (AF078815) isolated from cattle in South Africa and another with *Theileria* sp. MSD like (AF078816) a strain of *T. mutans* (Chaisi et al., 2013) reported from cattle in South Africa. This shows that there are at least two strains of *T. mutans* circulating in cattle in Malawi. The finding of closely related strains reported in South Africa may suggest that most strains circulating in cattle in southern Africa are closely related. Although *T. mutans* infection in cattle is considered less pathogenic, it has been reported that mixed infection with *T. parva* increases the pathogenicity of *T. parva* (Morrison, 2010).

Theileria taurotragi which is considered less pathogenic in cattle but when clinical disease occurs it has been reported to present with clinical signs like those of ECF was also detected in cattle in Malawi. The phylogenetic tree of the 18S rDNA supports the close relatedness with *T. parva* as they cluster in the same clade. The vector tick for both *T. parva* and *T. taurotragi* is the brown ear tick *R. appendiculatus* (Walker et al., 20003; Ringo et al., 2020). The positive detection rate of 12.6% reported in this study is lower than 30.9% reported

in Tanzania (Ringo et al., 2020) and 27.5% reported in Zambia (Simuunza et al., 2011). However, it is comparable to 11.4% reported by another study in Zambia (Tembo et al., 2018).

The detection of co-infections in 152 animals (79.6%) with both Anaplasmataceae and piroplasms is an important finding as it complicated the proper diagnosis and treatment of the animals that show clinical disease as some combinations need different treatment regimen. The co-infection rate reported in this study was higher than those reported in neighboring Tanzania of 36.4% (Ringo et al., 2020) but lower than those reported in Zambia of 52.9% and 75.6% by Simuunza et al. (2011) and Tembo et al. (2018), respectively. Mixed infections with TBPs that are individually less pathogenic than other species within the same genus have been reported to cause clinical disease (Jurković et al., 2020). However, another study has suggested that this may be beneficial to the animals as the less pathogenic species may help to generate immunity against the more pathogenic species by heterologous protection (Woolhouse et al., 2015). In this study, although there were many combinations of mixed infections, only four combinations were statistically significant suggesting that these pathogens have a high likelihood of being detected when there is co-infection and may be transmitted by same tick species. *Theileria mutans* and *T. velifera* co-infection correlation in cattle was also reported in Zambia (Tembo et al., 2018). However, in this study, there was no co-infection correlation between *T. parva* and the causative agents of benign theilerioses (*T. mutans*, *T. taurotragi* and *T. velifera*) in this study as also previously reported in Zambia (Tembo et al., 2018). This finding shows that the challenge to TBPs is widespread in the southern Africa and collaborative efforts to develop novel methods to control ticks and tick-borne diseases are warranted.

The high co-infection rate may have resulted from the examined animals being infested with different tick species which are vectors of different pathogens detected in this study at the same time (Foughali et al., 2021). The transmission of different pathogens by the same tick species such as *T. parva* and *T. taurotragi* which are both transmitted by *R. appendiculatus*

may also play a role in the high co-infection rate (Ringo et al., 2020). Since there is no data on tick's specificity, further studies focusing on tick tropism to understand the vector competence of the ticks infesting cattle in Malawi are required.

Declaration of Competing Interest

The authors declare no conflict of interest.

Acknowledgments

This study was financially supported in part by JSPS KAKENHI (grant numbers 19H03118, 20K21358, and 20KK0151). International Collaborative Research Programme for Tackling Neglected Tropical Disease (NTD) Challenges in African countries (JP18jm0510001) of Japan Agency for Medical Research and Development (AMED). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

R.N. and E.C. conceived and designed the experiments; R.N. and C.S. obtained funding; E.C, E.M, collected samples; E.C., G.P., A.E.A., performed the experiments; E.C. and W.M.A.M., analyzed the data; E.C. Writing- Original draft preparation; K.H., C.S., K.K., N.N. and R.N. Supervision; K.H., K.K., N.N., C.S. and R.N. Writing- Reviewing and Editing. All authors approved the final version of the manuscript.

References

- Alfredo, A., Jonsson, N., Finch, T., Neves, L., Molloy, J. and Jorgensen, W., 2005. Serological survey of *Babesia bovis* and *Anaplasma marginale* in cattle in Tete Province, Mozambique. *Trop. Anim. Health Prod.* 37, 121–131
- Allsopp, B.A., 2015. Heartwater - *Ehrlichia ruminantium* infection. *Rev. Sci. Tech.* 34, 557-568
- Alvasen, K., Johansson, S.M., Hoglund, J., Ssuna, R., Emanuelson, U., 2016. A field survey on parasites and antibodies against selected pathogens in owned dogs in Lilongwe, Malawi. *J. S. Afr. Vet. Assoc.* 87, e1–e6
- Ben Said, M., Belkahia, H., El Mabrouk, N., Saidani, M., Alberti, A., Zobba, R., Cherif, A., Mahjoub, T., Bouattour, A., Messadi, L., 2017. *Anaplasma platys*-like strains in ruminants from Tunisia. *Infect. Genet. Evol.* 49, 226–233
- Berggren, A., 1978. Cattle ticks in Malawi. *Vet. Parasitol.* 4, 289-297
- Kocan, K.M., de la Fuente, J., Guglielmone, A.A., Melendez, R.D., 2003. Antigens and alternatives for control of *Anaplasma marginale* infection in cattle. *Clin. Microbiol. Rev.* 16, 698–712
- Cabezas-Cruz, A., Zweggarth, E., Vancová, M., Broniszewska, M., Grubhoffer, L., Passos, L.M.F., Ribeiro, M.F.B., Alberdi, P., de la Fuente, J., 2016. *Ehrlichia minasensis* sp. nov., isolated from the tick *Rhipicephalus microplus*. *Int. J. Syst. Evol. Microbiol.* 66, 1426-1430
- Chatanga, E., Hayashida, K., Muleya, W., Kusakisako, K., Moustafa, M.A.M., Salim, B., Katakura, K., Sugimoto, C., Nonaka, N., Nakao, R., 2020. Genetic Diversity and Sequence polymorphism of two genes encoding *Theileria parva* antigens recognized by CD8⁺ T cells among vaccinated and unvaccinated cattle in Malawi. *Pathogens* 9, 334.

537 Chatanga, E., Kainga, H., Maganga, E., Hayashida, K., Katakura, K., Sugimoto, C., Nonaka,
 538 N., Nakao, R., 2021a. Molecular identification and genetic characterization of tick-
 539 borne pathogens in sheep and goats at two farms in the central and southern regions of
 540 Malawi. *Ticks Tick Borne Dis.* 12, 101629

541 Chatanga, E., Kainga, H., Razemba, T. Ssuna, R., Swennen, L., Hayashida, K., Sugimoto, C.,
 542 Katakura, K., Nonaka N., Nakao, R., 2021b. Molecular detection and characterization
 543 of tick-borne hemoparasites and Anaplasmatidae in dogs in major cities of Malawi.
 544 *Parasitol. Res.* 120, 267–276

545 Chinombo, D.O., Mzoma, F.J., Musisi, F.L., 1988. East Coast fever in Malawi (1983–1987).
 546 In theileriosis in Eastern, Central and Southern Africa. In proceedings of a workshop
 547 on East Coast fever immunization, Lilongwe, Malawi, 20–22 September 1988, Dolan,
 548 T.T., (Ed.), ILRAD: Nairobi, Kenya, pp. 12–16

549 DAHLD, 2006. Policy document on livestock in Malawi. Ministry of Agriculture, Irrigation
 550 and Water Development (MoAIWD); Department of Animal Health and Livestock
 551 Development (DAHLD): Lilongwe, Malawi.

552 DAHLD, 2020. 2019/2020 Third round agricultural production estimates (APES). Ministry
 553 of Agriculture, Irrigation and Water Development (MoAIWD); Department of Animal
 554 Health and Livestock Development (DAHLD): Lilongwe, Malawi.

555 Dumler, J.S., Barbet, A.F., Bekker, C.P.J., Dasch, G.A., Palmer, G.H., Ray, S.C., Rikihisa,
 556 Y., Rurangirwa, F.R., 2001. Reorganization of genera in the families Rickettsiaceae
 557 and Anaplasmatidae in the order Rickettsiales: Unification of some species of
 558 *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*,
 559 descriptions of six new species combinations and designation of *Ehrlichia equi* and
 560 'HGE agent' as subjective synonyms of *Ehrlichia phagocytophila*. *Int. J. Syst. Evol.*
 561 *Microbiol.* 51, 2145-2165

562 Foughali, A. A., Jedidi, M., Dhibi, M., Mhadhbi, M., Sassi, L., Berber, A., Bitam, I., Gharbi,
 563 M., 2021. Infection by haemopathogens and tick infestation of sheep during summer
 564 season in Constantine region, Northeast Algeria. *Vet. Med. Sci.* 7, 1769–1777.

565 Gofton, A.W., Doggett, S., Ratchford, A., Ryan, U., Irwin, P., 2016. Phylogenetic
 566 characterization of two novel Anaplasmataceae from Australian *Ixodes holocyclus*
 567 ticks: '*Candidatus* Neoehrlichia australis' and '*Candidatus* Neoehrlichia arcana'. *Int.*
 568 *J. Syst. Evol. Microbiol.* 66, 4256–4261

569 Inokuma, H., Oyamada, M., Kelly, P.J., Jacobson, L.A., Fournier, P.E., Itamoto, K., 2005.
 570 Molecular detection of a new *Anaplasma* species closely related to *Anaplasma*
 571 *phagocytophilum* in canine blood from South Africa. *J. Clin. Microbiol.* 43, 2934–
 572 2937

573 Johansson, S., 2015. General health conditions in the dog population of Lilongwe. In Epsilon
 574 archive for student projects. Examensarbete, Upsala, Sweden. 34, 1-31

575 Jurković, D., Mihaljević, Ž., Duvnjak, S., Silaghi, C., Beck, R., 2020. First reports of
 576 indigenous lethal infection with *Anaplasma marginale*, *Anaplasma bovis* and
 577 *Theileria orientalis* in Croatian cattle. *Ticks Tick Borne Dis.* 11, 101469

578 Kumar, S., Stecher, G., Tamura, K., 2016. 'MEGA7: molecular evolutionary genetics
 579 analysis Version 7.0 for bigger datasets'. *Mol. Biol. Evol.* 33, 1870–1874

580 Lawrence, J.A., Musisi, F.L., Mfitilodze, M.W., Tjornehoj, K., Whiteland, A.P., Kafuwa,
 581 P.T., Chamambala, K.E., 1996. Integrated tick and tick-borne disease control trials in
 582 crossbred dairy cattle in Malawi. *Trop. Anim. Health Prod.* 28, 280–288.

583 Liz, J.S., Anderes, L., Sumner, J.W., Massung, R.F., Gern, L., Rutti, B., Brossard, M., 2000.
 584 PCR detection of granulocytic ehrlichiae in *Ixodes ricinus* ticks and wild small
 585 mammals in western Switzerland. *J. Clin. Microbiol.* 38, 1002–1007

586 Lounsbury, C.P., 1904. Transmission of African Coast fever. Agr. J. Cape Good Hope. 24,
 587 428-432

588 Lubembe, D.M., Odongo D.O., Salih, D.A., Sibeko-Matjila, K.P., 2020. Microsatellite and
 589 minisatellite genotyping of *Theileria parva* population from southern Africa reveals
 590 possible discriminatory allele profiles with parasites from eastern Africa. Ticks Tick
 591 Borne Dis. 11, 101539

592 Makala, L.H., Mangani, P., Fujisaki, K., Nagasawa, H., 2003. The current status of major
 593 tick-borne diseases in Zambia. Vet. Res. 35, 467–483

594 Martins, T., Neves, L., Pedro, O., Fafetine, J., Do Rosário, V., Domingos, A., 2010.
 595 Molecular detection of *Babesia* spp. and other haemoparasitic infections of cattle in
 596 Maputo Province, Mozambique. Parasitology 137, 939-946

597 Martins, T.M., Pedro, O.C., Caldeira, R.A., do Rosário, V.E., Neves, L., Domingos, A., 2008.
 598 Detection of bovine babesiosis in Mozambique by a novel semi-nested hot-start PCR
 599 method. Vet. Parasitol. 153, 225-230

600 Masatani, T., Hayashi, K., Andoh, M., Tateno, M., Endo, Y., Asada, M., Kusakisako, K.,
 601 Tanaka, T., Gokuden, M., Hozumi, N., Nakadohzo, F., Matsuo, T., 2017. Detection
 602 and molecular characterization of *Babesia*, *Theileria*, and *Hepatozoon* species in hard
 603 ticks collected from Kagoshima, the southern region in Japan. Ticks Tick Borne Dis.
 604 8, 581–587

605 Moll, G., Lohding, A., Young, A., 1984. Epidemiology of theilerioses in the Trans-Mara
 606 division, Kenya: Husbandry and disease background and preliminary investigations
 607 on theilerioses in calves. Prev. Vet. Med. 2, 801–831

608 Moll, G., Lohding, A., Young, A., Leitch, B., 1986. Epidemiology of theileriosis in calves in
 609 an endemic area of Kenya. Vet. Parasitol. 19, 255–273

610 Morrison, W.I., 2010. Theilerias. In the Merck Veterinary Manual. Merck Publishing Co.
611 Inc. Whitehouse Station, New Jersey, USA.10, 33-35

612 Nene, V., Kiara, H., Lacasta, A., Pelle, R., Svitek, N., Steinaa, L., 2016. The biology of
613 Theileria parva and control of East Coast fever – Current status and future trends. Ticks
614 Tick Borne Dis. 7, 549–564

615 NSO, 2018. 2018 Malawi population and housing census main report. Zomba, Malawi.
616 [http://populationmalawi.org/wp1/wp-content/uploads/2019/10/2018-Malawi-](http://populationmalawi.org/wp1/wp-content/uploads/2019/10/2018-Malawi-Population-and-Housing-Census-Main-Report-1.pdf)
617 [Population-and-Housing-Census-Main-Report-1.pdf](http://populationmalawi.org/wp1/wp-content/uploads/2019/10/2018-Malawi-Population-and-Housing-Census-Main-Report-1.pdf) accessed on 19 April, 2020.

618 Ogata, S., Pereira, J.A.C., Jhonny, L.V.A., Carolina, H.P.G., Matsuno, K., Orba, Y., Sawa,
619 H., Kawamori, F., Nonaka, N., Nakao, R., 2021. Molecular Survey of *Babesia* and
620 *Anaplasma* infection in cattle in Bolivia. Vet. Sci. 8, 188

621 Oura, C.A.L., Bishop, R., Asiimwe, B.B., Spooner, P., Lubega, G.W., Tait, A., 2007. *Theileria*
622 *parva* live vaccination: parasite transmission, persistence and heterologous challenge
623 in the field. Parasitology 134, 1205–1213

624 Park, J., Han, D.G., Ryu, J.H., Chae, J.B., Chae, J.S., Yu, D.H., Park, B.K., Kim, H.C., Choi,
625 K.S., 2018. Molecular detection of *Anaplasma bovis* in Holstein cattle in the Republic
626 of Korea. Acta Vet. Scand. 60, 15

627 Peters, S.G., Aboge, G.O., Kariuki, H.W., Kanduma, E.G., Gakuya, D.W., Maingi, N., Mulei,
628 C.M., Mainga, A.O., 2020. Molecular prevalence of emerging *Anaplasma* and
629 *Ehrlichia* pathogens in apparently healthy dairy cattle in peri-urban Nairobi, Kenya.
630 BMC Vet. Res. 16, 364

631 Peters, S.G., Gakuya, D.W., Maingi, N., Mulei, C.M., 2019. Prevalence and risk factors
632 associated with *Ehrlichia* infections in smallholder dairy cattle in Nairobi City
633 County, Kenya. Vet. World, 12, 1599-1607

634 Qiu, Y., Kaneko, C., Kajihara, M., Saasa, N., Simulundu, E., Muleya, W., Thu, M.J.,
 635 Hang'ombe, M.B., Katakura, K., Takada, A., Sawa, H., Simuunza, M., Nakao, R.,
 636 2018. Tick-borne haemoparasites and Anaplasmataceae in domestic dogs in Zambia.
 637 Ticks Tick Borne Dis. 9, 988–995

638 Rar, V.A., Epikhina, T.I., Livanova, N.N., Panov, V.V., Doroschenko, E.K., Pukhovskaya,
 639 N.M., Vysochina, N.P., Ivanov. L.I., 2011. Genetic variability of *Anaplasma*
 640 *phagocytophilum* in *Ixodes persulcatus* ticks and small mammals in the Asian part of
 641 Russia. Vector Borne Zoonotic Dis. 11, 1013-1021

642 Richey, E.J., Palmer, G.H., 1990. Bovine anaplasmosis. Compend. Contin. Educ. Pract. Vet.
 643 12, 1661–1668

644 Ringo, A.E., Rizk, M.A., Moumouni, P.F.A., Liu, M., Galon, E.M., Li, Y., Ji, S.,
 645 Tumwebaze, M., Byamukama, B., Thekisoe, O., Xuan, X., 2020. Molecular detection
 646 and characterization of tick-borne haemoparasites among cattle on Zanzibar Island,
 647 Tanzania. Acta Trop. 211, 105598

648 Ringo, A.E., Adjou Moumouni, P.F., Lee, S.H., Liu, M., Khamis, Y.H., Gao, Y., Guo, H.,
 649 Zheng, W., Efstratiou, A., Galon, E.M., Li, J., Tiwananthagorn, S., Inoue, N., Suzuki,
 650 H., Thekisoe, O., Xuan, X., 2018. Molecular detection and characterization of tick-
 651 borne protozoan and rickettsial pathogens isolated from cattle on Pemba Island,
 652 Tanzania. Ticks Tick Borne Dis. 9, 1437–1445

653 Simuunza, M., Weir, W., Courcier, E., Tait, A., Shiels, B., 2011. Epidemiological analysis of
 654 tick-borne diseases in Zambia. Vet. Parasitol. 175, 331–342

655 Sivakumar, T., Tuvshintulga, B., Kothalawala, H., Silva, S.S.P., Lan, D.T.B., Long, P.T.,
 656 Ybanez, A.P., Ybanez, R.H.D., Francisco Benitez, D., Tayebwa, D.S., 2020. Host
 657 range and geographical distribution of *Babesia* sp. Mymensingh. Transbound. Emerg.
 658 Dis. 7, 2233–2239

659 Sivakumar, T., Tuvshintulga, B., Zhyldyz, A., Kothalawala, H., Yapa, P.R., Kanagaratnam,
 660 R., Vimalakumar, S.C., Abeysekera, T.S., Weerasingha, A.S., Yamagishi, J., Igarashi,
 661 I., Silva, S.S.P., Yokoyama, N., 2018. Genetic Analysis of *Babesia* Isolates from
 662 Cattle with Clinical Babesiosis in Sri Lanka. J. Clin. Microbiol. 56, e00895-18
 663 Squarre, D., Nakamura, Y., Hayashida, K., Kawai, N., Chambaro, H., Namangala, B.,
 664 Sugimoto, C., Yamagishi, J., 2020. Investigation of the piroplasm diversity circulating
 665 in wildlife and cattle of the greater Kafue ecosystem, Zambia. Parasit. Vectors, 13,
 666 599.

667 Swai, E.S., Karimuribo, E.D., Kambarage, D.M., Moshy, W.E., 2009. A longitudinal study
 668 on morbidity and mortality in young stock smallholder dairy cattle with special
 669 reference to tick-borne infections in Tanga region, Tanzania. Vet. Parasitol. 160, 34–
 670 42

671 Swai, E.S., Karimuribo, E.D., Kambarage, D.M., Moshy, W.E., Mbise, A.N., 2007. A
 672 comparison of sero-prevalence and risk factors for *Theileria parva* and *T. mutans* in
 673 smallholder dairy cattle in the Tanga and Iringa regions of Tanzania. Vet. J. 174, 390–
 674 396

675 Tembo, S., Collins, N.E., Sibeko-Matjila, K.P., Troskie, M., Vorster, I., Byaruhanga, C.,
 676 Oosthuizen, M.C., 2018. Occurrence of tick-borne haemoparasites in cattle in the
 677 Mungwi district, Northern Province, Zambia. Ticks Tick Borne Dis. 9, 707–717.

678 Uilenberg, G., 1995. International collaborative research: Significance of tick-borne
 679 hemoparasitic diseases to world animal health. Vet. Parasitol. 57, 19–41

680 Walker, A.R., Bouattour, A., Camicas, J.L., Estrada Peña, A., Horak, I., Latif, A., Pegram,
 681 R.G., Preston, P.M., 2003. Ticks of Domestic Animals in Africa: A guide to
 682 Identification of Species. Edinburgh (Scotland) Bioscience Reports, Edinburgh, UK.
 683 Pp 1-22

684 Woolhouse, M.E., Thumbi, S.M., Jennings, A., Chase-Topping, M., Callaby, R., Kiara, H.,
685 Oosthuizen, M.C., Mbole-Kariuki, M.N., Conradie, I., Handel, I.G., Poole, E.J., Njiiri,
686 E., Collins, N.E., Murray, G., Tapio, M., Auguet, O.T., Weir, W., Morrison, W.I.,
687 Kruuk, L.E., Bronsvoort, B.M., Hanotte, O., Coetzer, K., Toye, P.G., 2015. Co-
688 infections determine patterns of mortality in a population exposed to parasite
689 infection. Sci. Adv. 1, e1400026

690 Young, A.S., Grootenhuys, J.G., Kimber, C.D., Kanhai, G.K., Stagg, D.A., 1977. Isolation of
691 a *Theileria* species from Eland (*Taurotragus oryx*) infective for cattle. Tropenmed
692 Parasitol. 28, 185–194.