



Title	Serological characterization of lineage II insect-specific flaviviruses compared with pathogenic mosquito-borne flaviviruses
Author(s)	Tabata, Koshiro; Itakura, Yukari; Toba, Shinsuke et al.
Citation	Biochemical and biophysical research communications, 616, 115-121 https://doi.org/10.1016/j.bbrc.2022.05.080
Issue Date	2022-08-06
Doc URL	https://hdl.handle.net/2115/90263
Rights	© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	tabata BBRC Manuscript_com.pdf



1 **Serological characterization of lineage II insect-specific**
2 **flaviviruses compared with pathogenic mosquito-borne**
3 **flaviviruses**

4
5 Koshiro Tabata^a, Yukari Itakura^a, Shinsuke Toba^{a,b}, Kentaro Uemura^{a,b,c}, Mai
6 Kishimoto^a, Michihito Sasaki^a, Jessica J. Harrison^g, Akihiko Sato^{a,b}, William W. Hall^{d,e,f},
7 Roy A. Hall^g, Hirofumi Sawa^{a,d,e,h}, Yasuko Orba^{a,d,*}

8
9 **Author affiliations:**

10 ^a Division of Molecular Pathobiology, International Institute for Zoonosis Control,
11 Hokkaido University, Sapporo, Hokkaido, 001-0020, Japan

12 ^b Shionogi & Co., Ltd., Osaka, 541-0045, Japan

13 ^c Laboratory of Biomolecular Science, Faculty of Pharmaceutical Science, Hokkaido
14 University, Sapporo, 060-0812, Japan

15 ^d International Collaboration Unit, International Institute for Zoonosis Control,
16 Hokkaido University, Sapporo, Hokkaido, 001-0020, Japan

^e Global Virus Network, Baltimore, Maryland, 21201, USA

^f National Virus Reference Laboratory, School of Medicine, University College Dublin,
Belfield, Dublin, 4, Ireland

^g Australian Infectious Diseases Research Centre, School of Chemistry and Molecular
Biosciences, University of Queensland, St Lucia, Queensland, Australia

^h One Health Research Center, Hokkaido University, Sapporo, Hokkaido, 001-0020,
Japan

***Corresponding author:**

E-mail: orbay@czc.hokudai.ac.jp

Postal address: International Institute for Zoonosis Control, Hokkaido University, N20,
W10, Kita-ku, Sapporo, 001-0020, Hokkaido, Japan

Keywords: lineage II insect-specific flavivirus, BJV, PSFV, antibody, cross-reactivity,
antibody-dependent enhancement (ADE)

ABSTRACT

The genus *Flavivirus* includes pathogenic tick- and mosquito-borne flaviviruses as well as non-pathogenic insect-specific flaviviruses (ISFVs). Phylogenetic analysis based on whole amino acid sequences has indicated that lineage II ISFVs have similarities to pathogenic flaviviruses. In this study, we used reactive analysis with immune serum against Psorophora flavivirus (PSFV) as a lineage IIa ISFV, and Barkeji virus (BJV) as a lineage IIb ISFV, to evaluate the antigenic similarity among lineage IIa and IIb ISFVs, and pathogenic mosquito-borne flaviviruses (MBFVs). Binding and antibody-dependent enhancement assays showed that anti-PSFV sera had broad cross-reactivity with MBFV antigens, while anti-BJV sera had low cross-reactivity. Both of the lineage II ISFV antisera were rarely observed to neutralize MBFVs. These results suggest that lineage IIa ISFV PSFV has more antigenic similarity to MBFVs than lineage IIb ISFV BJV.

1. Introduction

Flavivirus is a genus of single-stranded, enveloped RNA viruses, which includes tick-borne flaviviruses (TBFVs), mosquito-borne flaviviruses (MBFVs), and insect-specific flaviviruses (ISFVs). TBFVs and MBFVs infect vertebrates sometimes

51 producing disease. ISFVs are phylogenetically divided into lineage I (also called
52 classical ISFV) and lineage II. In phylogenetic analyses, lineage II ISFVs cluster with
53 the MBFVs [1]. Therefore, lineage II ISFVs are considered to be dual-host affiliated
54 ISFVs, which may be transmitted to mammals other than mosquitoes, as are MBFVs
55 [2,3]. However, there have been no reports of lineage II ISFVs infecting vertebrate cells
56 under natural conditions [4,5].

57 Flaviviruses have positive-sense, single-stranded RNA genomes encoding a
58 polyprotein comprised of structural and nonstructural proteins [6]. Reactive analyses
59 using antibodies against the structural protein of MBFVs have demonstrated that those
60 antibodies are mutually cross-reactive among MBFVs, especially Dengue virus
61 (DENV), Zika virus (ZIKV), Japanese encephalitis virus (JEV), and West Nile virus
62 (WNV) [7-11]. These studies suggest that structural proteins are conserved among these
63 MBFVs. Recently, hemagglutination inhibition assays and reactive analyses using a
64 panel of anti-lineage II ISFV monoclonal antibodies showed that lineage II ISFV also
65 possessed antigenically similar structural proteins to MBFVs [3,12,13].

66 The lineage II ISFVs are phylogenetically further divided into lineage IIa and
67 IIb, and lineage IIa ISFVs seem to be associated *Aedes* species [14]. While the antigenic
68 relationship between lineage II ISFV and MBFV have been investigated, the

characterization of lineage IIa and IIb ISFVs compared with MBFVs are still poorly understood. In this study, we performed antibody-based analyses using anti-lineage IIa and IIb ISFV sera to MBFV antigens to identify antigenic similarity among lineage IIa and IIb ISFVs, and the MBFVs. Therefore, we used two lineage II ISFVs, Psorophora flavivirus (PSFV; lineage IIa ISFV) isolated from *Psorophora albigenu* in Bolivia, and Barkedji virus (BJV; lineage IIb ISFV) isolated from *Culex* spp. in Zambia [2,14].

2. Materials and methods

2.1. Cells

Vero9013 cells (Vero cells; JCRB, Ibaraki, Japan) were maintained in Dulbecco's Modified Eagle Medium (DMEM; Nissui, Tokyo, Japan) supplemented with 10% fetal bovine serum (FBS). K562 cells (JCRB) were maintained in RPMI medium (Thermo Fisher Scientific) supplemented with 10% FBS. Vero cells and K562 cells were cultured at 37°C and 5% CO₂. *Aedes albopictus*-derived C6/36 mosquito cells (ATCC, Manassas, VA, USA) were cultured at 28°C, 5% CO₂ in Eagle's Minimum Essential Medium (MEM; Nissui) supplemented with 10% FBS and non-essential amino acids (Thermo Fisher Scientific, Waltham, MA, USA).

2.2. Viruses

Dengue virus type 2 (DENV2) hu/INDIA/09-74 strain (GenBank accession no. LC367234), ZIKV MR766-NIID strain (GenBank accession no. LC002520), JEV Beijing strain (GenBank accession no. L48961), and WNV zmq16m11 strain (GenBank accession no. LC318700) were used as vertebrate-infecting mosquito-borne flaviviruses (MBFVs). Psorophora flavivirus (PSFV) (GenBank accession no. LC567151) and Barkedji virus (BJV) zmq17L31 strain (GenBank accession no. LC497470) were used as lineage IIa and IIb ISFVs, respectively. All viral strains were propagated in C6/36 cells in MEM containing 2% FBS and incubated for three to five days, after which culture supernatants were collected and stored at -80°C until use.

2.3. Analysis of homology and molecular phylogenetics

The genome sequences of the flaviviruses used were obtained from GenBank. The sequence similarities of the MBFVs and the lineage II ISFVs used in this study were evaluated using BLAST analysis. For phylogenetical analyses, amino acid sequence alignments based on whole amino acid sequences or precursor membrane and envelope (prME) proteins were generated using the MUSCLE algorithm of the CLC Genomics Workbench 20.1 (Qiagen, Hilden, Germany). Maximum-likelihood (ML)

phylogenetic trees were constructed using the best-fit models of the Le_Gascuel_2008 model with gamma rate categories with invariant sites, with empirical frequencies (LG+G+I+F) for the whole amino acid sequences and LG+G+I for the prME protein, and bootstrap values of 1000 replicates in MEGA X software [15]. The phylogenetic trees were visualized using Interactive Tree Of Life version 6.5.2 [16].

2.4. Focus forming assays (FFA)

Vero cells (1.5×10^5 cells/well) for MBFVs or C6/36 cells (1.0×10^6 cells/well) for lineage II ISFVs, grown in 24-well plates (Corning, Tewksbury, MA, USA) were inoculated with 10-fold serial dilutions of each virus and incubated for 1 h at 37°C (Vero cells) or 28°C (C6/36 cells). An overlay medium was added (MEM containing both 0.5% methyl cellulose and 2% FBS) and incubated for 32 h (ZIKV, JEV, and WNV) or 72 h (PSFV, BJV, and DENV2). After incubation, the cells were fixed with pre-chilled methanol at -30°C for more than 10 min. The fixed cells were blocked with phosphate-buffered saline (PBS) containing 1% bovine serum albumin (BSA) at room temperature (RT) for 30 min. Anti-flavivirus nonstructural protein 1 (NS1) monoclonal 4G4 antibody [17] for MBFVs or anti-flavivirus envelope monoclonal BJ-6E6 antibody [13] for lineage II ISFVs in 0.1% BSA-PBS was then added to each well and incubated

at RT for 1 h. Alexa Fluor 488-conjugated anti-mouse IgG antibody (Invitrogen, Waltham, MA, USA) in 0.1% BSA-PBS was applied as a secondary antibody at RT for 1 h, and viral focus images were acquired using a fluorescence microscope (IX73, Olympus, Tokyo, Japan). The virus titers were calculated based on the number of foci and expressed as focus-forming units (FFU) per ml.

2.5. Ethical statement

All animal experiments were performed in accordance with the National University Corporation, Hokkaido University, Regulations on Animal Experimentation. The protocol was approved by the Institutional Animal Care and Use Committee of Hokkaido University (approval nos. 18-0036 and 18-0149).

2.6. MBFV-infected serum

AG129 mice deficient in interferon- α/β and - γ receptors were obtained from Marshall BioResources (Hull, East Yorkshire, UK), and bred in-house in an animal facility. Ten-week-old male AG129 mice were inoculated subcutaneously with 1.0×10^5 FFU of DENV2 or ZIKV. Six-week-old female BALB/c mice (Japan SLC, Shizuoka, Japan) were inoculated subcutaneously with 1.0×10^6 FFU of JEV or 1.0×10^4 FFU of WNV. The body weight of the virus-inoculated mice was monitored daily. When mice

showed a 10% decrease in their body weight, they were euthanized using excess isoflurane anesthesia, and sera were collected. The mouse sera were inactivated by incubation at 56°C for 30 min and stored at –80°C until use.

2.7. Virus purification

At four days post inoculation with either PSFV or BJV in C6/36 cells, the culture supernatants were harvested and filtrated through a 0.45 µm syringe filter and pelleted by ultracentrifugation with a 20% sucrose cushion using an SW32Ti rotor (Beckman Coulter, Brea, CA, USA) at 153,720 ×g for 3 h [18]. The pellets were resuspended in PBS. The protein concentrations of the resuspended samples were determined using BCA assays (Thermo Fisher Scientific) conducted according to the manufacture's protocol. Virus suspensions were stored at –80°C until immunization.

2.8. Immunization

Six-week-old female BALB/c mice were subcutaneously inoculated with 20 µg of PSFV or BJV, or PBS as a negative control. Four weeks post inoculation, the mice were euthanized using excess isoflurane anesthesia, and the sera were collected, inactivated by incubation at 56°C for 30 min and stored at –80°C until use.

2.9. Enzyme-linked immunosorbent assay (ELISA)

Vero cells (6.0×10^5 cells/well) or C6/36 cells (4.0×10^6 cells/well) in six-well plates (Corning) were inoculated with MBFV or lineage II ISFV at a multiplicity of infection (MOI) of 0.1. At 72 hours post infection (hpi) (PSFV, BJV, and DENV2) or 48 hpi (ZIKV, JEV, and WNV), the cells were lysed in lysis buffer [1% NP-40, 20 mM Tris-HCl (pH 7.5), 150 mM NaCl, 5 mM EDTA]. After centrifugation to remove the cell debris, the virus-infected cell lysates were collected from the supernatant. Flat-bottom 384-well plates (Thermo Fisher Scientific) were coated with 50-fold diluted cell lysates at 37°C for 3 h. The plates were blocked with 5% skim milk-PBS at RT for 30 min. Immunized mouse sera and pan-flavivirus 4G2 monoclonal antibody (ATCC) used as a positive control were serially 10-fold diluted with 1% skim milk-PBS and added to each well. Following incubation for 2 h at 37°C, the plates were washed five times with 0.01% PBS-Tween20 (PBS-T). Horseradish peroxidase (HRP) conjugated anti-mouse IgG antibody (Thermo Fisher Scientific) was added and incubated at 37°C for 1 h. After washing five times, 1-Step Ultra TMB-ELISA Substrate Solution (Thermo Fisher Scientific) was added as a HRP substrate. After incubation at RT for 15 min, the reaction was stopped by the addition of 2 M sulfuric acid. The optical density value was

measured at 450 nm using a GroMax Discover Microplate Reader (Promega, Madison, WI, USA). The binding titer was defined as the endpoint serum dilution, which is the reciprocal of the highest dilution over the cut-off value; the mean plus three times the standard deviation of non-immunized control sera. The values lower than the cut-off value were expressed as the limit of detection.

2.10. Foci reduction neutralization test (FRNT)

Vero cells (7.5×10^4 cells/well) or C6/36 (5.0×10^5 cells/well) cells were seeded in 48-well plates (Corning) 24 h before infection. Serum samples were serially 10-fold diluted with DMEM containing 2% FBS and incubated with 50 FFU of each virus at 37°C for 1 h. The cells were inoculated with the serum-virus complexes and incubated at 37°C for 1 h. After incubation, the inoculum was removed, and the cells were treated with overlay media and incubated for 32 h (ZIKV, JEV, and WNV) or 72 h (PSFV, BJV, and DENV2). Fixation and primary antibody treatments were performed, as described in Section 2.4. After inoculation with HRP-conjugated anti-mouse IgG antibody (Thermo Fisher Scientific) as a secondary antibody at 37°C 1 h, virus-derived foci were stained with TrueBlue Peroxidase Substrate (Sera Care Life Sciences Inc., Milford, MA, USA) and counted. Neutralizing antibody titers were defined as the reciprocal of the

highest serum dilution showing a 50% reduction in the number of foci compared to non-immunized control serum (FRNT₅₀), as previously reported [19].

2.11. Antibody-dependent enhancement (ADE) of MBFV infection

Flow cytometry-based assays were performed to measure the ADE of MBFV infection, as previously reported [8]. Serial 10-fold dilutions of serum samples were incubated with MBFV (MOI of 0.1) for 1 h at 37°C, and 5.0×10^4 K562 cells in RPMI containing 10% FBS were added to each well in round-bottom 96-well plates (Corning). At 48 hpi (JEV and WNV) or 72 hpi (DENV2 and ZIKV), cells were fixed with 4% paraformaldehyde, and permeabilized with PBS containing 0.2% BSA and 0.05% saponin. Viral antigens were stained with 4G2 antibody, and Alexa Fluor 488-conjugated goat anti-mouse IgG antibody (Invitrogen). The virus-infected cells were counted using an FACSCanto (BD Biosciences, Franklin Lakes, NJ, USA) and analyzed using FlowJo software version 10.7.1 (Treestar Inc., Ashland, OR, USA). Data were expressed as overall ADE, calculating the area under the curve using GraphPad Prism software version 8.0 (GraphPad Software, San Diego, CA, USA).

2.12. Statistical analysis

All statistical analyses were performed using GraphPad Prism software version 8.0. The methods of statistical analysis are described in the Figure Legends for each experiment.

3. Results

3.1. The prME proteins of lineage IIa ISFVs are phylogenetically closely related to those of MBFVs

ML phylogenetic analysis based on whole amino acid sequences showed that both lineage IIa and IIb ISFVs (PSFV and BJV) were grouped within the cluster of MBFVs (Fig. 1A). The cluster of lineage I ISFVs was distinct from those of MBFVs and lineage II ISFVs, as previously reported [20], indicating that lineage II ISFVs have a greater amino acid sequence similarity to MBFVs compared to lineage I ISFVs.

In general, the amino acid sequences of the prME proteins were highly similar among flaviviruses [21,22]. We investigated the prME amino acid sequences of the MBFVs and lineage II ISFVs used in this study. The prME of DENV2, ZIKV, JEV, and WNV had 45-78% amino acid sequence identity and over 60% similarity to each other. The prMEs of PSFV and BJV also had 59–65% amino acid sequence similarity to those of DENV2, ZIKV, JEV, and WNV, which was comparable to the similarity among these

MBFVs (Table S1). The fusion loop domain in prME, which is known to be highly conserved in MBFVs, was conserved in both lineage II ISFVs (Fig. S1) [23]. We also performed ML phylogenetic analysis on the prME amino acid sequences of the flaviviruses. Lineage IIa ISFVs belonged to a clade of MBFVs, while lineage IIb ISFVs were associated with a different clade (Fig. 1B). This result suggested that the prME protein of lineage IIa ISFVs is more closely related to MBFVs than that of lineage IIb ISFVs.

3.2. Anti-PSFV sera cross-react with MBFV antigens with ADE activity

To evaluate the antigenic similarity between MBFV and lineage II ISFV, we performed serological analysis using antisera from mice subcutaneously immunized with purified PSFV from lineage IIa ISFV or BJV from lineage IIb ISFV.

The production of antibodies against PSFV or BJV in mice sera was evaluated using ELISAs of lineage II ISFV-infected cell lysates. All anti-PSFV or BJV sera ($n = 7$ each) reacted to cellular lysates infected with each lineage II ISFV, indicating that immunization with each lineage II ISFV *via* SC route induced anti-PSFV or BJV antibody in each serum (Figs. 2A, S2A and S2B). We evaluated the reactivity of antisera to cellular lysates infected with MBFVs, including DENV2, ZIKV, JEV, and WNV.

Pan-flavivirus 4G2 antibody was used as a positive control and reacted against all the lysates infected with MBFVs (Figs. S2C and S2D). There were no differences in non-specific binding to cell lysates of non-infected Vero cells between the anti-PSFV sera, anti-BJV sera, and non-immunized control sera (Fig. S2E). ELISA revealed that anti-PSFV sera possessed broad binding activity to cell lysates infected with DENV2 (positivity rate 6/7), ZIKV (positivity rate 6/7), JEV (positivity rate 1/7), and WNV (positivity rate 3/7) (Figs. 2B and S2F). On the other hand, anti-BJV sera had binding activity to DENV2 (positivity rate 4/7), ZIKV (positivity rate 3/7), and WNV (positivity rate 1/7) but not to JEV (positivity rate 0/7) (Figs. 2B and S2F).

Antibodies which recognize structural proteins such as prME enhance flavivirus infection by incorporating virus-bound antibodies *via* the Fcγ receptor [24,25]. This enhancement of antibody-dependent viral infection has been called antibody dependent enhancement (ADE). We found that anti-PSFV and BJV sera reacted to cell lysates infected with DENV2, ZIKV, JEV, or WNV, suggesting that anti-PSFV and BJV sera may elicit ADE. Therefore, we used flow cytometry-based ADE assays using Fcγ receptor-expressing K562 cells to evaluate the ADE activity of anti-PSFV or BJV sera. Enhancement of the number of K562 cells infected with DENV2, ZIKV, JEV, or WNV was observed in the presence of anti-PFSV sera (Fig. 3A), and the overall ADE activity

of anti-PSFV sera was significantly higher than those of control sera (Fig. 3B). In contrast, anti-BJV sera had only a slightly enhanced number of infected cells with each MBFV (Fig. 3A), and their overall ADE activity was not significantly higher than those of control sera (Fig. 3B). These results indicated that anti-PSFV sera potentially cross-reacted to DENV2, ZIKV, JEV, and WNV, accompanied by ADE activity. It has been reported that ADE is elicited by antibodies showing reactivity with structural proteins in flavivirus infection [10]. We confirmed that structural proteins of MBFVs were recognized by anti-PSFV or BJV sera using western blotting and found that both sera reacted with the prM and E proteins of MBFVs (Fig. S3).

3.3. Anti-lineage II ISFV sera had no or low neutralizing activity against MBFVs

We evaluated the neutralizing activity of lineage II ISFV antisera against the lineage II ISFV or MBFV infection. First, inhibition of PSFV or BJV infection in C6/36 cells by anti-PSFV or BJV sera was confirmed using FRNT (Figs. 4A and 4B). Some of those antisera neutralized not only their own virus infection but also other lineage II ISFV infections. Next, we confirmed that the positive controls of MBFV antisera obtained from MBFV-infected mice completely inhibited the foci formation of each virus in Vero cells in a serum dilution-dependent manner (Fig. 4C). In contrast,

anti-PSFV and BJV sera showed few or no foci reductions against DENV2, ZIKV, JEV, or WNV (Fig. 4C). The FRNT₅₀ titers of both anti-ISFV sera were significantly lower than those of positive control sera against all MBFVs (Fig. 4D). However, a few anti-PSFV sera suppressed foci formation to below 50% for DENV2 (positivity rate 1/7), ZIKV (positivity rate 1/7), WNV (positivity rate 1/7), and anti-BJV sera also decreased the foci formation of WNV (positivity rate 1/7). These results suggested that the epitopes recognized by anti-MBFV neutralizing antibodies might be present in PSFV and BJV.

4. Discussion

Previous studies using MBFV antisera showed that hemagglutination activities of lineage IIa ISFVs such as Aripo virus, La Tina virus, Marisma mosquito virus and lineage IIb ISFV such as Nanay virus were inhibited by various MBFV antisera [3,12], suggesting that these lineage II ISFVs antigens were cross-reactive with MBFV antisera. In this study, we evaluated cross-reactivities of both lineage IIa and IIb ISFVs antisera with MBFVs antigens. Our results showed that the lineage IIa ISFV PSFV antisera had higher positivity for binding to MBFV antigens and induced more ADE of MBFV infections than lineage IIb ISFV BJV antisera. These results suggest that PSFV is more

antigenically similar to MBFVs than BJV. We further found that both lineage II antisera recognized prM and E proteins of each MBFV (Fig. S3). Taken together, these observations suggest that lineage IIa PSFV had prME proteins more antigenically similar to those of DENV2, ZIKV, JEV, and WNV than lineage IIb BJV. This finding was consistent with the result of the ML phylogenetic analysis using prME protein sequences (Fig. 1B).

Our results and previous studies showed lineage II ISFV possessed antigenically similar viral proteins to MBFVs [3,12,13]. Furthermore, this study demonstrated the existence of antigenic differences between genetically closely related lineage IIa and IIb ISFVs by antibody-based analyses. It should be noted, however, we only used a single virus strain as representative of each lineage IIa or IIb ISFV in this study. To clarify the differences in the viral proteins of the lineage II ISFVs, further studies, such as structural analysis or reactive analysis using monoclonal antibodies with other virus strains, will be required.

Declaration of competing interest

Shinsuke Toba, Kentaro Uemura and Akihiko Sato are employees of Shionogi & Co., Ltd. Other authors declare no competing interests.

322

323 **Acknowledgments**

324 This work was supported by the Japan Agency for Medical Research and
325 Development (AMED) under grant number JP21wm0225018 (Y.O.), Japan Program for
326 Infectious Diseases Research and Infrastructure (AMED) under grant number
327 JP21wm0125008 (H.S.), by the Ministry of Education, Culture, Sports, Science and
328 Technology, Japan (MEXT) /Japan Society for the Promotion of Science (JSPS)
329 KAKENHI under the grant number JP20K21298 (Y.O.) and JP20J22269 (K.T.), by the
330 Japan Science and Technology Agency (JST) Moonshot R&D under the grant number
331 JPMJMS2025 (Y.O.). We would like to thank Enago (www.enago.jp) for English
332 language review.

333

- [1] J. Hobson-Peters, J.J. Harrison, D. Watterson, et al., A recombinant platform for flavivirus vaccines and diagnostics using chimeras of a new insect-specific virus, *Sci Transl Med* 11 (2019). 10.1126/scitranslmed.aax7888.
- [2] C.E. Wastika, H. Harima, M. Sasaki, et al., Discoveries of Exoribonuclease-Resistant Structures of Insect-Specific Flaviviruses Isolated in Zambia, *Viruses* 12 (2020). 10.3390/v12091017.
- [3] A.J. Auguste, R.M. Langsjoen, D.L. Porier, et al., Isolation of a novel insect-specific flavivirus with immunomodulatory effects in vertebrate systems, *Virology* 562 (2021) 50-62. 10.1016/j.virol.2021.07.004.
- [4] T.B.H. Piyasena, N.D. Newton, J. Hobson-Peters, et al., Chimeric viruses of the insect-specific flavivirus Palm Creek with structural proteins of vertebrate-infecting flaviviruses identify barriers to replication of insect-specific flaviviruses in vertebrate cells, *J Gen Virol* 100 (2019) 1580-1586. 10.1099/jgv.0.001326.
- [5] A.M.G. Colmant, J. Hobson-Peters, T.A.P. Slijkerman, et al., Insect-Specific Flavivirus Replication in Mammalian Cells Is Inhibited by Physiological Temperature and the Zinc-Finger Antiviral Protein, *Viruses* 13 (2021). 10.3390/v13040573.
- [6] C.J. Neufeldt, M. Cortese, E.G. Acosta, et al., Rewiring cellular networks by members of the Flaviviridae family, *Nat Rev Microbiol* 16 (2018) 125-142. 10.1038/nrmicro.2017.170.
- [7] W.D. Crill, G.J. Chang, Localization and characterization of flavivirus envelope glycoprotein cross-reactive epitopes, *J Virol* 78 (2004) 13975-13986. 10.1128/JVI.78.24.13975-13986.2004.
- [8] S.V. Bardina, P. Bunduc, S. Tripathi, et al., Enhancement of Zika virus pathogenesis by preexisting ant flavivirus immunity, *Science* 356 (2017) 175-180. 10.1126/science.aal4365.
- [9] H. Garg, R. Yeh, D.M. Watts, et al., Enhancement of Zika virus infection by antibodies from West Nile virus seropositive individuals with no history of clinical infection, *BMC Immunol* 22 (2021) 5. 10.1186/s12865-020-00389-2.
- [10] W. Dejnirattisai, A. Jumnainsong, N. Onsirisakul, et al., Cross-reacting antibodies enhance dengue virus infection in humans, *Science* 328 (2010) 745-748. 10.1126/science.1185181.
- [11] Y. Saito, M.L. Moi, N. Takeshita, et al., Japanese encephalitis vaccine-facilitated dengue virus infection-enhancement antibody in adults, *BMC Infect Dis* 16 (2016) 578. 10.1186/s12879-016-1873-8.
- [12] H. Guzman, M.A. Contreras-Gutierrez, A.P.A. Travassos da Rosa, et al., Characterization of Three New Insect-Specific Flaviviruses: Their Relationship to the Mosquito-Borne Flavivirus Pathogens, *Am J Trop Med Hyg* 98 (2018) 410-419. 10.4269/ajtmh.17-0350.
- [13] J.J. Harrison, J. Hobson-Peters, A.M.G. Colmant, et al., Antigenic Characterization of New Lineage II Insect-Specific Flaviviruses in Australian Mosquitoes and Identification of Host Restriction Factors, *mSphere* 5 (2020). 10.1128/mSphere.00095-20.

- [14] Y. Orba, K. Matsuno, R. Nakao, et al., Diverse mosquito-specific flaviviruses in the Bolivian Amazon basin, *J Gen Virol* 102 (2021). 10.1099/jgv.0.001518.
- [15] S. Kumar, G. Stecher, M. Li, et al., MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms, *Mol Biol Evol* 35 (2018) 1547-1549. 10.1093/molbev/msy096.
- [16] I. Letunic, P. Bork, Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation, *Nucleic Acids Res* 49 (2021) W293-W296. 10.1093/nar/gkab301.
- [17] D.C. Clark, M. Lobigs, E. Lee, et al., In situ reactions of monoclonal antibodies with a viable mutant of Murray Valley encephalitis virus reveal an absence of dimeric NS1 protein, *J Gen Virol* 88 (2007) 1175-1183. 10.1099/vir.0.82609-0.
- [18] M. Sasaki, P.D. Anindita, W. Phongphaew, et al., Development of a rapid and quantitative method for the analysis of viral entry and release using a NanoLuc luciferase complementation assay, *Virus Res* 243 (2018) 69-74. 10.1016/j.virusres.2017.10.015.
- [19] J.L. Slon Campos, M. Poggianella, S. Marchese, et al., DNA-immunisation with dengue virus E protein domains I/II, but not domain III, enhances Zika, West Nile and Yellow Fever virus infection, *PLoS One* 12 (2017) e0181734. 10.1371/journal.pone.0181734.
- [20] B.J. Blitvich, A.E. Firth, Insect-specific flaviviruses: a systematic review of their discovery, host range, mode of transmission, superinfection exclusion potential and genomic organization, *Viruses* 7 (2015) 1927-1959. 10.3390/v7041927.
- [21] T.C. Pierson, M.S. Diamond, The continued threat of emerging flaviviruses, *Nat Microbiol* 5 (2020) 796-812. 10.1038/s41564-020-0714-0.
- [22] E. Krol, G. Brzuska, B. Szewczyk, Production and Biomedical Application of Flavivirus-like Particles, *Trends Biotechnol* 37 (2019) 1202-1216. 10.1016/j.tibtech.2019.03.013.
- [23] B.S. Berneck, A. Rockstroh, J. Fertey, et al., A Recombinant Zika Virus Envelope Protein with Mutations in the Conserved Fusion Loop Leads to Reduced Antibody Cross-Reactivity upon Vaccination, *Vaccines (Basel)* 8 (2020). 10.3390/vaccines8040603.
- [24] L. Priyamvada, K.M. Quicke, W.H. Hudson, et al., Human antibody responses after dengue virus infection are highly cross-reactive to Zika virus, *Proc Natl Acad Sci U S A* 113 (2016) 7852-7857. 10.1073/pnas.1607931113.
- [25] A.P. Goncalvez, R.E. Engle, M. St Claire, et al., Monoclonal antibody-mediated enhancement of dengue virus infection in vitro and in vivo and strategies for prevention, *Proc Natl Acad Sci U S A* 104 (2007) 9422-9427. 10.1073/pnas.0703498104.



Figure Legends

Fig. 1 Phylogenetic analysis of insect-specific flaviviruses (ISFVs) based on amino acid sequences. Pathogenic mosquito-borne flaviviruses (MBFVs), including Dengue virus type 2 (DENV2), Zika virus (ZIKV), Japanese encephalitis virus (JEV) and West Nile virus (WNV) used in this study are highlighted in the black squares. Lineage II ISFVs, Psorophora flavivirus (PSFV) and Barkedji virus (BJV), are highlighted in the dot-lined square. Phylogenetic trees were generated based on (A) whole amino acid sequence and (B) precursor membrane and envelope protein (prME) by the maximum-likelihood method using best-fit models with bootstrap values of 1,000 replicates.


Fig. 2 Binding activity of anti-PSFV or BJV sera to cell lysates infected with each virus. Binding activity of anti-PSFV or BJV sera to cell lysates infected with (A) lineage II ISFVs and (B) MBFVs was evaluated using enzyme-linked immunosorbent assay (ELISA). The dotted line indicates the limit of detection (LOD). The values in the graphs are expressed as the mean $\pm$ SD of seven serum samples ($n = 7$).

Fig. 3 Antibody-dependent enhancement (ADE) of MBFV infection by anti-PSFV

or BJV sera. (A) Non-immunized control sera, anti-PSFV sera and anti-BJV sera were evaluated for their enhancement of MBFV infections in K562 cells. (B) Overall ADE activity was determined from the area under the curve (AUC) shown in Fig. 3A. The values in the graphs are expressed as the mean $\pm$ SD of seven serum samples ($n = 7$). **: $p < 0.005$, ***: $p < 0.001$ from one-way ANOVA with Dunn's multiple comparison test.

Fig. 4 Neutralizing activities of anti-PSFV or BJV sera against MBFVs. (A) Foci reduction neutralization tests (FRNTs) against PSFV or BJV were performed using C6/36 cells. (B) The FRNT₅₀ titer was determined from the data shown in Fig. 4A. *: $p < 0.01$ using t -tests with Mann-Whitney tests. (C) FRNT against MBFVs using Vero cells. (D) The FRNT₅₀ titer was determined from data shown in Fig. 4C. Dotted lines indicate 50% foci reduction (A and C) and detection limits (B and D). The values in the graphs are expressed as the mean $\pm$ SD of seven serum samples ($n = 7$). *: $p < 0.01$, **: $p < 0.005$ using one-way ANOVA with Dunn's multiple comparison test.

A



B





Fig. 2

A



B




Fig. 3




Fig. 4