



Title	Lead concentrations and isotope ratios in blood, breastmilk and feces : contribution of both lactation and soil/dust exposure to infants in a lead
Author(s)	Toyomaki, Haruya; Yabe, John; Nakayama, Shouta M. M. et al.
Citation	Environmental pollution, 286, 117456 <a href="https://doi.org/10.1016/j.envpol.2021.117456">https://doi.org/10.1016/j.envpol.2021.117456</a>
Issue Date	2021-10-01
Doc URL	<a href="https://hdl.handle.net/2115/90474">https://hdl.handle.net/2115/90474</a>
Rights	© 2021. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Rights(URL)	<a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Type	journal article
File Information	Manuscript 20210518 Without Tracking.pdf



1 **Lead concentrations and isotope ratios in blood, breastmilk and feces:**  
2 **contribution of both lactation and soil/dust exposure to infants in a lead mining**  
3 **area, Kabwe, Zambia**

4

5 Haruya Toyomaki<sup>1</sup>, John Yabe<sup>2, 8</sup>, Shouta M.M. Nakayama<sup>1\*</sup>, Yared B. Yohannes<sup>1,3</sup>,  
6 Kaampwe Muzandu<sup>1,2</sup>, Tiza Mufune<sup>4</sup>, Hokuto Nakata<sup>1</sup>, Yoshinori Ikenaka<sup>1,5,9,10</sup>,  
7 Takeshi Kuritani<sup>6</sup>, Mitsuhiro Nakagawa<sup>6</sup>, Kennedy Choongo<sup>2, 7</sup> and Mayumi Ishizuka<sup>1</sup>

8 (\*Corresponding author)

9

10 1) Laboratory of Toxicology, Department of Environmental Veterinary Sciences,  
11 Faculty of Veterinary Medicine, Hokkaido University, Japan

12 2) The University of Zambia, School of Veterinary Medicine, Zambia

13 3) Department of Chemistry, College of Natural and Computational Science, University  
14 of Gondar, Ethiopia

15 4) Ministry of Health, District Health Office, Kabwe, Zambia

16 5) Water Research Group, School of Environmental Sciences and Development, North-  
17 West University, South Africa

18 6) Department of Earth and Planetary Sciences, Graduate School of Science, Hokkaido  
19 University, Sapporo, Japan

20 7) Fiji National University, College of Agriculture, Fisheries & Forestry, School of  
21 Animal and Veterinary Sciences, Koronivia Campus, Suva, Fiji

22 8) Department of Pathobiology, School of Veterinary Medicine, University of Namibia,  
23 Windhoek, Namibia

24 9) Translational Research Unit, Veterinary Teaching Hospital, Faculty of Veterinary  
25 Medicine, Hokkaido University, Sapporo, 060-0818, Japan

26 10) One Health Research Center, Hokkaido University, Japan

27

28

29

30

31 \*Address Correspondence to

32 Shouta M.M. Nakayama

33 Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Faculty  
34 of Veterinary Medicine, Hokkaido University, Kita 18, Nishi 9, Kita-ku, Sapporo 060-  
35 0818, Japan.

36 Tel: +81-11-706-5105, Fax: +81-11-706-5105.

37 E-mail: [shouta-nakayama@vetmed.hokudai.ac.jp](mailto:shouta-nakayama@vetmed.hokudai.ac.jp)

38 or [shoutanakayama0219@gmail.com](mailto:shoutanakayama0219@gmail.com)

39

40

41 **Abstract**

42           Lead (Pb) poses a serious public health concern. Breastmilk may be a possible  
43 source of Pb exposure in infants, as Pb can be transferred from the maternal blood to  
44 breastmilk. The present study was undertaken to determine the Pb exposure and the  
45 contribution of lactation as one of the exposure pathways to infants in a Pb mining area,  
46 Kabwe, Zambia. Blood, breastmilk and infants' feces were collected from 418 pairs of  
47 infants and mothers. The Pb concentrations, isotope ratios in the samples, and  
48 biochemistry in mothers' plasma were analyzed. The overall mean of blood lead levels  
49 (BLLs) in infants and mothers were 18.0 and 11.3  $\mu\text{g/dL}$ , respectively. High Pb  
50 concentration in breastmilk (range: 0.4–51.9, mean: 5.3  $\mu\text{g/L}$ ) above the WHO  
51 acceptable level between 2 and 5  $\mu\text{g/L}$  were found and could be one of the sources of Pb  
52 exposure in infants. The Pb isotope ratios in infants' feces were the most similar to Pb  
53 ratios in the soil samples. The results suggest that infants are also exposed to Pb from  
54 the environment. Pb exposure in infants through breastfeeding and soil ingestion could  
55 potentially exceed daily intake of Pb which causes neurodevelopmental toxicity. In  
56 contrast to the high BLLs in mothers, the plasma biochemical profiles of most analyzed  
57 parameters were interestingly within, or close to, the standard reference values. Our data  
58 suggest that environmental remediation is urgently needed to reduce the Pb exposure in  
59 infants and mothers from the environment in Kabwe in parallel with chelation therapy.

60

61 **Keywords:** Lead poisoning, Infant, Mother, Breastmilk, Lead stable isotope

62

## 63 **1. Introduction**

64           Lead (Pb) poses a serious public health concern, accounting for 0.6% of the  
65 global burden of disease (WHO, 2010). Serious cases of Pb exposure have been  
66 reported in both developed and developing countries (Ajumobi et al., 2014; Haefliger et  
67 al., 2009; Ruckart et al., 2019). Lead poisoning causes various symptoms, including  
68 anemia, nephropathy, and death (Meyer et al., 2008). Children, especially infants are  
69 more vulnerable to Pb, compared to adults. To measure exposure to Pb, blood lead level  
70 (BLL) has been widely used. In the blood, Pb has a short half-life of 30–40 days  
71 (Barbosa et al., 2005). Even at low levels, Pb exposure can cause pediatric  
72 neurodevelopmental impairments, such as a reduction in intelligence quotient (IQ)  
73 (Canfield et al., 2003). Due to this, the blood Pb reference value for Pb exposure has  
74 been set to 5 µg/dL (CDC, 2019, 2012). A BLL above 45 µg/dL is considered the level  
75 where treatment is required (CDC, 2012, 2002; Needleman, 2004), and a BLL above  
76 100 µg/dL is considered a fatal level in children, which causes serious clinical  
77 symptoms such as encephalopathy, even in adults (Meyer et al., 2008; NAS, 1972). The  
78 European Food Safety Authority (EFSA) observed that Pb dietary intake of 0.5 µg/kg  
79 body weight (bw)/day would be associated with developmental neurotoxicity in young  
80 children (EFSA, 2010).

81           Breastmilk is vital for infants, to ensure normal development and to prevent  
82 infectious diseases. However, breastmilk may be a possible source of Pb exposure in  
83 infants, as Pb can be transferred from the maternal blood to breastmilk. As a result,  
84 BLLs in mothers should be monitored to prevent Pb exposure in infants via  
85 breastfeeding. Therefore, to minimize Pb exposure in infants through breastmilk, WHO  
86 (1989) has set the acceptable level of breastmilk Pb concentration to be between 2 and 5

87  $\mu\text{g/L}$ . Mothers with confirmed BLLs above 40  $\mu\text{g/dL}$  should pump and discard their  
88 breastmilk (CDC, 2010).

89         Lead exposure primarily occurs via ingestion and inhalation, and can be traced  
90 to numerous sources, including battery recycling, gasoline, paint, as well as mining  
91 (Calabrese and Stanek, 1995; Meyer et al., 2008; Schoning et al., 1996; Yabe et al.,  
92 2010). Identifying the source of Pb exposure is important to prevent exposure. One such  
93 method for identifying sources is the use of Pb isotopic tracing (Komarek et al., 2008;  
94 Gulson, 2008). Lead is present in the environment as four main isotopes:  $^{208}\text{Pb}$  (52%),  
95  $^{207}\text{Pb}$  (23%),  $^{206}\text{Pb}$  (24%), and  $^{204}\text{Pb}$  (1%) (Komárek et al., 2008). The compositions of  
96 these isotopes are not affected to a measurable extent by physicochemical fractionation  
97 processes (Bollhöfer and Rosman, 2001; Veysseyre et al., 2001).

98         The Zambian town of Kabwe accommodates a Pb-zinc (Zn) mining area which,  
99 up until its closure in 1994, was operated without adequate pollution laws to regulate  
100 mining emissions. Elevated BLLs and Pb concentrations in the feces and urine of  
101 children near the mine have been reported, all of which exceeded the 5  $\mu\text{g/dL}$  blood Pb  
102 reference value (Yabe et al., 2018, 2015). Yabe et al. (2015) found that BLLs in  
103 children between the ages of one and two years old were higher compared with those in  
104 children between the ages of four and seven years old in Kabwe, as has been observed  
105 in many other studies. It is necessary to reduce and prevent Pb exposure in children.  
106 This is especially important in infants, who are more vulnerable to Pb poisoning.  
107 Moreover, a more recent study has revealed a high Pb exposure also in mothers, where  
108 approximately 5% of mothers were found to have a BLL above 45  $\mu\text{g/dL}$  which  
109 indicated that treatment was required (Yabe et al., 2020, Nakata et al., 2021). The  
110 breastfeeding practices of mothers with high BLLs are a possible source of Pb exposure

111 for infants in Kabwe. However, the precise sources and routes of Pb exposure in infants  
112 have not yet been determined. Furthermore, no clinical studies of Pb poisoning have  
113 been done in Kabwe, despite high BLLs being reported in the local people. Some  
114 previous studies have reported that metallothionein concentrations, which is a cysteine-  
115 rich protein that binds and detoxifies toxic metals, increase as metal concentrations in  
116 the blood increase (Bizoń and Milnerowicz, 2014; Kowalska et al., 2015).  
117 Metallothionein may therefore play an important role in reducing Pb toxicity in the  
118 people of Kabwe.

119 The current study aimed to determine the Pb exposure and the contribution of  
120 lactation as one of the exposure pathways to infants in a lead mining area, Kabwe,  
121 Zambia. Pb concentrations in mothers' breastmilk, infants' feces, and both infants' and  
122 mothers' blood, were analyzed. Daily intake of Pb in infants through breastfeeding and  
123 soil ingestion was calculated to estimate the burden of routes of Pb exposure. The Pb  
124 isotope ratios in samples were analyzed in a limited number of samples to determine the  
125 source of Pb exposure. Furthermore, a plasma biochemical analysis including  
126 metallothionein concentrations was conducted in the mothers to evaluate the health  
127 impact of Pb exposure in this population.

128


## 129 **2. Materials and Methods**

### 130 **2.1 Sampling Sites**

131 The town of Kabwe is located at about 28°26'E and 14°27'S, and is the  
132 provincial capital of Zambia's Central Province. It is the fourth largest town in Zambia,  
133 with a population of about 230,000 inhabitants and an area of 1547 km<sup>2</sup>. In Kabwe,  
134 metallic residues from abandoned tailings and waste stored in the mine have continued

135 to serve as potential sources of metal pollution even after the closure of the mine. Dust  
136 emanates from the mine dumps, and residents in townships close to the mine may be  
137 exposed to high levels of Pb in contaminated dust and soil.

138 The present study was conducted at health centers in four sites near the mine  
139 (Kasanda, Makululu, Chowa and Katondo) and one site far from the mine (Bwacha) in  
140 the rainy season from January to March of 2017 (Fig. 1). Kasanda, Makululu, Chowa,  
141 Katondo, and Bwacha health centers were located about 0.9, 2.6, 1.4, 4.5, and 6.1 km  
142 from the mine, respectively. Kasanda and Makululu are located on the western side of  
143 the mine and in the direction of prevailing winds.



144  
145 Fig. 1. Map of the sampling sites in Kabwe, Zambia (image modified from Google  
146 Earth).

147

148 **2.2 Sampling**

149 This study was approved by the University of Zambia Research Ethics  
150 Committee (UNZAREC; REF. No. 012-04-16). Further approvals were granted by the  
151 Ministry of Health through the Zambia National Health Research Ethics Board and the  
152 Kabwe District Health Office.

153 A sensitization campaign about the research activities was conducted by  
154 community health workers before sampling in their catchment areas around the health  
155 centers. Mothers and guardians were encouraged to participate in the study, and were  
156 asked to take their breastfed infants under the age of 1 year and 6 months to the selected  
157 health centers for sample collection. Only infants with mothers/guardians that willingly  
158 agreed to participate and signed the informed consent were included in the present study.  
159 After informed and written consent were obtained from the mothers/guardians, blood  
160 samples were collected as described by Yabe et al. (2015). The mothers/guardians were  
161 also interviewed to obtain necessary personal details about themselves and their infants,  
162 such as age and sex. Sample collection and questionnaire administration were  
163 undertaken by certified local nurses. In accordance with ethical requirements,  
164 confidentiality was upheld in the study.

165 Blood samples up to 2 mL and 5 mL were collected from the cephalic veins of  
166 each infant and mother, respectively, and were placed into heparinized blood collection  
167 tubes. Breastmilk samples from mothers were collected in clean sample cups by gentle  
168 compression of the breast and transferred to 2 mL sample tubes for storage and  
169 transportation. To avoid contamination, the hands as well as venipuncture and breast  
170 sites were cleaned and sanitized with an ethanol swab before the sample collection.  
171 Plasma samples were separated only from the mothers' blood after centrifugation. For

172 infants' fecal samples, mothers/guardians were handed 30 mL stool containers equipped  
173 with scoops and were instructed to scoop feces into the container from a used diaper in  
174 the morning of the following day. Household soil samples were collected in June 2016  
175 from Kasanda (n = 12) and Makululu (n = 20) as a reference of environmental samples  
176 for Pb stable isotope analysis.

177 The processed samples were transported to the laboratory of The University of  
178 Zambia, School of Veterinary Medicine, Zambia, and stored at -20 °C. The material  
179 transfer agreement (MTA) for human samples from the Zambia National Health  
180 Research Ethics Committee (approval No. E00417) was obtained before transportation.  
181 Similarly, the phytosanitary certificate from plant quarantine and phytosanitary service,  
182 Zambia Ministry of Agriculture, and import permission by plant protection station,  
183 Japanese Ministry of Agriculture, Forestry and Fisheries (approval No. 28-313) was  
184 also granted for soil samples. The human samples were transported in temperature-  
185 controlled boxes with ice packs, and the soil samples in temperature-controlled boxes,  
186 for further analysis at Hokkaido University, Japan.

187

### 188 **2.3 Pb and Metal Concentration Analysis**

189 Pb and other metals (iron (Fe), copper (Cu), Zn, and silver (Ag)) were extracted  
190 from the samples. Thawed fecal and bulk soil samples were weighed on heat-resistant  
191 tissue drying plates and dried for 48 h in a tissue drying oven at 60 °C, whereas whole  
192 blood and breastmilk samples were only thawed. Each blood and breastmilk sample  
193 (1mL) and 50 mg of each dried fecal and soil sample were analyzed. Detailed method  
194 was described in the supplementary materials.

195

196 **2.4 Calculation of Daily Intake of Pb in Infants through Breastfeeding and Soil**  
197 **Ingestion**

198 Daily intake of Pb in infants through breastfeeding and soil ingestion was  
199 calculated. The calculation was conducted using the maximum, mean, and minimum Pb  
200 concentrations in breastmilk and soil samples in the present study using the formulas  
201 below. The amounts of daily breastmilk intake and soil ingestion in infants were set as  
202 0.78 L/day and 30 mg/day reported by Costa et al. (2010) and United States  
203 Environmental Protection Agency (US EPA, 2011), respectively.

204

205 *Daily intake of Pb = (Pb concentrations in breastmilk × Amount of daily breastmilk*  
206 *intake) + (Pb concentrations in soil × amount of daily soil ingestion)*

207

208 In the present study, 0.5 µg/kg bw/day was used as the reference value as the  
209 limit of daily intake of Pb for infants (EFSA, 2010).

210

211 **2.5 Stable Pb Isotope Analysis**

212 Only 26 sample sets with high Pb concentrations of infants' and mothers' blood,  
213 breastmilk, and infants' feces from Kasanda and Makululu, were chosen and analyzed  
214 during Pb isotope analysis as well as 32 soil samples. As mentioned above, these soil  
215 samples were collected at the preliminary survey in 2016. The sampling area between  
216 blood and soil are same, but not all the locations were exactly from the same household.  
217 The sample dissolution procedure was similar to previously described methods  
218 (Kuritani and Nakamura, 2002; Nakayama et al., 2019). The detailed methods of the  
219 high precision isotope analysis were described in the supplementary materials. We have

220 measured duplicates of breastmilk samples and the results of Pb isotope ratios had the  
221 similar trends. As sample amounts decrease, the uncertainty of ion beam intensity  
222 measurements for the minor isotope  $^{204}\text{Pb}$  tends to increase, and the accuracy and  
223 precision of the isotopic ratios involving  $^{204}\text{Pb}$  decrease ( $^{204}\text{Pb}$  error; Hamelin et al.,  
224 1985). Therefore, the isotopic composition of Pb is commonly expressed as ratios of  
225  $^{208}\text{Pb}$ ,  $^{207}\text{Pb}$ , and  $^{206}\text{Pb}$ . However, since normalization to  $^{204}\text{Pb}$  yields the largest  
226 variability between reservoirs (Komárek et al., 2008), we included  $^{204}\text{Pb}$  data with  
227 correction (Kuritani et al., 2003, 2002). This was done in order to observe the detail  
228 variability of isotope ratio among the breastmilk, blood, feces and soil samples in this  
229 study.

230

## 231 **2.6 Plasma Biochemical Analysis and Metallothionein ELISA**

232 A conventional blood biochemical analyzer (FUJI DRICHEM 7000V;  
233 FUJIFILM corporation, Tokyo, Japan) was used to analyze the concentrations of  
234 alanine aminotransferase (ALT), alkaline phosphatase (ALP), aspartate  
235 aminotransferase (AST), gamma glutamyl transpeptidase (GGT), lactase dehydrogenase  
236 (LDH), total bilirubin (T-Bil), total protein (TP), albumin (Alb), blood urea nitrogen  
237 (BUN), creatinine (Cre), and urea acid (UA) in mothers' plasma samples.  
238 Metallothionein in mothers' plasma samples was measured by ELISA using an antibody  
239 against iso-Metallothionein I and II (Metallothionein ELISA kit; Frontier Institute Co.,  
240 Ltd., Hokkaido, Japan). The standard reference ranges for each parameter in humans  
241 were provided by the kit manufacturers.

242

243 **2.7 Statistical Analysis**

244 All data from the experiments and questionnaires were combined into a single  
245 electronic database and checked for accuracy and outliers. All statistical analyses were  
246 performed at a significance level of  $p < 0.05$  using JMP 13.1.0 (SAS Institute, USA).  
247 Mean values were indicated in addition to standard deviation (SD) values. The  
248 collinearity between factors was analyzed using Spearman's rank correlation test. A  
249 Steel–Dwass multiple comparisons test was used to compare the differences between  
250 the factors among areas and samples.

251

## 252 3. Results

### 253 3.1 Characteristics of the Infants and Mothers

254 A total of 418 pairs of infants and mothers participated in this study. Of these,  
255 333 participants came from four sites near the mine, and 85 came from Bwacha, which  
256 is located about 6 km from the mine (Table 1; Fig. 1). None of the infants and mothers  
257 had overt signs of Pb poisoning. Infants from Chowa were younger than those from  
258 other areas, except for Bwacha ( $p < 0.05$ ). In regard to height, infants from Chowa were  
259 significantly shorter and infants from Katondo were significantly taller than infants  
260 from other areas ( $p < 0.05$ ). Body weight of the infants from Chowa was significantly  
261 lower than that of the infants from other areas ( $p < 0.05$ ).

262 The overall height of boys was greater than that of girls overall ( $p < 0.01$ ), with  
263 boys found to be significantly taller in the four areas near the mine considered together  
264 ( $p < 0.01$ ) and in Makululu alone ( $p < 0.01$ , Supplementary Table S3). In Katondo, the  
265 height of boys tended to be greater than that of girls, but no significant difference was  
266 recorded ( $p = 0.09$ ). The weight of boys was significantly higher than that of girls when  
267 considering all infants ( $p < 0.01$ ), and for infants from near the mine ( $p < 0.01$ ), and  
268 infants in Kasanda ( $p < 0.05$ ), Makululu ( $p < 0.01$ ), Katondo ( $p < 0.01$ ), and Bwacha ( $p$   
269  $<$  0.045) alone.

270 Table 1. General characteristics of infants and mothers, as well as Pb concentrations in blood, breastmilk and fecal samples in Kabwe,  
 271 Zambia; mean  $\pm$  SD values (sample size, minimum–maximum).

Area (N)	Sex of infants, boy:girl	Age of infants, months	Height, Cm	Weight, kg	Age of mothers, years	BLLs in infants, $\mu\text{g/dL}$	BLLs in mothers, $\mu\text{g/dL}$	Pb in breastmilk, $\mu\text{g/L}$	Pb in infants' feces, mg/kg dry weight
Overall (418)	221:197	7.1 $\pm$ 3.8 (417, 0.1–16.8)	64.2 $\pm$ 7.7 (361, 45.0–95.0)	7.3 $\pm$ 1.7 (411, 1.9–12.1)	26.1 $\pm$ 6.5 (412, 16.3–46.1)	18.0 $\pm$ 18.1 <sup>A</sup> (406, 0.8–93.4)	11.3 $\pm$ 9.2 <sup>B</sup> (417, 1.5–82.6)	5.3 $\pm$ 7.0 <sup>C</sup> (407, 0.4–51.9)	39.2 $\pm$ 217.7 <sup>D</sup> (212, 0.08–3002.7)
Near the mine (333)	171:162	7.2 $\pm$ 4.0 (332, 0.1–16.8)	64.5 $\pm$ 8.0 (280, 45.0–95.0)	7.2 $\pm$ 1.7 (329, 1.9–12.1)	26.1 $\pm$ 6.6 (329, 16.3–46.1)	21.4 $\pm$ 18.9 <sup>**A</sup> (321, 1.6–93.4)	13.0 $\pm$ 9.5 <sup>**B</sup> (333, 1.9–82.6)	6.1 $\pm$ 7.5 <sup>**C</sup> (324, 0.4–51.9)	49.2 $\pm$ 248.0 <sup>**D</sup> (162, 0.09–3002.7)
Kasanda (82)	43:39	7.3 $\pm$ 3.8 <sup>a</sup> (82, 0.6–16.7)	64.5 $\pm$ 8.4 <sup>a</sup> (74, 49.0–95.0)	7.6 $\pm$ 1.5 <sup>a</sup> (79, 4.0–12.1)	27.1 $\pm$ 6.7 (82, 17.4–45.4)	24.8 $\pm$ 20.9 <sup>a,A</sup> (80, 3.7–93.4)	15.8 $\pm$ 10.6 <sup>a,B</sup> (82, 2.3–82.6)	9.2 $\pm$ 8.9 <sup>a,C</sup> (79, 1.9–50.4)	38.8 $\pm$ 88.9 <sup>ab,D</sup> (28, 0.6–451.0)
Makululu (102)	44:58	7.7 $\pm$ 3.7 <sup>a</sup> (102, 1.5–16.7)	63.0 $\pm$ 7.0 <sup>a</sup> (102, 45.0–85.0)	7.4 $\pm$ 1.4 <sup>a</sup> (101, 4.1–11.9)	26.2 $\pm$ 7.0 (102, 16.6–45.9)	30.8 $\pm$ 19.4 <sup>b,A</sup> (102, 5.6–82.7)	17.6 $\pm$ 10.1 <sup>a,B</sup> (102, 3.3–67.3)	7.1 $\pm$ 7.1 <sup>ab,C</sup> (100, 1.1–40.2)	82.4 $\pm$ 362.7 <sup>a,D</sup> (69, 1.1–3002.7)
Chowa (58)	35:23	5.3 $\pm$ 3.9 <sup>b</sup> (58, 0.1–15.2)	56.2 $\pm$ 7.1 <sup>b</sup> (13, 49.0–67.0)	5.6 $\pm$ 1.4 <sup>b</sup> (58, 3.5–8.7)	26.9 $\pm$ 6.3 (58, 17.6–41.8)	15.7 $\pm$ 15.5 <sup>c,A</sup> (49, 3.2–62.3)	11.2 $\pm$ 6.1 <sup>b,A</sup> (58, 3.0–36.2)	5.6 $\pm$ 6.5 <sup>b,B</sup> (57, 1.4–42.1)	33.7 $\pm$ 139.4 <sup>bc,C</sup> (31, 0.09–780.4)
Katondo (91)	49:42	7.7 $\pm$ 4.3 <sup>a</sup> (90, 0.9–16.8)	67.5 $\pm$ 7.6 <sup>c</sup> (91, 48.0–94.0)	7.8 $\pm$ 1.7 <sup>a</sup> (91, 1.9–11.1)	24.6 $\pm$ 5.9 (87, 16.3–46.1)	10.9 $\pm$ 9.9 <sup>c,A</sup> (90, 1.6–51.0)	6.3 $\pm$ 3.5 <sup>c,B</sup> (91, 1.9–21.6)	2.3 $\pm$ 5.5 <sup>c,C</sup> (88, 0.4–51.9)	4.4 $\pm$ 5.1 <sup>c,D</sup> (34, 0.1–19.4)
Bwacha (85)	50:35	6.9 $\pm$ 3.2 <sup>ab</sup> (85, 1.6–13.5)	63.0 $\pm$ 6.6 <sup>a</sup> (81, 51.0–86.0)	7.6 $\pm$ 1.6 <sup>a</sup> (82, 4.6–11.4)	26.2 $\pm$ 5.9 (83, 16.6–43.3)	5.2 $\pm$ 4.1 <sup>d,A</sup> (85, 0.8–22.0)	4.7 $\pm$ 3.4 <sup>d,A</sup> (84, 1.5–23.1)	2.3 $\pm$ 2.9 <sup>c,B</sup> (83, 0.5–17.9)	6.9 $\pm$ 27.6 <sup>d,C</sup> (50, 0.08–184.0)

272 Note: \*\* indicates a significant difference ( $p < 0.01$ ) between sites near the mine and Bwacha. Various small letters indicate a significant  
 273 difference among areas ( $p < 0.05$ ). Various capital letters indicate a significant difference among infants' and mothers' blood, breastmilk,  
 274 and infants' feces ( $p < 0.05$ ).

275 **3.2 Pb and Other Metals Concentrations in Blood, Breastmilk, Infants' Feces, and**  
276 **Soil.**

277 The overall mean of Pb concentrations in the soil samples ( $n = 32$ ) was  $1048 \pm$   
278  $1470$  mg/kg (dry weight) and ranged from 346 to 6327 mg/kg.


279 The overall mean values of BLLs in infants and mothers were  $18.0 \pm 18.1$  and  
280  $11.3 \pm 9.2$   $\mu\text{g/dL}$ , respectively (Table 1 and Fig. 2a). We found 76.8% of infants  
281 (312/406) and 73.6% of mothers (307/417) had BLLs above the reference value for Pb  
282 exposure (5  $\mu\text{g/dL}$ ; CDC, 2019, 2012). Moreover, BLLs in 8.9% of infants (36/406) and  
283 1.2% of mothers (5/417) were above 45  $\mu\text{g/dL}$ , the recommended threshold BLL for  
284 chelation therapy (Meyer, 2008). No infants or mothers had BLLs above the lethal level  
285 for Pb exposure (100  $\mu\text{g/dL}$ ), however, the highest BLL in an infant in the present study  
286 was 93.4  $\mu\text{g/dL}$ . The overall mean of Pb concentrations in breastmilk and infants' feces  
287 were  $5.3 \pm 7.0$   $\mu\text{g/L}$  and  $39.2 \pm 217.7$  mg/kg (dry weight), respectively (Fig. 2b and 2c).  
288 Overall, 30.0% of breastmilk samples (122/407) had Pb concentrations of more than 5  
289  $\mu\text{g/L}$ , which is above the accepted level for breastfeeding (WHO 1989).

290 There were significant differences in Pb concentrations among sample types in  
291 the samples from all sites: infants' feces > infants' blood > mothers' blood > breastmilk  
292 ( $p < 0.05$ ). Among the samples from Chowa and Bwacha, there were no significant  
293 differences in Pb concentrations between infants' and mothers' blood. Pb concentrations  
294 in infants' blood were  $1.8 \pm 1.5$  and  $60.2 \pm 67.6$  times higher than those in the mothers'  
295 blood and Pb concentrations in breastmilk, respectively (Supplementary Table S4). On  
296 the other hand, Pb concentrations in infants' blood were  $5.3 \pm 7.4\%$  of the Pb  
297 concentrations in infants' feces.


298 Pb concentrations in infants' and mothers' blood, breastmilk, and infants' feces  
299 from sites near the mine (Kasanda, Makululu, Chowa, and Katondo) were significantly  
300 higher than the concentrations in samples from Bwacha. Among sites, the mean of Pb  
301 concentrations in each sample type from Makululu were the highest, except in  
302 breastmilk. Pb concentrations in samples from Bwacha were significantly lower than  
303 those from other sites ( $p < 0.05$ ), except in breastmilk.

304 BLLs in boys were significantly higher than those in girls ( $p = 0.04$ ) in  
305 Makululu, and higher in Bwacha ( $p = 0.06$ , Supplementary Table S5). Pb concentrations  
306 in infants' feces of boys were significantly higher than those of girls in Chowa ( $p <$   
307  $0.01$ ). The same trend was found in all infants, although the association was not  
308 statistically significant ( $p = 0.08$ ).


309 Supplementary Tables S6, S7, S8, and S9 show Fe, Cu, Zn, and Ag  
310 concentrations in the samples, respectively. Fe concentrations in infants' blood and Fe,  
311 Cu, Zn, and Ag concentrations in mothers' blood from sites near the mine were  
312 significantly lower than the concentrations in samples from Bwacha. On the other hand,  
313 Fe concentrations in breastmilk from sites near the mine were significantly higher than  
314 the concentrations in samples from Bwacha.



315



316



317

318 Fig. 2. BLLs in infants and mothers (2a), Pb concentrations in breastmilk (2b), and  
 319 infants' feces (2c) among areas. The blue, red, orange, violet, and green colors indicate  
 320 samples from Kasanda, Makululu, Chowa, Katondo, and Bwacha, respectively. Solid  
 321 and dot box plots in Fig. 2a indicate BLLs in infants and mothers, respectively. Red and  
 322 black lines in Fig. 2a indicate 45 and 5  $\mu\text{g/dL}$ , respectively. The red dotted line in Fig  
 323 2b indicates the 5  $\mu\text{g/L}$  level.

324

325

### 326 3.3 Relationships among Samples and Factors

327 In the samples from all areas, significant positive correlations existed among  
 328 infants in the factors of age, height, and weight (Table 2,  $p < 0.001$ ). There were  
 329 significant positive correlations of Pb concentrations among all sample types ( $p < 0.001$ ,  
 330 Supplementary Fig. S2). BLLs in infants and Pb concentrations in infants' feces had a

331 significant positive correlation with the age (Supplementary Fig. S3), height, and weight  
332 of infants ( $p < 0.001$ ).

333 In all samples from sites near the mine, the same trend was found  
334 (Supplementary Table S10). Moreover, the height of infants had a significant negative  
335 correlation with BLLs in mothers ( $p < 0.01$ ,  $\rho = -0.19$ ) and Pb concentrations in  
336 breastmilk ( $p < 0.05$ ,  $\rho = -0.13$ ). In the samples from Bwacha, the same trend was  
337 found as in the samples from all other sites, except for the relationship between Pb  
338 concentrations in breastmilk and other samples, and the BLLs in mothers and Pb  
339 concentrations in infants' feces (Supplementary Table S11). Supplementary Tables S12,  
340 13, and 14 show the relationships among Pb and other metals in the samples from all  
341 areas, from sites near the mine, and Bwacha, respectively. In the samples from all areas,  
342 Pb concentrations in infants' blood significantly increased as Cu ( $p < 0.001$ ,  $\rho = 0.34$ )  
343 and Ag ( $p < 0.001$ ,  $\rho = 0.28$ ) concentrations in infants' blood. There were significant  
344 negative correlations between Pb and Fe concentrations in infants' blood as well as in  
345 mothers' blood.

346 Table 2. Correlation coefficients ( $R^2$ ) among factors and Pb concentrations in samples in all infants and mothers in the present study.

	Age of infants	Height	Weight	Age of mothers	of BLLs in infants	BLLs in mothers	Pb in breastmilk	Pb in infants' feces
Age of infants		0.64***	0.59***	NS	0.46***	NS	NS	0.44***
Height			0.76***	NS	0.36***	NS	NS	0.27***
Weight				NS	0.32***	NS	NS	0.28***
Age of mothers					NS	NS	NS	NS
BLLs in infants						0.68***	0.52***	0.83***
BLLs in mothers							0.58***	0.57***
Pb in breastmilk								0.46***
Pb in infants' feces								

347 Note: \*\*\* indicates  $p < 0.001$ . BLL, blood lead level; NS, not significant.

### 348 **3.4 Daily Intake of Pb in Infants through Breastfeeding and Soil Ingestion**

349 Table 3 shows the results of daily intake of Pb in infants through breastfeeding  
350 and soil ingestion. The results ranged from 0.3 to 40.5  $\mu\text{g}/\text{day}$  through breastfeeding and  
351 from 10.4 to 189.8  $\mu\text{g}/\text{day}$  through soil ingestion depending on Pb concentrations in the  
352 samples. Daily intake of Pb through combined breastfeeding and soil ingestion ranged  
353 from 10.7 to 230.3  $\mu\text{g}/\text{day}$ . The reference value of daily intake of Pb which would be  
354 associated with developmental neurotoxicity in infants in the present study was 3.7  
355  $\mu\text{g}/\text{day}$  (the mean bw: 7.3 kg) and ranged from 1.0  $\mu\text{g}/\text{day}$  (the minimum bw: 1.9 kg) to  
356 6.1  $\mu\text{g}/\text{day}$  (the maximum bw: 12.1) based on the EFSA's reference value (0.5  $\mu\text{g}/\text{kg}$   
357 bw/day) (EFSA, 2010). The minimum of calculated daily intake of Pb even only  
358 through soil ingestion exceeded the maximum of the limit.  
359

360

361 Table 3. Sum of possible daily intake of Pb ( $\mu\text{g/day}$ ) through breastfeeding and soil ingestion.

		Daily intake of Pb through soil ingestion (30 mg/day)				
		Maximum Pb (6326.7 mg/kg)	Mean Pb (1047.6 mg/kg)	Minimum Pb (345.7 mg/kg)		
		189.8 $\mu\text{g/day}$	31.4 $\mu\text{g/day}$	10.4 $\mu\text{g/day}$	364	
Daily intake of Pb through breastmilk (0.78 L/day)	Maximum Pb (51.9 $\mu\text{g/L}$ )	40.5 $\mu\text{g/day}$	230.3	71.9	50.8	365
	Mean Pb (5.3 $\mu\text{g/L}$ )	4.1 $\mu\text{g/day}$	193.9	35.5	14.5	366
	Minimum Pb (0.4 $\mu\text{g/L}$ )	0.3 $\mu\text{g/day}$	190.1	31.8	10.7	367
						368
						369
						370

371 Note: The reference value of daily intake Pb in the present study was 3.7  $\mu\text{g}$  (the mean bw: 7.3 kg) ranged from 1.0  $\mu\text{g}$  (the minimum  
 372 bw: 1.9 kg) to 6.1  $\mu\text{g}$  (the maximum bw: 12.1) based on the EFSA's reference value (0.5  $\mu\text{g/kg bw/day}$ ) (EFSA, 2010).

373

### 374 3.5 Pb Isotope Ratio Analysis

375 Table 4 shows the mean values of the  $^{208}\text{Pb}/^{206}\text{Pb}$ ,  $^{207}\text{Pb}/^{206}\text{Pb}$ ,  $^{208}\text{Pb}/^{204}\text{Pb}$ , and  
 376  $^{207}\text{Pb}/^{204}\text{Pb}$  ratios in the samples. The  $^{208}\text{Pb}/^{206}\text{Pb}$  Pb isotope ratios in mothers' blood  
 377 ( $2.127 \pm 0.006$ ) were significantly different from those in infants' blood and feces and  
 378 the soil ( $2.129 \pm 0.006$ ,  $2.130 \pm 0.004$ , and  $2.131 \pm 0.002$ , respectively,  $p < 0.05$ ). On  
 379 the other hand, there was no significant difference in  $^{207}\text{Pb}/^{206}\text{Pb}$  ratios between infants'  
 380 and mothers' blood. Both  $^{208}\text{Pb}/^{204}\text{Pb}$ , and  $^{207}\text{Pb}/^{204}\text{Pb}$  ratios in mothers' blood were  
 381 significantly different from those in other samples ( $p < 0.05$ ), except for those in  
 382 breastmilk. Pb isotope ratios in soil samples were similar to those reported for Kabwe  
 383 galena (Kamona et al., 1999).

384 Pb isotope ratios in infants' and mothers' blood, and infants' feces were closer  
 385 to those reported for Kabwe galena (Kamona et al., 1999) as the reciprocal of Pb  
 386 concentrations decreased (Supplementary Fig. S4).

387 The  $^{208}\text{Pb}/^{204}\text{Pb}$  and  $^{207}\text{Pb}/^{204}\text{Pb}$  ratios (Fig. 3b) clearly show trends of samples  
 388 compared to  $^{208}\text{Pb}/^{206}\text{Pb}$  and  $^{207}\text{Pb}/^{206}\text{Pb}$  ratios (Fig. 3a and Supplementary Fig. S5).

389


390

391 Table 4. Mean  $\pm$  SD values of the Pb isotope ratios in different samples from Kasanda  
 392 and Makululu.


Samples (N)	$^{208}\text{Pb}/^{206}\text{Pb}$	$^{207}\text{Pb}/^{206}\text{Pb}$	$^{208}\text{Pb}/^{204}\text{Pb}$	$^{207}\text{Pb}/^{204}\text{Pb}$	$^{206}\text{Pb}/^{204}\text{Pb}$
Infants' blood (26)	$2.129 \pm 0.006$ <sup>ac</sup>	$0.8724 \pm 0.003$ <sup>ac</sup>	$38.271 \pm 0.063$ <sup>acd</sup>	$15.691 \pm 0.012$ <sup>a</sup>	$17.997 \pm 0.062$ <sup>ab</sup>
Mothers' blood (26)	$2.127 \pm 0.006$ <sup>b</sup>	$0.8726 \pm 0.002$ <sup>a</sup>	$38.223 \pm 0.080$ <sup>bc</sup>	$15.680 \pm 0.015$ <sup>b</sup>	$17.970 \pm 0.059$ <sup>a</sup>
Breastmilk (26)	$2.129 \pm 0.001$ <sup>bc</sup>	$0.8729 \pm 0.0005$ <sup>b</sup>	$38.271 \pm 0.046$ <sup>c</sup>	$15.691 \pm 0.010$ <sup>b</sup>	$17.972 \pm 0.018$ <sup>a</sup>
Infants' feces (26)	$2.130 \pm 0.004$ <sup>ac</sup>	$0.8725 \pm 0.002$ <sup>c</sup>	$38.349 \pm 0.067$ <sup>d</sup>	$15.709 \pm 0.012$ <sup>a</sup>	$18.005 \pm 0.048$ <sup>b</sup>
Soil (32)	$2.131 \pm 0.002$ <sup>a</sup>	$0.8724 \pm 0.001$ <sup>d</sup>	$38.407 \pm 0.036$ <sup>e</sup>	$15.723 \pm 0.005$ <sup>c</sup>	$18.022 \pm 0.027$ <sup>c</sup>
Kabwe galena (Kamona et al., 1999)	$2.134 \pm 0.0009$	$0.8731 \pm 0.0003$	$38.410 \pm 0.033$	$15.713 \pm 0.010$	$17.997 \pm 0.007$

393 Note: Various small letters indicate a significant difference among areas ( $p < 0.05$ ).

394 Note: The number of significant digits in each ratio value were adjusted with those of  
395 the reference (Kamona et al., 1999)



396



397

398

399 Fig. 3. Pb isotope ratios of individual sample (3a:  $^{208}\text{Pb}/^{206}\text{Pb}$  and  $^{207}\text{Pb}/^{206}\text{Pb}$  and 3b:  
400  $^{208}\text{Pb}/^{204}\text{Pb}$  and  $^{207}\text{Pb}/^{204}\text{Pb}$ ) in different samples from Kasanda and Makululu. The blue  
401 circle, red triangle, green square, orange rhombus, and black cross markers indicate  
402 infants' blood, mothers' blood, breastmilk, infants' fecal, and soil samples, respectively.  
403 The reference value of Kabwe galena was obtained from a report by Kamona et al.  
404 (1999) and is indicated by a plus sign.

405

### 406 **3.6 Plasma Biochemical Analysis**

407 Supplementary Table S15 shows the mean values of the plasma biochemical  
408 analysis on mothers' plasma. The mean values of all parameters in mothers were within  
409 the reference range or slightly higher, except for ALP. The ALT and ALP  
410 concentrations in the mothers from sites near the mine were significantly higher than  
411 those in the mothers from Bwacha ( $p < 0.05$ ). On the other hand, LDH ( $p < 0.05$ ), Alb  
412 ( $p = 0.049$ ), and metallothionein ( $p < 0.001$ ) concentrations in the mothers from Bwacha  
413 were significantly higher than those in mothers from sites near the mine. T-Bil  
414 concentrations in the mothers from Bwacha were higher than those in the mothers from  
415 sites near the mine, although this difference was not significant.

416 In all mothers, there was a significant positive correlation between the Pb and  
417 AST concentrations (Supplementary Table S16,  $p < 0.05$ ,  $\rho = 0.16$ ), and Fe and  
418 metallothionein concentrations (Supplementary Table S17,  $p < 0.05$ ,  $\rho = 0.17$ ). The  
419 ALT concentrations displayed an almost significant increase as Pb concentrations

420 increased ( $p = 0.08$ ,  $\rho = 0.13$ ). The Alb ( $p < 0.05$ ,  $\rho = -0.14$ ) and metallothionein  
421 concentrations ( $p < 0.001$ ,  $\rho = -0.53$ ) significantly decreased as the Pb concentrations  
422 increased.

423           In the mothers from sites near the mine, the Pb concentrations were significantly  
424 associated with increased AST ( $p < 0.05$ ,  $\rho = 0.17$ ) and LDH ( $p < 0.05$ ,  $\rho = 0.19$ )  
425 concentrations. On the other hand, the BUN ( $p < 0.05$ ,  $\rho = -0.26$ ) and metallothionein  
426 concentrations ( $p < 0.001$ ,  $\rho = -0.43$ ) significantly decreased as Pb concentrations  
427 increased. Moreover, the metallothionein concentrations significantly associated with  
428 decreased Cu concentrations ( $p < 0.05$ ,  $\rho = -0.21$ ). In the mothers from Bwacha, there  
429 were no significant correlations between Pb and plasma biochemical factors.

#### 430 **4. Discussion**

431 To the authors' knowledge, the present study undertook the first ever analysis of  
432 Pb in breastmilk of mothers from the sites around the mine in Kabwe. The overall mean  
433 Pb concentration in breastmilk was 5.3 µg/L, which was marginally above the  
434 acceptable level of 2 to 5 µg/L for breastfeeding (WHO, 1989), and 30.0% of breastmilk  
435 samples contained Pb levels above the acceptable level. Compared with previous  
436 studies that reported elevated Pb concentrations in breastmilk ranged from 8.8 to 35.4  
437 µg/L (Isaac et al., 2012; Turan et al., 2001), Pb concentrations in the breastmilk of  
438 mothers from sites near the mine were comparable or even lower. On the other hand, the  
439 mean values of Pb concentrations in breastmilk of mothers from Katondo and Bwacha  
440 (2.3 µg/L) were within the acceptable level, which agrees with the results obtained in  
441 unpolluted areas in other reports (Ettinger et al., 2014; Klein et al., 2017), although the  
442 highest individual Pb breastmilk concentration in this study was found in Katondo (51.9  
443 µg/L). Pb in breastmilk may be one of the sources of Pb exposure in infants. Pb  
444 concentrations in breastmilk in this study were 5.6% of Pb concentrations in maternal  
445 blood. This result agreed with previous studies reporting breastmilk/mothers' blood  
446 ratios between 1% and 10% (Anastácio et al., 2004; Ettinger et al., 2005; Koyashiki et  
447 al., 2010; Koyashiki, et al., 2010; Gulson et al., 1998). In an evaluation of  
448 breastmilk/mothers' blood relationships, Gulson et al. (1998) suggested that a ratio of  
449 Pb concentration in breast milk to the concentration of Pb in maternal blood greater than  
450 15% should be treated with caution, higher values than this arising probably from  
451 sampling and/or analytical contamination.

452 High overall mean Pb concentration in infants' feces (39.2 mg/kg) was recorded  
453 in the present study. The highest Pb concentration in infants' feces was 3002 mg/kg.

454 These results are in agreement with a previous study conducted in Kabwe (Yabe et al.,  
455 2018). Even in Bwacha, which is far from the mine, high Pb concentrations were found  
456 in infants' feces. This suggests that infants in Kabwe were exposed to Pb via ingestion.

457 Among sample types, Pb concentrations in infants' feces were significantly  
458 higher than in blood and breastmilk. There were significant positive correlations among  
459 samples. BLLs in infants significantly increased with BLLs in mothers ( $p < 0.001$ ,  $\rho =$   
460  $0.68$ ), Pb concentrations in breastmilk ( $p < 0.001$ ,  $\rho = 0.43$ ), and infants' feces ( $p <$   
461  $0.001$ ,  $\rho = 0.82$ ). These results suggest that mothers, as well as infants, are exposed to  
462 Pb from the environment, as they share the same living conditions. Pb concentrations in  
463 infants' feces may be a useful indicator of Pb exposure in infants, in addition to BLLs.

464 The present study found different trends of Pb exposure in infants and mothers  
465 among the studied areas in Kabwe. The mean BLL in infants and mothers from  
466 Makululu, which is further from the mine than Kasanda, were the highest among the  
467 area. Following Makululu, those from Kasanda and Chowa were the second and the  
468 third highest, respectively. Yabe et al. (2015) reported a similar trend for BLLs in  
469 children under seven years old, but the highest BLL mean was found in Kasanda,  
470 among the three sites. Since most areas in Makululu are dusty and unpaved compared to  
471 Kasanda, residents could easily come in to contact with polluted soils or dusts in the  
472 area. Pb exposure in infants and mothers from Kasanda and Makululu, which are  
473 located on the western side of the mine and in the direction of prevailing winds, could  
474 be higher than that in Chowa, which lies in the opposite direction. Pb concentrations in  
475 breastmilk and infants' feces showed similar trends to BLLs in infants and mothers.  
476 These trends agreed with those determined in earlier studies by Tembo et al. (2006) and  
477 Nakayama et al. (2011), who reported similar trends in the soils in Kabwe. Toyomaki et

478 al. (2020) reported that Pb concentrations in dog blood decreased with the distance from  
479 the mine in Kabwe, and that the exposure to Pb in dogs remarkably decreased about 5  
480 km away from the mine. These findings suggested that location of the townships in the  
481 relation to the wind direction and distance from the mine influence the extent of the  
482 exposure to Pb in infants and mothers in Kabwe, as has been shown in other studies  
483 (Soto-Jiménez and Flegal, 2011; Yun et al., 2018).

484 In the present study, Fe, Cu, Zn and Ag were also analyzed. In contrast to Pb,  
485 the concentrations of Fe, Cu, Zn and Ag in the blood of infants and mothers tended to  
486 be lower in areas surrounding the mine area than in Bwacha. Given that Pb poisoning is  
487 known to cause anemia, the negative correlation between Pb and Fe concentrations  
488 suggested that the higher Pb concentrations negatively affected Fe metabolism,  
489 especially absorption (Hegazy et al., 2010; Chen et al., 2019). Positive correlations were  
490 found between Pb and Cu, Zn, as well as Ag. These positive co-exposures could be  
491 attributed to the presence of Pb, Zn and Ag in the galena in Kabwe as confirmed in  
492 previous studies (Nakayama et al., 2011).

493 Blood lead levels in boys were significantly higher than those in girls in  
494 Makululu. The same trend for children under seven years old in the same township was  
495 previously reported (Yabe et al., 2015). Moreover, Pb concentrations in infants' feces of  
496 boys were significantly higher than those of girls in Chowa. This difference could be  
497 attributed to differences in breastmilk consumptions as reported by Costa et al. (2010)  
498 where boys consumed breastmilk 0.05 kg/d more than girls. These results suggest that  
499 boys are more exposed to Pb via ingestion than girls.

500 Significant positive correlations between BLLs in infants and age of the infants  
501 were found. The hand-to-mouth or object-to-mouth (pica) behavior of children, and

502 high absorbance of ingested Pb from the gastrointestinal tract are well known factors  
503 attributed to high Pb exposure in children (Wani et al., 2015). However, younger infants  
504 who are not ambulatory could display less hand-to-mouth behaviors, as they are under  
505 the care of their mothers or guardians although inhalation may be an important pathway  
506 for very young children in windy environments (Gulson et al., 2009). Given that BLLs  
507 in infants increases from birth to around two years of age in Kabwe, it is important to  
508 pay more attention to activities of infants during this period. Moreover, cleaning floors  
509 in the house where infants spend most of the time and the kitchen where food could be  
510 contaminated by house dust would be important to reduce Pb exposure in infants. On  
511 the other hand, only BLLs in mothers from Bwacha significantly decreased as the age of  
512 mothers increased. Adults in Bwacha, which is farthest from the mine, could be less  
513 exposed to Pb, thus, Pb in the blood of adults may mainly occur from redistribution of  
514 endogenous bone-derived Pb (Gulson et al. 1998; Manton et al. 2003). From this point  
515 of view, adults even from sites far from the mine in Kabwe could be chronically  
516 exposed to Pb via endogenous exposure.

517         The Pb isotope ratios in infants' samples, especially feces, were almost identical  
518 to those in the soil samples. The soil samples exhibited Pb isotope ratios similar to those  
519 in Kabwe galena (Kamona et al., 1999). Furthermore, a Pb exposure study on rats  
520 exposed to lead in the Kabwe soil revealed that the Pb isotope ratios in these biological  
521 samples were also similar to those in Kabwe galena (Kamona et al., 1999; Nakayama et  
522 al., 2019). These results suggest that contaminated soil or dust from the mine could be  
523 one of the important sources of Pb exposure in infants of Kabwe as well as breastmilk.  
524 Understanding which infant behaviors and activities are related to their Pb exposure is  
525 required to determine the routes and to minimize the exposure. Pb isotope ratios in

526 infants' blood were similar to those in mothers' blood as infants could be exposed to Pb  
527 through breastfeeding and through the placenta before birth. On the other hand, Pb  
528 isotope ratios in mothers' samples, especially blood were different from those in  
529 infants' feces and the soil samples. Pb isotope ratios in chicken, goats, and vegetables in  
530 Kabwe reported by Nakata et al. (2016) were similar to the results in the present study,  
531 suggesting that consuming contaminated food could be an important route of Pb  
532 exposure in mothers. In the present study, both  $^{206}\text{Pb}$  and  $^{204}\text{Pb}$  ratios were used to  
533 compare the differences of Pb isotope ratios among sample types. Both results were  
534 similar, but  $^{204}\text{Pb}$ -based ratios displayed clear differences among sample types.  
535 Therefore,  $^{204}\text{Pb}$  ratios could be more useful than  $^{206}\text{Pb}$  ratios to elucidate the source of  
536 Pb exposure as recommended by Gulson (2008).

537         Daily intake of Pb in infants through breastfeeding and soil ingestion was  
538 calculated to estimate the burden of the possible routes of Pb exposure. Given that Pb  
539 exposure can cause pediatric neurodevelopmental impairments even at low level  
540 (Canfield et al., 2003), the reference value 0.5  $\mu\text{g}/\text{kg}$  bw/day was used (US EPA, 2011).  
541 From the results obtained in the present study, daily intake of Pb even through soil  
542 ingestion alone highly exceeded the reference value. Although the bioavailability of Pb  
543 in galena is known to be minimal (Rasmussen et al., 2011), Pb exposure through soil  
544 ingestion could be the important source due to the high concentrations. Moreover, the  
545 results of daily intake of Pb through breastfeeding using the maximum Pb  
546 concentrations were larger than the reference value. These results suggest that both Pb  
547 though breastfeeding and soil ingestion are important sources of Pb exposure in infants.  
548 In the present study, the daily intake of Pb from the environment was only calculated  
549 from soil ingestion. However, US EPA (2011) estimated the same amount (30 mg/day)

550 of dust ingestion in parallel with soil ingestion. This implies that the actual daily intake  
551 in the field could be underestimated in the present study. Thus, future studies need to  
552 evaluate the detail of Pb exposure in infants as well as mothers in Kabwe including  
553 other routes, such as dust ingestion.

554 In contrast to the high BLLs in mothers, the plasma biochemical profiles of most  
555 analyzed parameters were interestingly within, or close to, the standard reference values,  
556 except in the case of ALP. These results indicate that Pb exposure in Kabwe mothers  
557 did not significantly impact their health, as was observed during sampling, where all  
558 sampled mothers appeared healthy. More specifically, the ALT and ALP values in the  
559 mothers from sites near the mine were significantly higher than those in the mothers  
560 from Bwacha although these values were not significantly correlated with BLLs in  
561 mothers. Therefore, these results could not only be attributed to Pb exposure, but also  
562 other factors. On the other hand, LDH and AST, which are indicators of liver function,  
563 significantly increased as BLLs increased. These results suggest that Pb exposure in  
564 mothers may have caused some mild liver damage. High Pb exposure is known to cause  
565 kidney damage in conjunction with an increase in BUN and a decrease in Alb. However,  
566 both biomarkers significantly decreased as BLLs increased in mothers. During a  
567 previous study on Pb poisoning in refugee children in the United States, the CDC  
568 reported chronic and acute malnutrition as risk factors for Pb poisoning (CDC, 2005).  
569 Further studies in Kabwe should therefore focus on the relationship between Pb  
570 exposure and nutrition status. In our study, metallothionein concentrations significantly  
571 decreased as BLLs increased. A previous study by Mustonen et al. (2014) found  
572 constant metallothionein expression in earthworms from a contaminated site, and  
573 therefore suggested that the inducibility of the metallothionein response could be lost in

574 earthworms with a history of metal exposure. It is probable that local people in the sites  
575 near the mine may be chronically exposed to metals, including Pb, over a long period of  
576 time compared to people in sites far from the mine, such as Bwacha. Therefore,  
577 metallothionein expression in people residing near the mine was lower than that in  
578 people residing far from the mine. Further studies should focus on both metallothionein  
579 concentrations and gene expression. In the present study, it is difficult to exclude the  
580 possibility of other factors or diseases since a detailed questionnaire survey or medical  
581 check-ups were not performed.

582         The findings in the present study suggest that one of the important sources of Pb  
583 exposure in infants could be Pb from the environment, especially from soils. Daily  
584 intake of Pb in infants through soil ingestion could be enough to exceed the EFSA's  
585 reference value of daily intake of Pb. It is important to minimize Pb exposure in  
586 infants from soil and dust.. Thus, remediation of the environment in Kabwe is  
587 urgently needed to reduce Pb exposure. High BLLs in mothers could also be one of the  
588 important sources of Pb exposure in infants via breastfeeding. Also, high BLLs in  
589 mothers may cause their fetus to be exposed to Pb during pregnancy. In the current  
590 situation, chelation therapy for Pb poisoning is prioritized more in children than in  
591 adults, as children are more vulnerable to Pb. However, in utero exposure to  
592 environmental lead may be adversely associated with neurodevelopment at two years of  
593 age (Lin et al., 2013). Pilsner et al. (2009) reported that the epigenome of the  
594 developing fetus can be influenced by the maternal cumulative lead burden, which may  
595 influence long-term epigenetic programming and disease susceptibility throughout a  
596 child's life. Reducing the Pb exposure in mothers is important to reduce Pb exposure in  
597 fetuses via the placenta, as well as Pb exposure in infants, via breastfeeding. Thus, it is

598 necessary that mothers with high BLLs are treated with chelation therapy, as well as

599 their children.

600

601 **5. Conclusions**

602 High Pb concentrations in breastmilk, which were above the WHO acceptable  
603 level for breastfeeding, could be one of the important sources of Pb exposure in infants.  
604 The results of the isotope ratio analysis suggest that Pb from the environment, such as  
605 contaminated soil is one of the important sources of Pb exposure in infants. Moreover,  
606 Pb exposure in infants through breastfeeding and soil ingestion potentially exceeded the  
607 EFSA's reference value of daily intake of Pb. Therefore, environmental remediation, in  
608 parallel with chelation therapy, is urgently needed to reduce the Pb exposure in infants  
609 and mothers in Kabwe. Moreover, mothers with high BLLs in Kabwe should be treated  
610 with chelation therapy to reduce the Pb exposure of their infants via breastfeeding.

611

612 **Acknowledgements**

613 We are grateful to the infants and mothers who participated in this study. We are  
614 also grateful to Takahiro Ichise and Nagisa Hirano (Laboratory of Toxicology, Faculty  
615 of Veterinary Medicine, Hokkaido University) for technical support with metal  
616 concentrations and isotope analyses. This work was supported by the Grants-in-Aid for  
617 Scientific Research from the Ministry of Education, Culture, Sports, Science and  
618 Technology of Japan awarded to M. Ishizuka (No. 16H0177906, 18K1984708,  
619 18KK028708, 21H02351), Ikenaka (18H0413208), and S.M.M. Nakayama (No.  
620 17KK0009 and 20K20633). This work was also supported by the foundation of JSPS  
621 Bilateral Open Partnership Joint Research Projects (JPJSBP120209902). This work was  
622 also supported by JSPS CORE to CORE program (MI), The Japan Prize Foundation;  
623 and Hokkaido University's SOUSEI Support Program for Young Researchers in  
624 FY2020 (SMMN); Hokkaido University Specific Research Projects (MI). We also  
625 acknowledge financial support from the Program for Leading Graduate Schools  
626 “Fostering Global Leaders in Veterinary Science for Contributing to One Health” (F01),  
627 Grant-in-Aid for JSPS Research Fellow (17J05756), “Global Leadership Training  
628 Programme in Africa (GLTP)” of United Nations University, as well as the Japan Prize  
629 Foundation. This research was also supported by JST/JICA, SATREPS (Science and  
630 Technology Research Partnership for Sustainable Development; No. JPMJSA1501) and  
631 aXis (Accelerating Social Implementation for SDGs Achievement; No. JPMJAS2001)  
632 funded by JST as well as Program for supporting introduction of the new sharing  
633 system (JPMXS0420100619). The English was proofread by Uni-edit ([https://uni-  
634 edit.net/usa](https://uni-edit.net/usa)).

635

636 **Conflict of interest**

637 The authors declare no conflict of interest.

638

639 **References**

- 640 ➤ Ajumobi, O.O., Tsofo, A., Yango, M., Aworh, M.K., Anagbogu, I.N., Mohammed,  
641 A., Umar-Tsafe, N., Mohammed, S., Abdullahi, M., Davis, L., Idris, S., Poggensee,  
642 G., Nguku, P., Gitta, S., Nsubuga, P., 2014. High concentration of blood lead levels  
643 among young children in Bagega community, Zamfara – Nigeria and the potential  
644 risk factor. *Pan Afr Medical J* 18, 14.  
645 <https://doi.org/10.11694/pamj.suppl.2014.18.1.4264>
- 646 ➤ Anastácio, A. da S., Silveira, C.L.P. da, Miekeley, N., Donangelo, C.M., 2004.  
647 Distribution of Lead in Human Milk Fractions: Relationship with Essential  
648 Minerals and Maternal Blood Lead. *Biol Trace Elem Res* 102, 027–038.  
649 <https://doi.org/10.1385/bter:102:1-3:027>
- 650 ➤ Barbosa, F., Tanus-Santos, J.E., Gerlach, R.F., Parsons, P.J., 2005. A Critical  
651 Review of Biomarkers Used for Monitoring Human Exposure to Lead: Advantages,  
652 Limitations, and Future Needs. *Environ Health Persp* 113, 1669–1674.  
653 <https://doi.org/10.1289/ehp.7917>
- 654 ➤ Bizoń, A., Milnerowicz, H., 2014. Participation of metallothionein and superoxide  
655 dismutase in the blood of smoking smelters. *Int J Occup Med Env* 27, 326–334.  
656 <https://doi.org/10.2478/s13382-014-0258-8>
- 657 ➤ Bollhöfer, A., Rosman, K.J.R., 2001. Isotopic source signatures for atmospheric  
658 lead: the Northern Hemisphere. *Geochim Cosmochim Acta* 65, 1727–1740.  
659 [https://doi.org/10.1016/S0016-7037\(00\)00630-X](https://doi.org/10.1016/S0016-7037(00)00630-X)
- 660 ➤ Calabrese, E.J., Stanek, E.J., 1995. A Dog's Tale: Soil Ingestion by a Canine.  
661 *Ecotox Environ Safe* 32, 93–95. <https://doi.org/10.1006/eesa.1995.1090>
- 662 ➤ Canfield, R.L., Henderson, C.R., Cory-Slechta, D.A., Cox, C., Jusko, T.A.,  
663 Lanphear, B.P., 2003. Intellectual Impairment in Children with Blood Lead  
664 Concentrations below 10 µg per Deciliter. *New Engl J Medicine* 348, 1517–1526.  
665 <https://doi.org/10.1056/nejmoa022848>
- 666 ➤ Centers for Disease Control and Prevention, 2019. CDC - Lead - Blood Lead  
667 Levels in Children. Available: [https://www.cdc.gov/nceh/lead/prevention/blood-](https://www.cdc.gov/nceh/lead/prevention/blood-lead-levels.htm)  
668 [lead-levels.htm](https://www.cdc.gov/nceh/lead/prevention/blood-lead-levels.htm) [accessed 20 February 2020]

- 669 ➤ Centers for Disease Control and Prevention ,2012. CDC Response to Advisory  
670 Committee on Childhood Lead Poisoning Prevention Recommendations in “Low  
671 Level Lead Exposure Harms Children: A Renewed Call of Primary Prevention.”.  
672 Available:  
673 [https://www.cdc.gov/nceh/lead/ACCLPP/Final\\_Document\\_030712.pdf](https://www.cdc.gov/nceh/lead/ACCLPP/Final_Document_030712.pdf) [accessed  
674 20 February 2020]
- 675 ➤ Centers for Disease Control and Prevention, 2010. Guidelines for the identification  
676 and management of lead exposure in pregnant and lactating women. Available:  
677 <https://www.cdc.gov/nceh/lead/publications/leadandpregnancy2010.pdf> [accessed  
678 19 February 2020]
- 679 ➤ Centers for Disease Control and Prevention, 2005. Elevated blood lead levels in  
680 refugee children--New Hampshire, 2003-2004. *Mmwr Morbidity Mortal Wkly Rep*  
681 54, 42–6.
- 682 ➤ Centers for Disease Control and Prevention, 2002. Managing elevated blood lead  
683 levels among young children: recommendations from the Advisory Committee on  
684 Childhood Lead Poisoning Prevention. Available:  
685 <https://www.cdc.gov/nceh/lead/casemanagement/managingEBLLs.pdf> [accessed 19  
686 February 2020]
- 687 ➤ Chen C-J, Lin T-Y, Wang C-L, Ho C-K, Chuang H-Y, Yu H-S. Interactive Effects  
688 between Chronic Lead Exposure and the Homeostatic Iron Regulator Transport  
689 HFE Polymorphism on the Human Red Blood Cell Mean Corpuscular Volume  
690 (MCV). *International Journal of Environmental Research and Public Health*. 2019;  
691 16(3):354. <https://doi.org/10.3390/ijerph16030354>
- 692 ➤ Costa, T.H.M. da, Haisma, H., Wells, J.C.K., Mander, A.P., Whitehead, R.G.,  
693 Bluck, L.J.C., 2010. How much human milk do infants consume? Data from 12  
694 countries using a standardized stable isotope methodology. *J Nutrition* 140, 2227–  
695 32. <https://doi.org/10.3945/jn.110.123489>
- 696 ➤ Ettinger, A.S., Roy, A., Amarasiriwardena, C.J., Smith, D., Lupoli, N., Mercado-  
697 García, A., Lamadrid-Figueroa, H., Tellez-Rojo, M.M., Hu, H., Hernández-Avila,  
698 M., 2014. Maternal Blood, Plasma, and Breast Milk Lead: Lactational Transfer and

- 699 Contribution to Infant Exposure. *Environ Health Persp* 122, 87–92.  
700 <https://doi.org/10.1289/ehp.1307187>
- 701 ➤ Ettinger, A.S., Téllez-Rojo, M.M., Amarasinghwardena, C., Peterson, K.E., Schwartz,  
702 J., Aro, A., Hu, H., Hernández-Avila, M., 2005. Influence of Maternal Bone Lead  
703 Burden and Calcium Intake on Levels of Lead in Breast Milk over the Course of  
704 Lactation. *Am J Epidemiol* 163, 48–56. <https://doi.org/10.1093/aje/kwj010>
- 705 ➤ European Food Safety Authority, 2010. Scientific Opinion on Lead in Food. *Efsa J*  
706 8, 1570. <https://doi.org/10.2903/j.efsa.2010.1570>
- 707 ➤ Gulson, B., 2008. Stable lead isotopes in environmental health with emphasis on  
708 human investigations. *Sci Total Environ* 400, 75–92.  
709 <https://doi.org/10.1016/j.scitotenv.2008.06.059>
- 710 ➤ Gulson, B., Korsch, M., Matison, M., Douglas, C., Gillam, L., McLaughlin, V.,  
711 2009. Windblown Lead Carbonate as the Main Source of Lead in Blood of  
712 Children from a Seaside Community: An Example of Local Birds as “Canaries in  
713 the Mine.” *Environ Health Persp* 117, 148–154. <https://doi.org/10.1289/ehp.11577>
- 714 ➤ Gulson, B.L., Jameson, C.W., Mahaffey, K.R., Mizon, K.J., Patison, N., Law, A.J.,  
715 Korsch, M.J., Salter, M.A., 1998. Relationships of lead in breast milk to lead in  
716 blood, urine, and diet of the infant and mother. *Environ Health Persp* 106, 667–674.  
717 <https://doi.org/10.1289/ehp.98106667>
- 718 ➤ Gulson, B.L., Mahaffey, K.R., Jameson, C.W., Mizon, K.J., Korsch, M.J., Cameron,  
719 M.A., Eisman, J.A., 1998. Mobilization of lead from the skeleton during the  
720 postnatal period is larger than during pregnancy. *J Lab Clin Med* 131, 324–329.  
721 [https://doi.org/10.1016/s0022-2143\(98\)90182-2](https://doi.org/10.1016/s0022-2143(98)90182-2)
- 722 ➤ Haefliger, P., Mathieu-Nolf, M., Locicero, S., Ndiaye, C., Coly, M., Diouf, A., Faye,  
723 A.L., Sow, A., Tempowski, J., Pronczuk, J., Junior, A.P.F., Bertollini, R., Neira, M.,  
724 2009. Mass lead intoxication from informal used lead-acid battery recycling in  
725 dakar, senegal. *Environ Health Persp* 117, 1535–40.  
726 <https://doi.org/10.1289/ehp.0900696>
- 727 ➤ Hamelin, B., Manhès, G., Albarede, F., Allègre, C.J., 1985. Precise lead isotope  
728 measurements by the double spike technique: a reconsideration. *Geochimica et*  
729 *Cosmochimica Acta* 49, 173–182.

- 730 ➤ Hegazy AA, Zaher MM, Abd El-Hafez MA, Morsy AA, Saleh RA. Relation  
731 between anemia and blood levels of lead, copper, zinc and iron among children.  
732 BMC Res Notes. 2010 May 12;3:133. doi: 10.1186/1756-0500-3-133. PMID:  
733 20459857; PMCID: PMC2887903.
- 734 ➤ Isaac, C.P.J., Sivakumar, A., Kumar, C.R.P., 2012. Lead Levels in Breast Milk,  
735 Blood Plasma and Intelligence Quotient: A Health Hazard for Women and Infants.  
736 B Environ Contam Tox 88, 145–149. <https://doi.org/10.1007/s00128-011-0475-9>
- 737 ➤ Kamona, A.F., Lévêque, J., Friedrich, G., Haack, U., 1999. Lead isotopes of the  
738 carbonate-hosted Kabwe, Tsumeb, and Kipushi Pb-Zn-Cu sulphide deposits in  
739 relation to Pan African orogenesis in the Damaran-Lufilian Fold Belt of Central  
740 Africa. Miner Deposita 34, 273–283. <https://doi.org/10.1007/s001260050203>
- 741 ➤ Klein, L.D., Breakey, A.A., Scelza, B., Valeggia, C., Jasienska, G., Hinde, K., 2017.  
742 Concentrations of trace elements in human milk: Comparisons among women in  
743 Argentina, Namibia, Poland, and the United States. Plos One 12, e0183367.  
744 <https://doi.org/10.1371/journal.pone.0183367>
- 745 ➤ Komárek, M., Ettler, V., Chrastný, V., Mihaljevič, M., 2008. Lead isotopes in  
746 environmental sciences: A review. Environ Int 34, 562–577.  
747 <https://doi.org/10.1016/j.envint.2007.10.005>
- 748 ➤ Kowalska, K., Bizoń, A., Zalewska, M.,  
749 Milnerowicz, H., 2015. The influence of biological and environmental factors on  
750 metallothionein concentration in the blood. J Trace Elem Med Bio 29, 99–103.  
751 <https://doi.org/10.1016/j.jtemb.2014.05.001>
- 752 ➤ Koyashiki, G.A.K., Paoliello, M.M.B., Matsuo, T., Oliveira, M.M.B. de,  
753 Mezzaroba, L., Carvalho, M. de F., Sakuma, A.M., Turini, C., Vannuchi, M.T.O.,  
754 Barbosa, C.S.D., 2010. Lead levels in milk and blood from donors to the Breast  
755 Milk Bank in Southern Brazil. Environ Res 110, 265–71.  
756 <https://doi.org/10.1016/j.envres.2009.12.001>
- 757 ➤ Koyashiki, G.A.K., Paoliello, M.M.B., Tchounwou, P.B., 2010. Lead Levels in  
758 Human Milk and Children's Health Risk: A Systematic Review. Rev Environ  
Health 25, 243–254. <https://doi.org/10.1515/reveh.2010.25.3.243>

- 759 ➤ Kuritani, T., Nakamura, E., 2003. Highly precise and accurate isotopic analysis of  
760 small amounts of Pb using  $^{205}\text{Pb}$ – $^{204}\text{Pb}$  and  $^{207}\text{Pb}$ – $^{204}\text{Pb}$ , two double spikes.  
761 *J Anal Spectrom* 18, 1464–1470. <https://doi.org/10.1039/b310294g>
- 762 ➤ Kuritani, T., Nakamura, E., 2002. Precise isotope analysis of nanogram-level Pb for  
763 natural rock samples without use of double spikes. *Chem Geol* 186, 31–43.  
764 [https://doi.org/10.1016/s0009-2541\(02\)00004-9](https://doi.org/10.1016/s0009-2541(02)00004-9)
- 765 ➤ Lin, C.-C., Chen, Y.-C., Su, F.-C., Lin, C.-M., Liao, H.-F., Hwang, Y.-H., Hsieh,  
766 W.-S., Jeng, S.-F., Su, Y.-N., Chen, P.-C., 2013. In utero exposure to  
767 environmental lead and manganese and neurodevelopment at 2 years of age.  
768 *Environ Res* 123, 52–57. <https://doi.org/10.1016/j.envres.2013.03.003>
- 769 ➤ Manton, W.I., Angle, C.R., Stanek, K.L., Kuntzelman, D., Reese, Y.R.,  
770 Kuehnemann, T.J., 2003. Release of lead from bone in pregnancy and lactation.  
771 *Environ Res* 92, 139–151. [https://doi.org/10.1016/s0013-9351\(03\)00020-3](https://doi.org/10.1016/s0013-9351(03)00020-3)
- 772 ➤ Meyer, P.A., Brown, M.J., Falk, H., 2008. Global approach to reducing lead  
773 exposure and poisoning. *Mutat Res Rev Mutat Res* 659, 166–175.  
774 <https://doi.org/10.1016/j.mrrev.2008.03.003>
- 775 ➤ Mustonen, M., Haimi, J., Väisänen, A., Knott, K.E., 2014. Metallothionein gene  
776 expression differs in earthworm populations with different exposure history.  
777 *Ecotoxicology* 23, 1732–1743. <https://doi.org/10.1007/s10646-014-1338-z>
- 778 ➤ Nakata, H., Nakayama, S.M.M., Yabe, J., Liyambi, A., Mizukawa, H., Darwish,  
779 W.S., Ikenaka, Y., Ishizuka, M., 2016. Reliability of stable Pb isotopes to identify  
780 Pb sources and verifying biological fractionation of Pb isotopes in goats and  
781 chickens. *Environ. Pollut.* 208, 395e403. PMID: 26549754. <https://doi.org/10.1016/j.envpol.2015.10.006>.
- 782 [1016/j.envpol.2015.10.006](https://doi.org/10.1016/j.envpol.2015.10.006).
- 783 ➤ Nakata H, Nakayama SMM, Yabe J, Muzandu K, Toyomaki H, Yohannes YB,  
784 Kataba A, Zyambo G, Ikenaka Y, Choongo K, Ishizuka M. Assessment of  
785 LeadCare® II analysis for testing of a wide range of blood lead levels in  
786 comparison with ICP–MS analysis. *Chemosphere*. 129832 (2021) DOI:  
787 10.1016/j.chemosphere.2021.129832
- 788 ➤ Nakayama, S.M.M., Ikenaka, Y., Hamada, K., Muzandu, K., Choongo, K., Teraoka,  
789 H., Mizuno, N., Ishizuka, M., 2011. Metal and metalloid contamination in roadside

- 790 soil and wild rats around a Pb–Zn mine in Kabwe, Zambia. *Environ Pollut* 159,  
791 175–181. <https://doi.org/10.1016/j.envpol.2010.09.007>
- 792 ➤ Nakayama, S.M.M., Nakata, H., Ikenaka, Y., Yabe, J., Oroszlany, B., Yohannes,  
793 Y.B., Bortey-Sam, N., Muzandu, K., Choongo, K., Kuritani, T., Nakagawa, M.,  
794 Ishizuka, M., 2019. One year exposure to Cd- and Pb-contaminated soil causes  
795 metal accumulation and alteration of global DNA methylation in rats. *Environ*  
796 *Pollut* 252, 1267–1276. <https://doi.org/10.1016/j.envpol.2019.05.038>
- 797 ➤ National Academy of Science, 1972. Lead: Airborne lead in perspective: Biological  
798 effects of atmospheric pollutants. Washington, DC: National Academy of Sciences,  
799 71-177, 281-313.
- 800 ➤ Nassir, I., Al-Sharify, A., Baiee, H., 2013. Lead and cadmium in the breast milk of  
801 lactating mothers living in Hilla City, Babylon, Iraq, during the year 2012. *Journal*  
802 *of Babylon University*. 8(21).
- 803 ➤ Needleman, H., 2004. Lead Poisoning. *Annu Rev Med* 55, 209–222.  
804 <https://doi.org/10.1146/annurev.med.55.091902.103653>
- 805 ➤ Pilsner, J.R., Hu, H., Ettinger, A., Sánchez, B.N., Wright, R.O., Cantonwine, D.,  
806 Lazarus, A., Lamadrid-Figueroa, H., Mercado-García, A., Téllez-Rojo, M.M.,  
807 Hernández-Avila, M., 2009. Influence of Prenatal Lead Exposure on Genomic  
808 Methylation of Cord Blood DNA. *Environ Health Persp* 117, 1466–1471.  
809 <https://doi.org/10.1289/ehp.0800497>
- 810 ➤ Rasmussen, P.E., Beauchemin, S., Chénier, M., Levesque, C., MacLean, L.C.W.,  
811 Marro, L., Jones-Otazo, H., Petrovic, S., McDonald, L.T., Gardner, H.D., 2011.  
812 Canadian house dust study: lead bioaccessibility and speciation. *Environ Sci*  
813 *Technol* 45, 4959–65. <https://doi.org/10.1021/es104056m>
- 814 ➤ Ruckart, P.Z., Ettinger, A.S., Hanna-Attisha, M., Jones, N., Davis, S.I., Breyse,  
815 P.N., 2019. The Flint Water Crisis. *J Public Health Man* 25, S84–S90.  
816 <https://doi.org/10.1097/phh.0000000000000871>
- 817 ➤ Schoning, P., Abraham, J., Burnett, B., 1996. Silicate and metal dust in lungs of  
818 Greyhounds. *Am J Vet Res* 57, 1006–9.

- 819 ➤ Soto-Jiménez, M.F., Flegal, A.R., 2011. Childhood lead poisoning from the smelter  
820 in Torreón, México. *Environ Res* 111, 590–596.  
821 <https://doi.org/10.1016/j.envres.2011.01.020>
- 822 ➤ Tembo, B.D., Sichilongo, K., Cernak, J., 2006. Distribution of copper, lead,  
823 cadmium and zinc concentrations in soils around Kabwe town in Zambia.  
824 *Chemosphere* 63, 497–501. <https://doi.org/10.1016/j.chemosphere.2005.08.002>
- 825 ➤ Toyomaki, H., Yabe, J., Nakayama, S.M.M., Yohannes, Y.B., Muzandu, K.,  
826 Liazambi, A., Ikenaka, Y., Kuritani, T., Nakagawa, M., Ishizuka, M., 2020. Factors  
827 associated with lead (Pb) exposure on dogs around a Pb mining area, Kabwe,  
828 Zambia. *Chemosphere* 125884. <https://doi.org/10.1016/j.chemosphere.2020.125884>
- 829 ➤ Turan, S., Saygi, Ş., Kiliç, Z., Acar, O., 2001. Determination of Heavy Metal  
830 Contents in Human Colostrum Samples by Electrothermal Atomic Absorption  
831 Spectrophotometry. *J Trop Pediatrics* 47, 81–85.  
832 <https://doi.org/10.1093/tropej/47.2.81>
- 833 ➤ United States Environmental Protection Agency, 2011. Exposure Factors  
834 Handbook 2011 Edition (Final Report). U.S. Environmental Protection Agency,  
835 Washington, DC, EPA/600/R-09/052F.
- 836 ➤ Veysseyre, A.M., Bollhöfer, A.F., Rosman, K.J.R., Ferrari, C.P., Boutron, C.F.,  
837 2001. Tracing the Origin of Pollution in French Alpine Snow and Aerosols Using  
838 Lead Isotopic Ratios. *Environ Sci Technol* 35, 4463–4469.  
839 <https://doi.org/10.1021/es0105717>
- 840 ➤ Wani, A., Ara, A., Usmani, J., 2015. Lead toxicity: a review. *Interdiscip Toxicol* 8,  
841 55–64. <https://doi.org/10.1515/intox-2015-0009>
- 842 ➤ World Health Organization, 2010. Childhood lead poisoning. Available:  
843 <https://www.who.int/ceh/publications/leadguidance.pdf> [accessed 19 February  
844 2020]
- 845 ➤ World Health Organization, 1989. Minor and trace elements in human milk.  
846 Geneva, Switz: World Health Organization. Available:  
847 [https://apps.who.int/iris/bitstream/handle/10665/39678/9241561211.pdf?sequence=](https://apps.who.int/iris/bitstream/handle/10665/39678/9241561211.pdf?sequence=1)  
848 [1](https://apps.who.int/iris/bitstream/handle/10665/39678/9241561211.pdf?sequence=1) [accessed 19 February 2020]

- 849 ➤ Yabe, J., Ishizuka, M., Umemura, T., 2010. Current Levels of Heavy Metal  
850 Pollution in Africa. *J Vet Med Sci.* 72, 1257–1263.  
851 <https://doi.org/10.1292/jvms.10-0058>
- 852 ➤ Yabe, J., Nakayama, S., Ikenaka, Y., Yohannes, Y.B., Bortey-Sam, N., Oroszlany,  
853 B., Muzandu, K., Choongo, K., Kabalo, A., Ntapisha, J., Mweene, A., Umemura, T.,  
854 Ishizuka, M., 2015. Lead poisoning in children from townships in the vicinity of a  
855 lead–zinc mine in Kabwe, Zambia. *Chemosphere* 119, 941–947.  
856 <https://doi.org/10.1016/j.chemosphere.2014.09.028>
- 857 ➤ Yabe, J., Nakayama, S.M., Nakata, H., Toyomaki, H., Yohannes, Y.B., Muzandu,  
858 K., Kataba, A., Zyambo, G., Hiwatari, M., Narita, D., Yamada, D., Hangoma, P.,  
859 Munyinda, N.S., Mufune, T., Ikenaka, Y., Choongo, K., Ishizuka, M., 2020.  
860 Current trends of blood lead levels, distribution patterns and exposure variations  
861 among household members in Kabwe, Zambia. *Chemosphere* 243, 125412.  
862 <https://doi.org/10.1016/j.chemosphere.2019.125412>
- 863 ➤ Yabe, J., Nakayama, S.MM., Ikenaka, Y., Yohannes, Y.B., Bortey-Sam, N., Kabalo,  
864 A.N., Ntapisha, J., Mizukawa, H., Umemura, T., Ishizuka, M., 2018. Lead and  
865 cadmium excretion in feces and urine of children from polluted townships near a  
866 lead-zinc mine in Kabwe, Zambia. *Chemosphere* 202, 48–55.  
867 <https://doi.org/10.1016/j.chemosphere.2018.03.079>
- 868 ➤ Yun, S.-W., Baveye, P.C., Kim, D.-H., Kang, D.-H., Lee, S.-Y., Kong, M.-J., Park,  
869 C.-G., Kim, H.-D., Son, J., Yu, C., 2018. Analysis of metal(loid)s contamination  
870 and their continuous input in soils around a zinc smelter: Development of  
871 methodology and a case study in South Korea. *Environ Pollut* 238, 140–149.  
872 <https://doi.org/10.1016/j.envpol.2018.03.020>