HOKKAIDO UNIVERSITY

Title	Assessing long-term coral reef degradation in Indonesia's Tiworo strait marine conservation area using remote sensing and rapid appraisal for fisheries approaches
Author(s)	Yasir Haya, La Ode Muhammad; Sadarun, Baru; Tadjudah, Muslim et al.
Citation	Modeling Earth Systems and Environment, 8(4), 4731-4748 https://doi.org/10.1007/s40808-022-01421-7
Issue Date	2022-11
Doc URL	https://hdl.handle.net/2115/90624
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: http://dx.doi.org/10.1007/s40808-022-01421-7
Type	journal article
File Information	the latest paper_Yasir Haya et al. 2022.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Assessing long-term coral reef degradation in Indonesia's Tiworo Strait Marine Conservation Area using remote sensing and rapid appraisal for fisheries approaches

La Ode Muhammad Yasir Haya[1*], Baru Sadarun[1], Muslim Tadjudah[2], Analuddin Kangkuso[3], La Ode Alirman Afu[1], Asrin Ginong Pratikino[4], Amadhan Takwir[4], Masahiko Fujii[5]

[1]*Marine Science Department, Fisheries and marine Science Faculty, Halu Oleo University, Indonesia.*
[2]*Capture Fisheries Study Program, Fisheries and Marine Science Faculty, Halu Oleo University, Indonesia*.
[3] *Biotechnology Department, Mathematic and Natural Sciences Faculty, Halu Oleo University, Indonesia*.
[4]*Oceanography Study Program, Fisheries and Marine Science Faculty, Halu Oleo University, Indonesia*.
[5]*Faculty of Environmental Earth Science, Hokkaido University, Japan.*

✉ La Ode Muhammad Yasir Haya

laode.haya@uho.ac.id

Abstract

In Indonesia, the coral reef ecosystem in the Tiworo Strait Conservation Area (TSCA) faces various threats of natural and anthropogenic stressors that can damage the coral reef ecosystem's role and services. We analyzed changes in coral reef habitat at TSCA over the 25 years from 1994 to 2019 using multi-temporal and multi-sensor satellite imagery data combined with in-situ measurement data and social surveys. Our results show a decrease in live coral cover from 78.30 ha in 1994 to 8.01 ha in 2019, with a 2.81 ha/year degradation rate. Our analysis of 37 threat attributes shows that the TSCA coral reef ecosystem faces a "high threat" to very high threat levels. Threat scores for coral reefs assessed as facing severe conditions according to threat indices included contributions from the ecological dimension (16.87 = very high threat), economic dimension (31.00 = high threat), social dimension (34.83 = high threat), technological dimension (41.10 = high threat), and law and institutional dimension (26.83 = high threat). Coral reefs will undoubtedly go extinct if local threats continue without preventative measures. Therefore, the sustainability of coral reefs in the TSCA-one of the most important marine conservation sites in the

Coral Triangle Marine Eco-region should be the primary priority for all stakeholders. Appropriate policies and supervision in the field must be carried out rigorously and measurably, implementing the analyzed set of strategies.

Keywords: *coral cover change; rapid appraisal for fisheries (RAPFISH); remote sensing; threats; Tiworo Strait Conservation Area (TSCA), Coral Triangle Eco-region of Indonesia*

1. Introduction

Over the past four decades, numerous studies have identified the trend of continuous coral reef degradation due to natural and anthropogenic pressures. These pressures include enhanced local pollution of coastal areas (Burke *et al.*, 2011; Häder *et al.*, 2020; Randazzo-Eisemann *et al.*, 2021), overfishing (Moussa et al., 2019; Nichols et al., 2019), destructive fishing practices (Yasir Haya and Fujii 2020; Hampton-Smith et al. 2021), sea surface temperature (Hughes et al. 2018; Ngoc 2019), climate change and subsequent ocean warming (Pendleton et al. 2016; Hughes et al. 2017; Zhang et al. 2021), ocean acidification (Allemand and Osborn 2019; Zunino et al. 2021), coral diseases (Lapointe et al. 2019; Hazraty-Kari et al. 2021), nutrient enrichment (Lapointe et al. 2019; Adam et al. 2021), and other anthropogenic threats (Burke et al. 2011; Mellin et al. 2016; Hoegh-Guldberg et al. 2019; Hein et al. 2021). Degradation of coral reefs leads to a loss of their ecological and economic value, particularly the goods and services to coastal and other communities (Cesar and Chong 2004; Mehvar et al. 2018; Yasir Haya and Fujii 2019; Santavy et al. 2021). Given these conditions, coral reef ecosystems require urgent management, such as a marine protected area (MPA) approach (Mellin et al. 2016; Vaughan and Agardy 2020; Hampton-Smith et al. 2021).

MPAs have become one of the most frequently employed marine management tools worldwide for conserving species and habitats, maintaining ecosystem functionality, and ensuring the sustainable use of marine resources (Pagiola et al. 2004; Vaughan and Agardy 2020). In Indonesia, the types and sizes of MPAs differ according to their conservation goals and targets. The Tiworo Strait is a medium-sized marine conservation area that located at the Southeast of the Sulawesi mainland in the center of the Coral Triangle Area, Indonesia (**Fig. 1**). The Tiworo Strait exhibits tremendous marine biodiversity potential with its combination of coral reefs,

54 mangroves, seagrass, and fish, with a total area of around 27,936 ha (KKJI-KP3K, 2019). Because of this potential,
55 the Tiworo Strait was designated as a conservation area by the local government of the Muna Regency and has
56 become one of the most critical marine conservation sites in the Coral Triangle of Southeast Sulawesi, Indonesia.
57 Nevertheless, the coral reefs in the Tiworo Strait Conservation Area (TSCA) are thought to have experienced
58 massive degradation due to pressure from various environmentally unfriendly activities.

59 The primary sources of coral reef degradation are local anthropogenic factors i.e., blast fishing, trap
60 fishing (locally known as “Bubu”), and bottom-trawling net practices (locally known as “Dogol”). These three
61 fishing practices are considered environmentally unfriendly because they overly exploit fishery resources and
62 violate conservation rules (Gorris 2016; Chan and Hodgson 2017; Kularatne 2020) thus allegedly reducing coral
63 cover in this area. Information on coral reef changes in the TSCA, which is desperately needed for habitat
64 management, is not yet available. However, the combination of remote-sensing technology, in-situ data, and social
65 surveys can provide alternative approaches for monitoring changes in reef composition at high spatial and
66 temporal resolution (Andréfouët et al. 2003; Foo and Asner 2019; Chen et al. 2022). These approaches can
67 facilitate monitoring in remote areas that are not directly accessible including coral reef ecosystems (Purkis 2018;
68 De et al. 2021; Molder et al. 2022).

69 Remote-sensing technology has been used in a variety of coral reef studies, including the classification
70 of benthic habitats (Zhang 2015; Roelfsema et al. 2018; Zhang et al. 2018; Molder et al. 2022), detection of
71 changes in coral composition (Andréfouët et al. 2007; Hedley et al, 2016; Parsons et al. 2018; Purkis 2018;
72 Nimalan et al. 2021) and benthic reflectance at the bottom (Lee et al. 2013; Barnes et al. 2014; Thompson et al.
73 2017; Roelfsema et al. 2021) and estimation of coastal water bathymetry (Lyzenga et al. 2006; Mishra et al. 2006;
74 Pacheco et al. 2015; Minghelli et al. 2021). Frequently, these studies only analyze changes in benthic habitat cover
75 without identifying habitat status and types of threats.

76 To provide adequate coral data, this comprehensive study explores the ability of Landsat Satellites to
77 detect coral reef cover, using four multi-temporal Landsat that have not yet been applied to this region. Our study
78 illustrates the ability of Landsat satellite multi-sensor technology, combined with field data, to monitor changes in
79 coral cover over the past 25 years (from 1994 to 2019) and identify the threat status of coral reef ecosystems.

2. Materials and Methods

2.1. Study Site

The study site was located in the TSCA, Muna Barat Regency, in the Southeast Sulawesi Province of Indonesia, geographically located between 04°16'40"S – 04°32'20"S and 122°13'35"E – 122°32'40"E (**Fig. 1**). The Tiworo Strait was declared a MPA through a Decree of the Muna Regent (No.157/2004) and has a total area of 27,936 ha. For the last several decades, fishermen, fishing boat workers, seaweed and fish cage farmers, small-scale crab entrepreneurs, traders, and grocers living near the TSCA have depended on the marine resources in the Tiworo Strait.

2.2. Data Sources

2.2.1. Satellite Imagery Data

Landsat TM data consist of seven spectral bands with a spatial resolution of 30 m for bands 1–5 and 7. Landsat ETM+ data consist of eight spectral bands with a spatial resolution of 30 m for bands 1–7. Landsat 8 data have nine spectral bands with a spatial resolution of 30 m for bands 1–7 and 9. Landsat TM, ETM, ETM+, and OLI image classifications with a spatial resolution of 30 m can be used to examine the cover of five seabed substrates: live coral, rubble, sand, dead coral, and seagrass. Satellite imagery data used in this study were obtained from multi-temporal and multi-sensor Landsat imagery from 1994, 2002, 2013, and 2019. Landsat imagery data is freely downloadable from US Geological Survey-USGS (2019). The types and characteristics of Landsat imagery are summarized in Table 1.

2.2.2. Field Data

(i) Ground Control Points and Ground Truthing Data

Aside from coral observations, coordinate points as ground control points (GCPs) and for ground truthing (GT) were determined using global positioning system (GPS). GCPs at the location were typically road intersections, docks, or landmarks identifiable in the satellite images. GT data were focused on Pulau Tiga because

it has a large surrounding reef flat compared to other islands within the TSCA, making it easy for Landsat satellite sensors to capture spectral responses reflecting seabed cover. Coral reefs off Pulau Tiga can represent the condition of reefs in the TSCA at large because all of its islands can be accessed directly by local and non-local fishermen. GT data were obtained using GPS for each type of seabed cover, which included live corals, dead corals, rubble, seagrass, and sand (Nurdin et al. 2015). *In-situ* data on bottom cover types were obtained using a GPS unit (latitude-longitude data) and snorkeling or SCUBA equipment (data on bottom cover types). GT data were applied to check the accuracy of the image classification results.

(ii) Social Surveys

Socio-economic data were obtained through interviews and questionnaires as supporting information for image interpretation and Rapid Appraisal for Fisheries (RAPFISH) analysis. We used purposive sampling by selecting respondents according to types of fishery activities, occupations, academic positions, and stakeholder roles (Kothari 2004). To ensure the validity and currency of the data, we consulted with marine and fisheries scientists at the Faculty of Fisheries and Marine Science at Halu Oleo University. To strengthen the methodological assessment, we incorporated the opinions and perspectives of resource users (fishers), resource managers, and subject experts (Susilo 2003).

2.3. Data Analysis

2.3.1. Image Processing

Image processing was carried out in several phases, and refers to Nurdin et al. (2015) as follows:

a. Atmospheric Correction

Scattering and absorption of molecules by the atmosphere can decrease the quality of information in an image by up to 10%, depending on the spectral channels (Che and Price 1992). Atmospheric correction is therefore essential to minimize the effect of the atmosphere in multi-temporal images before comparing and analyzing the data (Hadjimitsis et al. 2010). Atmospheric correction was done for five series of Landsat imagery data using Dark Object Subtraction (DOS). In DOS, an absolute correction to the image is performed, such that

any reflection value (e.g., from shadows, deep clear water, or dense forest) is reduced to near zero percent. Then, signals recorded for each object by the sensor are a result of atmospheric scattering (Chavez 1996).

b. Geometric Correction

Geometric correction for three series of Landsat imagery data was conducted using the same coordinate points (GCPs). Geometric correction aims to improve the accuracy of and minimize the geometric error in Landsat imagery data. Eight GCP points were chosen for geometric correction using order polynomial transformation and a nearest-neighbor interpolation algorithm (Afwani and Danoedoro 2019). The corrected image was considered acceptable if the root mean square error was less than 0.5 (Baboo and Devi 2011).

c. Image Composites

True color in Landsat 5 TM and Landsat 7 ETM images is displayed with a RGB (red:green:blue) band composition of 3:2:1, whereas that in Landsat 8 OLI/TIRS images is displayed as RGB = 4:3:2 [38,40]. This band combination is often used to detect feature types on the bottom in shallow water at a preliminary stage. Three channels (bands) were selected because the band composites were considered the most appropriate for deducing the appearance of land cover. Red composites using band 3 (Landsat 5 TM/7 ETM) and band 4 (Landsat 8 OLI/TIRS) are suitable for detecting dry land or soil. Green composites were used to detect chlorophyll in vegetation from band 2 (Landsat 5 TM/7 ETM) and band 3 (Landsat 8 OLI/TIRS). High chlorophyll concentrations on the mainland provided high reflection digital values and were shown as dark green colors. Water was detected using band 1 (Landsat 5 TM/7 ETM) and band 2 (Landsat 8 OLI/TIRS) in blue composites so that it could be depicted in blue.

d. Image Subsetting (Cropping)

In image subsetting, or cropping, the aims are to delimit the area of interest, reinforce the geospatial phenomenon, and focus on the study area. In a subset image, objects appear larger such that existing information, such as color, can be seen more clearly.

158

159 *e. Application of the Depth Invariant Index*

160 The objective of the depth invariant index (DII) algorithm is to improve the accuracy of object
161 information (color) at the sea bottom in an image (Lyzenga 1981, 1985). We created at least 30 polygons in a
162 training area assumed to comprise coral reefs in the study area. The data obtained were analyzed in terms of
163 variance and covariance for blue and green bands. The variance and covariance were then used to determine the
164 attenuation coefficient to clarify the results of coral reef image classification using the following equations:

165 $$a = \frac{(\sigma ii - \sigma jj)}{2\sigma ij}, \quad (1)$$

166 $$\frac{ki}{kj} = a + \sqrt{(a^2 + 1)}, \quad (2)$$

167 $$(DII)ij = \ln(Li) - \left[\left(\frac{ki}{kj}\right) \times \ln(Lj)\right], \quad (3)$$

168 where i is the blue band, j is the green band, σ_{ii} is the variance of the blue band, σ_{jj} is the variance of the green
169 band, σ_{ij} is the covariance between blue and green bands, $\frac{ki}{kj}$ is the attenuation coefficient index ratio between
170 blue and green bands, and Li and Lj are the radiance pixels of blue and green bands, respectively.

171

172 *f. Classification, Ground Truth, and Reclassification*

173 Images produced by the DII algorithm were classified based on ISOCLASS unsupervised classification
174 (Call et al. 2003), resulting in 30 unlabeled classes with 100 iterations. These classification results were then
175 grouped according to their spectral and digital values. Each bottom type had a different spectral value identifiable
176 in the image. Reclassification was applied to the unsupervised-classification image based on visual interpretation
177 (spectral class color) and GT data with 75 sampling points in the field survey. Visual and field interpretations
178 were based on the interpretation guides by Suwargana (2014) and Nurdin et al. (2015), as seen in Table 2.

179

g. Post-Classification Process

At this stage, we produced maps of coral reef distribution and condition based on the extracted Landsat multi-temporal imagery data. Four maps were produced for 1994, 2002, 2013, and 2019, according to the time the images used were acquired. We also generated maps of changes in coral reef condition for the 25-year period using map overlay; coral reef changes from 1994 to 2002, 2002 to 2013, and 2013 to 2019 were mapped.

h. Accuracy Assessment

Overall accuracy is closely related to positional and thematic accuracy (Congalton and Green 2019), which is assessed using a confusion matrix. This method compares the image obtained from classification results (as the basis for the actual class) with field data, which are assumed to represent seabed cover. Data in rows were obtained from remote-sensing data classification, with the accuracy depending on the data producer, whereas data in columns were calculation results of field observations by researchers and were used in calculations of user accuracy. Greater consistency between classification and observation results would generate higher overall accuracy, which was calculated using the following equations:

$$Overall\ accuracy = \frac{\sum_{i=1}^{k} n_{ij}}{n} \times 100\%, \quad (4)$$

$$Producer\ accuracy\ j = \frac{n_{jj}}{n_{+j}} \times 100\%, \quad (5)$$

$$User\ accuracy\ i = \frac{n_{ii}}{n_{i+}} \times 100\%. \quad (6)$$

where n_{ij} is the number of observations at column j and row i, n_{ii} and n_{jj} are the number of observations categorized in the thematic class of i and j, respectively, n_{i+} and n_{+j} are the number of observations classified in the thematic class i from satellite data and j from *in-situ* data, respectively, and n is the total number of observations.

2.3.2. Rapid Appraisal for Fisheries Analysis

The Rapid Appraisal for Fisheries (RAPFISH) was developed at the University of British Columbia, Canada, and is used to evaluate fisheries sustainability with the application of a new multidisciplinary rapid assessment technique. The RAPFISH relies on the ordinance of assessed attributes that are grouped in several evaluation fields using multidimensional scaling (MDS). The fields or dimensions are ecology, economy,

206 technology, social, ethics/law, and institutions (Pitcher and Preikshot 2001; Alder et al. 2002; Fletcher 2006).
207 Each dimension is assigned a specific attribute, or indicator, associated with the threat to coral reefs, as
208 determined by Kennedy et al. (2013). Attributes are then parsed with reference to Pitcher and Preikshot (2001),
209 Susilo (2003), and RAPFISH (2019).

210 Technically, the stages of data collection and processing in RAPFISH are as follows: (i) identification
211 and determination of threat attributes, as defined according to the principles of the ecosystem approach to fisheries
212 management, (ii) definition and scoring of attributes, (iii) field verification and consultation with fisheries
213 scientists/experts, (iv) data processing, (v) scientific justification, and (vi) RAPFISH analysis (Pitcher 1999). The
214 RAPFISH procedure included data tabulation, data entry into the RAPFISH program, running the RAPFISH
215 program, analysis of multi-dimensional scaling (MDS), sensitivity analysis, Monte Carlo analysis, and
216 interpretation of results based on the threat index and sustainability status.

217

218 *a. Identification and Determination of Attributes*

219 Thirty-seven threat attributes were identified from a thorough review of all available data. The collected
220 attributes were classified into five dimensions: six attributes in the ecological dimension, eight attributes in the
221 economic dimension, seven attributes in the social dimension, nine attributes in the technological dimension, and
222 seven attributes in legal and institutional dimension (Table 3; (Pitcher and Preikshot 2001)). These attributes
223 represent the variation in threats to the sustainability of coral reefs in the TSCA.

224

225 *b. Defining and Scoring Attributes*

226 The second step in the analysis was defining and scoring attributes in RAPFISH (Pitcher and Preikshot 2001;
227 Kothari 2004). A "bad" score designates the worst possible condition of a coral reef ecosystem, and a "good" score
228 signifies the most favorable condition. Scoring was performed based on the method formulated by Good et al. (1999)
229 and Hershman et al. (1999).

230

231 *c. Feasibility of Threat Analysis*

Threat analysis feasibility is assessed by measuring stress values (particular measures of goodness-of-fit in MDS), which are calculated using the ALSCAL algorithm (Preikshot et al. 1998). This algorithm is used to optimize the distance of squared data to the original point (O_{ijk}) (d_{ijk}) in three-dimensional space (i, j, k), which is symbolized as stress (S), as described in the following equation:

$$S = \sqrt{\frac{1}{m}\sum_{k=1}^{m}\left[\frac{\sum_i\sum_j(d^2_{ijk}-\hat{d}^2_{ijk})^2}{\sum_i\sum_j O^4_{ijk}}\right]}, \quad (7)$$

where m is the number of analysts or experts from various scientific backgrounds. In the RAPFISH method, MDS analysis is terminated when the S value lies within a tolerable range, with an acceptable maximum value of 0.25 (25%) (Fauzi and Anna 2002).

d. Rotation Process

The projection of points onto the horizontal axis was done using a rotation process in which “bad” extreme positions were assigned to 0 and “good” extreme positions were assigned to 1. This process was carried out using RAPFISH scores. In addition, RAPFISH includes two "half-way" scores that are mirror images of each other for scaling in the vertical dimension and a set of pre-defined anchor points to avoid vertical "flipping" of the MDS ordinates. A more detailed description is provided in Kavanagh and Pitcher (2004).

e. Index Scale of Threats

We used a modified index scale to assess threats to the sustainability of coral reef ecosystems based on Pitcher and Preikshot (2001). Modifications included the following changes to descriptions of index levels: from unsustainable to very high threats, from less sustainable to high threats, from somewhat sustainable to fewer threats, and from sustainable to no threats. The threat index ranged from 0 to 100 and was divided into four categories (Table 4).

f. Sensitivity Analysis

Sensitivity analysis was done to determine which attribute was most sensitive to the sustainability of coral

reef ecosystems. The effect level of each attribute on the sustainability index was analyzed using leverage analysis in RAPFISH to determine the degree to which the ordination result changed when particular attributes were omitted from the dataset. The effect of each attribute was determined by observing the change in root squared correlation (RSQ) of the ordination, especially at the x-axis or accountability scale. Transformation of the RSQ at higher levels due to omission of a particular attribute causes a greater effect on the threat index. In other words, such an attribute is more sensitive to how a coral reef ecosystem is managed. Attributes with higher levels of sensitivity were subsequently used to formulate recommendations for saving coral reef ecosystems (Adiga et al. 2016).

g. Monte Carlo Analysis

Monte Carlo analysis was used to analyze errors in the assessment of attributes caused by several factors, such as 1) inconsistency in assessment, 2) differences in the assessment, 3) level of iterative stability, 4) data errors, and 5) high stress values (Kavanagh and Pitcher 2004). This method was also used to calculate the effect of random errors on the process of estimating/calculating ordination values. The acceptable stress value was set to <25% (Kavanagh and Pitcher 2004).

3. Results and Discussion

3.1. Change in Coral Reefs

The digital number obtained by Landsat multi-sensors is relatively small for the categories of live coral, dead coral, and seagrass, implying higher absorption of visible spectrum radiation (i.e., blue, green, and red channels) by the seabed compared to rubble and sand (**Fig. 2**). Therefore, the three categories appear darker in an image than does rubble or sand. Regression analysis results showed that the ratio of the blue channel vs. the green channel was the highest (RSQ > 94%) compared to other ratios in the visible spectrum. This is coincident with the results in numerous previous studies stating that the ratio of blue to green channels has a better ability to detect seabed cover, including coral reefs (El-Askary et al. 2014; Lillesand et al. 2015; Nurdin et al. 2015). Therefore, blue and green channels in the Landsat imagery data were used to detect seabed cover (**Fig. 3**).

283 Based on the results of an accuracy test using a confusion matrix, an accuracy test on the classification
284 results of Landsat 8 OLI/TIRS images from 2019 was carried out with field data obtained in 2019 for five classes
285 of seabed cover: live corals, dead corals, rubble, seagrass, and sand. The test yielded a producer accuracy of 83%,
286 user accuracy of 82%, and overall accuracy of 82%. This indicates that the interpretation results are considered
287 acceptable because the accuracies are greater than 80% (Lillesand et al. 2015).
288 The Landsat multi-temporal images from 1994, 2002, 2013 and 2019 showed that coral reefs have
289 changed over the 25-year period from 1994 to 2019, with a decline in live coral and increased dead coral and
290 rubble (**Fig. 4**). In 1994, the area for each category was 78.30 ha for live coral, 15.43 ha for dead coral, 20.07 ha
291 for rubble, 53.06 ha for seagrass, and 49.41 ha for sand, implying relatively small proportions of damaged coral
292 reefs (7.13%) and rubble (9.28%) compared to live coral (36.20%) (**Fig.4(a)**). The dominance of live coral is
293 considered to benefit the marine ecosystem and local people. Live coral was widely distributed (in cyan) around
294 the island, although some dead coral (in red) and rubble (in dark green) were also found in a small area.
295 There were significant changes in seabed cover from 1994 to 2003, particularly for live and dead coral
296 and rubble (**Fig. 4(b)**; Table 5). The live coral area decreased significantly by 41.95%, from 78.30 ha in 1994 to
297 45.45 ha in 2003. Dead coral doubled from 15.43 ha to 30.33 ha during the same period. The rubble area slowly
298 increased by 48.43%, from 20.07 ha in 1994 to 29.79 ha in 2003. Seagrass decreased slightly by 3.10 ha, whereas
299 sandy areas substantially increased from 49.41 ha in 1994 to 54.54 ha in 2003.
300 By 2013, live coral area had decreased, whereas dead corals had become dominant, especially off the
301 western and southern parts of the island (**Fig.4(c)**). The remaining live coral area was estimated to be 22.32 ha,
302 whereas dead coral cover doubled from 30.33 ha in 2003 to 61.35 ha in 2013 (Table 5). The rubble area increased
303 from 29.79 ha in 2003 to 32.13 ha in 2003, whereas the cover of seagrass and sand decreased slightly.
304 By 2019, live coral cover had declined to 8.01 ha (Table 5). Dead coral and rubble had slowly increased
305 to 66.42 and 33.66 ha, respectively. The areas of seagrass and sand fell slightly in the same period. Dead coral and
306 rubble occupied 30.71% and 15.56%, respectively, of the total area of the intertidal island (**Fig. 4(d)**).
307 We observed three types of changes in benthic habitat cover over the 25-year period from 1994 to 2019:
308 an increase, decrease, and fluctuative increase (Table 6). Visually, the greatest decrease in live coral cover

occurred between 1994 and 2003, with a reduction of 41.95%, followed by 2003–2013 (29.54%) and 2013–2019 (18.28%). Conversely, the greatest increase in dead coral cover occurred between 2003 and 2013 (201.04%), followed by 1994–2003 (96.57%) and 2013–2019 (32.86%). Rubble cover increased throughout the study period, although the rate gradually decreased. Seagrass exhibited fluctuations in cover: a decrease of 5.84% from 1994 to 2003, a slight increase of 3.79% from 2003 to 2013, and a decrease of 2.60% from 2013 to 2019. Similar to seagrass, sand cover fluctuated with a 10.38% increase from 1994 to 2003, a 16.64% decrease from 2003 to 2013, and a 12.81% increase from 2013 to 2019.

Live coral decreased dramatically by 89.77% from 78.30 ha in 1994 to 8.01 ha in 2019. By contrast, dead coral and rubble increased at an average rate of 2.19 ha/year for dead coral and 0.54 ha/year for rubble. This means that the loss in coral reefs during the period was 70.29 ha, or 2.81 ha/year.

Changes in habitat cover were be classified as follows: 1) live to dead coral, 2) dead coral to rubble, 3) rubble to sand, and 4) unidentified (**Fig. 5**). The largest percentage of change was from live to dead coral (72.54%), followed by dead coral to rubble (19.33%), rubble to sand (4.61%), and unidentified (3.15%). The large amount of change from live to dead coral implies that anthropogenic activities led to high coral reef mortality.

3.2. MDS Analysis

Results of MDS analysis for the 37 threat attributes using the RAPFISH approach showed that the cumulative stress values for the five dimensions were all in the range of 0.11–0.12 (Table 8). As this range was below 0.25, the threat values were statistically valid and fulfilled statistical procedures with a significant interval (RSQ) of 97%. Overall, threats to the TSCA and its coral reefs come from human activities surrounding the area.

3.3. Threat Index of Coral Reefs and Sensitivity

The RAPFISH ordination results indicated that the coral reef threat level varied according to the threat dimension (**Fig. 6**). Threats to coral reefs from the ecological dimension contributed the most (16.87), followed by legal and institutional (26.83), economic (31.00), social (34.83), and technological (41.10) dimensions. Overall, the

333 average threat index was 30.13, indicating a high threat to coral reef ecosystems.

334 The leverage analysis indicated that the sensitivities of the threat attributes in the five dimensions varied
335 greatly (**Fig. 7**). In this study, attributes that were ranked first to third in terms of contribution to index value in each
336 dimension were selected as the most sensitive attributes. For the ecological dimension, the top three most sensitive
337 attributes were "state of fishing exploitation" with a sensitivity score of 7.59, followed by "state of coral destruction"
338 (7.50) and "predators on corals" (i.e., crown-of-thorns seastar) (6.73). For the economic dimension, the top three
339 most sensitive attributes were "profitability" (5.83), "price of fish caught by destructive fishing" (5.78), and
340 "fishermen's income level" (5.30). For the social dimension, "level of public awareness" had the highest sensitivity
341 score (6.33). The second and third highest scores were for "habits of dumping waste into the sea" (6.04) and
342 "education level" (5.59). For the technological dimension, the three most sensitive attributes of nine attributes were
343 "intensity of blast fishing for the last 10 years" (5.36), followed by "intensity of trawl net fishing for the last 10
344 years" (5.17) and "ratio of fishing ground to coral reef area" (5.12). Of the seven attributes analyzed in the legal and
345 institutional dimension, those with the highest index scores were "involvement of law officers" (6.27), "lack of law
346 enforcement" (6.21), and "lack of monitoring and supervision" (5.50).

347 The 15 attributes were classified according to their high sensitivity values within each dimension for
348 consideration among the TSCA's stakeholders regarding coral reef management. These 15 attributes were also the
349 basis for developing technical programs and activities for conservation.

350 ### *3.4. Multi-dimensional Sustainability Index*

351 The results of the threat analysis for shallow water habitats in the TSCA showed that index values for
352 each dimension varied between very high threat (0.00–25.00) and high threat (25.01–50.00) levels. The indices for
353 MDS are visually depicted in **Fig. 8**. Threat scores for coral reefs assessed as facing severe conditions according
354 to threat indices included contributions from the ecological dimension (16.87 = very high threat), economic
355 dimension (31.00 = high threat), social dimension (34.83 = high threat), technological dimension (41.10 = high
356 threat), and law and institutional dimension (26.83 = high threat).

357 ### *3.5. Causal-loops Diagrams (CLDs) Model*

In general, these 15 threat attributes are presented in the causal-loops diagram model of coral reef ecosystem sustainability in the TSCA (**Fig. 9**). Causal-loops diagrams (CLDs) are used to qualitatively model causal relationships among a set of variables in a system; these diagrams capture our dynamic hypotheses and communicate essential feedback loops. An arrow with a "+" indicates that the effect increases when the cause goes up. On the other hand, an arrow with "**-**" indicates that when the cause increases, the effect decreases or harms the affected variable.

As shown in **Fig. 9**, there are three main CLDs variables: threats to coral reef ecosystems, sustainability of coral reef ecosystems, and changes in coral cover. In general, massive threats to coral reefs (i.e., ecological, economic, social, technological, and legal and institutional attributes) can affect the sustainability of coral reef ecosystems. Therefore, an appropriate strategy is needed to reduce pressure on coral reef ecosystems at the study site, ultimately increasing the percentage of coral cover as an indicator of healthy coral reefs.

3.6. Strategies to reduce threats to coral reef ecosystems

Long-term changes in the TSCA's coral reefs during the 25-year period from 1994 to 2019 have influenced the ecological, economic, and social sustainability of the surrounding communities. Coral reefs off Pulau Tiga and its surrounding areas are under "high threat," which means that the reefs and fisheries in the region are in danger. Overall, the TSCA is classified in the "high threat" category, with an average index value of 30.16 (**Fig. 8**). The most highly sensitive and influential threat index was the ecological dimension, with an index value of 16.87. Attributes of the ecological dimension must be seriously considered by all stakeholders if restoration of the TSCA's coral reefs were to occur. Critical attributes of the other dimensions i.e., law and institutional, economic, social, and technological must be addressed with appropriate policies and programs to prevent further degradation of the TSCA's coral reef ecosystems. We offer the following suggestions based on our results to reduce the threat status from highly threatened to threatened, and even to low threat levels.

Better Law Enforcement

Fair and firm law enforcement of illegal and destructive fishing activities is necessary. This is based on

our observations at the study site of fishermen who still use bombs and trawl nets. The intensive use of bombs and trawling nets not only damages coral reefs but also depletes and damages aquatic biota (Kularatne 2020). These destructive fishing activities are enabled by a ready supply of materials and a lack of monitoring facilities and personnel.

Improvement of Primary Education Facilities and Infrastructure

Lack of education and awareness within the local communities is a distinct, persistent problem. In some cases, destructive fishing behaviors are taken up by those with little education or awareness and no alternative livelihoods (Destructive Fishing Watch 2003). There is a shortage of teachers and there are no high schools in the TSCA area. More than 80% of the local fishers possess only a junior high school education level (Statistics of Muna Barat Regency 2019). Low levels of education significantly affect the understanding of coral reef resources and result in misconduct. Higher levels of education may promote a better understanding of the principles of sustainable coral reef management.

Enhancement of Community Income and Fishermen's Welfare

In addition to market and supply factors, low wages earned by fishermen are a strong contributing factor to destructive fishing practices. The imbalance of profits received by fishermen and intermediary buyers/traders is problematic and must be examined. Raising community incomes through collaborative management arrangements between the community (fishermen) and local government should be pursued. In some countries, such as Fiji, Chile, St. Lucia, Brazil, the Philippines, and other Pacific countries, collaborative management has been considered a success, especially in small-scale fisheries (d'Armengol et al. 2018). Success depends primarily on i) conducive social and institutional conditions, ii) market access and user dependence on resources, iii) local institutional characteristics, and iv) equitable distribution of benefits in the use of resources.

Regulation of Ocean Pollution from the Land

Agricultural land use and mangrove forest conversion in the Tondasi coastal area contributes to its

409 increasing turbidity and sedimentation. Besides muddiness, increased turbidity in the Tondasi coastal area affects
410 shallow water habitats, particularly coral reefs, off the surrounding islands. Most coral reefs off the surrounding
411 islands (near the mainland) were found to be mostly dead, disrupting their ability to function as habitat for fish
412 and other marine organisms. Implementation of coral reef rehabilitation programs for habitat recovery in this area
413 will be difficult because one of the limiting factors for healthy coral growth is an excessive concentration of
414 suspended sediments (>10 mg/L; (Erftemeijer et al. 2012)).
415
416 *Strengthening Local Institutions and Role Models*
417 Strong interactions between local fishers and their leaders (role models) can influence positive
418 community behavior in terms of the use of natural resources. The conditions of coral reef ecosystems in the TSCA
419 indicate destructive practices that are difficult to avoid. Leaders and role models in the local communities who are
420 aware of the issues facing local fishers and the coral reef ecosystems should be given roles and responsibilities
421 (Ho et al. 2016). Thus far, the roles of local community leaders have been taken over by village heads appointed
422 by the local government, but most village heads do not have good environmental awareness. Strong local
423 institutions are needed to gather people who are concerned about the future of their marine resources.
424
425 *Conflict Resolution*
426 Managing and defining zones in the TSCA is necessary to accommodate various economic, social, and
427 environmental interests (Day et al. 2019). The Tiworo Strait as a marine conservation area has not yet
428 implemented a zoning system, such as core zones and sustainable fishing zones. The implementation of zones
429 needs to be accompanied by socialization within the local communities so that all parties support and follow the
430 guidelines.
431
432 **4. Conclusion**
433 There were significant changes in live coral, dead coral, and rubble cover in the TSCA during the 1994–
434 2019 period. By analyzing satellite images, our results showed that the area of live coral decreased dramatically

by 89.77%, from 78.30 ha in 1994 to 8.01 ha in 2019, with a degradation rate of 2.81 ha/yr. Our RAPFISH analysis shows that a multidimensional threat has affected coral reefs during this period with an average threat index value of 30.13, which means that the threat status of coral reefs is categorized as high. This can be seen in the ecological dimension was the most sensitive dimension and was strongly influenced by threat level, with a threat index of 16.87 (very high threat status). If local threats continue without preventative measures, coral reefs will undoubtedly go extinct. In the future, sustainable management of coral reefs should be the primary goal for all stakeholders concerned with implementing the analyzed set of strategies. A multi-stakeholder collaboration model in the area can be the best option to improve coordination and cooperation in minimizing damage to coral reefs.

ACKNOWLEDGMENTS

We would like to thank Professors Muhammad Zamrun and La Sara for their support. This study was funded by the Halu Oleo University research program (no. 1584j/UN29.20/PPM/2019). We would also like to acknowledge support from the Integrated Research Program for Advancing Climate Models (TOUGOU Program) of the Ministry of Education, Culture, Sports, Science, and Technology of Japan (MEXT).

Conflicts of Interest: The authors confirm that they have no conflicts of interest to declare in relation to this submission.

454 **References**

455 Adam TC, Burkepile DE, Holbrook SJ et al (2021) Landscape-scale patterns of nutrient enrichment in a coral reef
456 ecosystem: implications for coral to algae phase shifts. Ecological Applications, 31(1):e2227.
457 https://doi.org/10.1002/eap.2227

458 Adiga MS, Ananthan PS, Kumari HVD, Ramasubramanian V (2016) Multidimensional analysis of marine fishery
459 resources of Maharashtra, India. *O*cean & Coastal Management, 130:13–20.

460 Afwani MZ, Danoedoro P (2019) The effects of polynomial interpolation and resampling methods in geometric
461 correction on the land-cover classification accuracy of Landsat-8 OLI imagery: a case study of Kulon Progo
462 area, Yogyakarta. Sixth Geoinformation Science Symposium 11311, 113110T.

463 Alder J, Zeller D, Pitcher T, Sumaila R (2002) A method for evaluating marine protected area management.
464 Coastal Management, 30(2):121–131. https://doi.org/10.1080/089207502753504661

465 Allemand D, Osborn D (2019) Ocean acidification impacts on coral reefs: From sciences to solutions. Regional
466 Studies in Marine Science, 28, 100558.

467 Andréfouët S, Hochberg EJ, Chevillon C et al. (2007) Multi-scale remote sensing of coral reefs. In Remote
468 sensing of coastal aquatic environments (pp. 297–315). Springer.

469 Andréfouët S, Kramer P, Torres-Pulliza D et al (2003) Multi-site evaluation of IKONOS data for classification of
470 tropical coral reef environments. Remote Sensing of Environment, *88*(1–2):128–143.
471 https://doi.org/10.1016/j.rse.2003.04.005

472 Baboo DSS, Devi MR (2011) Geometric Correction in Recent High Resolution Satellite Imagery: A Case Study in
473 Coimbatore, Tamil Nadu. International Journal of Computer Applications, 14(1):32–37.
474 https://doi.org/10.5120/1808-2324

475 Barnes BB, Hu C, Cannizzaro JP et al (2014) Estimation of diffuse attenuation of ultraviolet light in optically
476 shallow Florida Keys waters from MODIS measurements. Remote Sensing of Environment, 140:519–532.

477 Burke L, Selig L, Spalding M, Perry A (2011) *Reefs at risk Revisited.* https://www.icriforum.org/wp-
478 content/uploads/2019/12/factsheet_reefs_southeast_asia.pdf. Accessed 15 July 2019.

479 Call KA, Hardy JT, Wallin DO (2003) Coral reef habitat discrimination using multivariate spectral analysis and
480 satellite remote sensing. International Journal of Remote Sensing, 24(13):2627–2639.

481 Cesar H, Chong CK (2004) Economic Valuation and Socioeconomics of Coral Reefs : Methodological Issues and
482 Three Case Studies 1. *World Fish Center Conference Proceedings,* (1721):14–40.

483 Chan A, Hodgson PA (2017) A Systematic Analysis of Blast Fishing in South-East Asia and Possible Solutions.
484 IEEE Underwater Technology (UT), 1–6. https://doi.org/DOI: 10.1109/UT.2017.7890330

485 Chavez PS (1996) Image-based atmospheric corrections - Revisited and improved. Photogrammetric Engineering
486 and Remote Sensing, 62(9):1025–1036.

487 Che N, Price JC (1992) Survey of radiometric calibration results and methods for visible and near infrared
488 channels of NOAA-7,-9, and-11 AVHRRs. Remote Sensing of Environment, 41(1):19–27.

489 Chen J, Chen S, Fu R et al (2022) Remote Sensing Big Data for Water Environment Monitoring: Current Status,
490 Challenges, and Future Prospects. Earth's Future, 10(2):1–33. https://doi.org/10.1029/2021ef002289

491 d'Armengol L, Prieto Castillo M, Ruiz-Mallén I, Corbera E (2018) A systematic review of co-managed small-

scale fisheries: Social diversity and adaptive management improve outcomes. Global Environmental Change, 52:212–225. https://doi.org/10.1016/j.gloenvcha.2018.07.009

Day JC, Kenchington RA, Tanzer JM, Cameron DS (2019) Marine zoning revisited: How decades of zoning the Great Barrier Reef has evolved as an effective spatial planning approach for marine ecosystem-based management. Aquatic Conservation: Marine and Freshwater Ecosystems, 29:9–32.

De K, Nanajkar M, Arora M, Nithyanandan M, Mote S, Ingole B (2021) Application of remotely sensed sea surface temperature for assessment of recurrent coral bleaching (2014–2019) impact on a marginal coral ecosystem. Geocarto International. https://doi.org/10.1080/10106049.2021.1886345

Destructive Fishing Watch (2003) Profile of Destructive Fishing in Spermonde Islands. http://dfw.or.id/wp-content/uploads/2011/publikasi/DF_REPORT_SPERMONDE.pdf. Accessed 13 June 2019.

El-Askary H, El-Mawla ASH, Li J, El-Hattab MM, El-Raey M (2014) Change detection of coral reef habitat using Landsat-5 TM, Landsat 7 ETM+ and Landsat 8 OLI data in the Red Sea (Hurghada, Egypt). International Journal of Remote Sensing, 35(6):2327–2346. https://doi.org/10.1080/01431161.2014.894656

Endean R, Stablum W (1973) A study of some aspects of the crwon of thorns starfish (Acanthaster planci) infestations of reefs of Australia great barrier reef. *Atoll Research Bulletin*, *161*

Erftemeijer PLA, Riegl B, Hoeksema BW, Todd PA (2012) Environmental impacts of dredging and other sediment disturbances on corals: a review. *M*arine Pollution Bulletin, 64(9):1737–1765.

FAO (1995) Code of Conduct for Responsible Fisheries. http://www.fao.org/3/a-v9878e.pdf. Accessed 17 July 2019.

Fauzi A, Anna S (2002) EVALUASI STATUS KEBERLANJUTAN PEMBANGUNAN PERIKANAN: APLIKASI PENDEKATAN RAPFISH (Studi Kasus Perairan Pesisir DKI Jakarta). *Jurnal Pesisir & Lautan*, *4*(3):43–55.

Fletcher WJ (2006) Frameworks for managing marine resources in Australia through ecosystem approaches: do they fit together and are they useful? Bulletin of Marine Science, 78(3):691–704.

Foo SA, Asner GP (2019) Scaling Up Coral Reef Restoration Using Remote Sensing Technology. Frontiers in Marine Science, 6(79):1–8. https://doi.org/doi:10.3389/fmars.2019.00079

Good N, Schafer JB, Konstan JA et al (1999) Combining collaborative filtering with personal agents for better recommendations. Aaai/Iaai, 439:1–8.

Gorris P (2016) Deconstructing the reality of community-based management of marine resources in a small island context in Indonesia. Frontiers in Marine Science, 3(120):1-15.

Häder DP, Banaszak AT, Villafañe VE et al (2020) Anthropogenic pollution of aquatic ecosystems: Emerging problems with global implications. Science of the Total Environment, 713, 136586.

Hadjimitsis DG, Papadavid G, Agapiou A et al (2010) Atmospheric correction for satellite remotely sensed data intended for agricultural applications: impact on vegetation indices. Natural Hazards and Earth System Sciences, 10(1):89–95.

Hampton-Smith M, Bower DS, Mika S (2021). A review of the current global status of blast fishing: Causes, implications and solutions. Biological Conservation, 262. https://doi.org/10.1016/j.biocon.2021.109307

Hazraty-Kari S, Tavakoli-Kolour P, Das RR et al (2021) Baseline assessment of coral diseases in an environmentally extreme environment of the northern Persian Gulf. Marine Pollution Bulletin, 171, 112707.

https://doi.org/10.1016/j.marpolbul.2021.112707
Hedley JD, Roelfsema CM, Chollett I et al (2016) Remote sensing of coral reefs for monitoring and management: a review. Remote Sensing, 8(2):118.
Hein MY, Vardi T, Shaver EC et al (2021) Perspectives on the Use of Coral Reef Restoration as a Strategy to Support and Improve Reef Ecosystem Services. Frontiers in Marine Science,1–13. https://doi.org/10.3389/fmars.2021.618303
Hershman MJ, Good JW, Bernd-Cohen T, et al (1999) The effectiveness of coastal zone management in the United States. Coastal Management, 27(2–3):113–138.
Hill J, Wilkinson C (2004) Methods for ecological monitoring of coral reefs. Australian Institute of Marine Science, Townsville, 117
Ho NTT, Ross H, Coutts J (2016) The influence of leadership in fisheries co-management: the case of Tam Giang lagoon, Vietnam. Society & Natural Resources, 29(12):1405–1420.
Hoegh-Guldberg O, Pendleton L, Kaup A (2019) People and the changing nature of coral reefs. In Regional Studies in Marine Science (Vol. 30). https://doi.org/10.1016/j.rsma.2019.100699
Hughes TP, Anderson, KD, Connolly SR et al (2018) Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. Science, 359(6371). https://doi.org/10.1126/science.aan8048
Hughes TP, Barnes ML, Bellwood DR et al (2017) Coral reefs in the Anthropocene. In Nature, 546 (7656). https://doi.org/10.1038/nature22901
Kavanagh P, Pitcher TJ (2004) Implementing Microsoft Excel software for Rapfish: a technique for the rapid appraisal of fisheries status.
Kennedy EV, Perry CT, Halloran PR et al (2013) Avoiding coral reef functional collapse requires local and global action. Current Biology, 23(10):912–918. https://doi.org/10.1016/j.cub.2013.04.020
KKJI-KP3K (2019). Marine Protected Area Data. http://kkji.kp3k.kkp.go.id/index.php/en/marine-protected-area-data/details/7/92. Accessed 1 Sept.2019.
Kothari CR (2004) Research Methodology; Methods and Techniques (second revised edition). New Age International Publishers.
Kularatne RKA (2020) Unregulated and illegal fishing by foreign fishing boats in Sri Lankan waters with special reference to bottom trawling in northern Sri Lanka: A critical analysis of the Sri Lankan legislation. Ocean & Coastal Management, 185, 105012.
Lapointe BE, Brewton RA, Herren LW, Porter JW, Hu C (2019) Nitrogen enrichment, altered stoichiometry, and coral reef decline at Looe Key, Florida Keys, USA: a3-decade study. Marine Biology, 166(108):107–138. https://doi.org/https://doi.org/10.1007/s00227-019-3538-9
Lee Z, Weidemann A, Arnone R (2013) Combined Effect of Reduced Band Number and Increased Bandwidth on Shallow Water Remote Sensing: The Case of World View 2. IEEE Transactions on Geoscience and Remote Sensing, 51(5):2577–2586.
Lillesand T, Kiefer RW, Chipman J (2015) Remote sensing and image interpretation. John Wiley & Sons.
Lyzenga DR (1981) Remote sensing of bottom reflectance and water attenuation parameters in shallow water using aircraft and Landsat data. International Journal of Remote Sensing, *2*(1):71–82.
Lyzenga DR (1985) Shallow-water bathymetry using combined lidar and passive multispectral scanner data.

International Journal of Remote Sensing, 6(1):115–125.

Lyzenga DR, Malinas NP, Tanis FJ (2006) Multispectral bathymetry using a simple physically based algorithm. IEEE Transactions on Geoscience and Remote Sensing, 44(8):2251–2259.

Moussa MR, Fogg L, Bertucci F et al (2019) Long-term coastline monitoring on a coral reef island (Moorea, French Polynesia). Ocean and Coastal Management, 180, 104928. https://doi.org/10.1016/j.ocecoaman.2019.104928

Mehvar S, Filatova T, Dastgheib A et al (2018) Quantifying economic value of coastal ecosystem services: A review. Journal of Marine Science and Engineering, 6(1). https://doi.org/10.3390/jmse6010005

Mellin C, Macneil AM, Cheal AJ, Emslie MJ, Caley JM. (2016) Marine protected areas increase resilience among coral reef communities. Ecology Letters, 19(6):629–637. https://doi.org/10.1111/ele.12598

Minghelli A, Vadakke-Chanat S, Chami M et al (2021) Estimation of bathymetry and benthic habitat composition from hyperspectral remote sensing data (Biodiversity) using a semi-analytical approach. Remote Sensing, 13(10). https://doi.org/10.3390/rs13101999

Mishra D, Narumalani S, Rundquist D, Lawson M (2006) Benthic habitat mapping in tropical marine environments using QuickBird multispectral data. Photogrammetric Engineering & Remote Sensing, 72(9):1037–1048.

Molder EB, Schenkein SF, McConnell AE, Benedict KK, Straub CL (2022) Landsat Data Ecosystem Case Study: Actor Perceptions of the Use and Value of Landsat. Frontiers in Environmental Science, 9:1–19. https://doi.org/10.3389/fenvs.2021.805174

Ngoc QTK (2019) Assessing the value of coral reefs in the face of climate change: The evidence from Nha Trang Bay, Vietnam. Ecosystem Services, 35:99–108.

Nichols CR, Zinner J, Young DR (2019) Degradation of coastal ecosystems: causes, impacts and mitigation efforts. In Tomorrow's Coasts: Complex and Impermanent (pp. 119–136). Springer.

Nimalan K, Thanikachalam M, Usha T (2021) Spectral characteristics of coral reef benthic compositions in Gulf of Mannar. Journal of Earth System Science, 130(1). https://doi.org/10.1007/s12040-021-01571-9

Nurdin N, Komatsu T, Akbar AS, Djalil AR, Amri K (2015) Multisensor and multitemporal data from Landsat images to detect damage to coral reefs, small islands in the Spermonde archipelago, Indonesia. Ocean Science Journal, 50(2):317–325.

Pacheco A, Horta J, Loureiro C, Ferreira (2015) Retrieval of nearshore bathymetry from Landsat 8 images: A tool for coastal monitoring in shallow waters. Remote Sensing of Environment, 159:102–116. https://doi.org/10.1016/j.rse.2014.12.004

Pagiola S, Ritter KV, Bishop J (2004) Assessing the Economic Value of Ecosystem Conservation. In Environmental Economic Series. http://www.cbd.int/doc/case-studies/inc/cs-inc-iucn-nc-wb-en.pdf

Parsons M, Bratanov D, Gaston KJ, Gonzalez F (2018) UAVs, hyperspectral remote sensing, and machine learning revolutionizing reef monitoring. Sensors (Switzerland), 18(7):1–20. https://doi.org/10.3390/s18072026

Pendleton LH, Hoegh-Guldberg O, Langdon C, Comte A (2016) Multiple Stressors and Ecological Complexity Require a New Approach to Coral Reef Research. Frontiers in Marine Science, 3:1–5. https://doi.org/10.3389/fmars.2016.00036

609 Pitcher TJ (1999) Rapfish, a rapid appraisal technique for fisheries, and its application to the Code of Conduct for
610 Responsible Fisheries. *FAO Fisheries Circular (FAO)*.
611 Pitcher TJ, Preikshot D (2001) RAPFISH: a rapid appraisal technique to evaluate the sustainability status of
612 fisheries. Fisheries Research, 49(3):255–270.
613 Preikshot D, Nsiku E, Pitcher TJ, Pauly D (1998) An interdisciplinary evaluation of the status and health of
614 African lake fisheries using a rapid appraisal technique. Journal of Fish Biology, 53:381–393.
615 Purkis SJ (2018) Remote sensing tropical coral reefs: The view from above. Annual Review of Marine Science,
616 10:149–168.
617 Randazzo-Eisemann Á, Garza-Pérez JR, Penié-Rodriguez I, Figueroa-Zavala B (2021) 25 years of multiple
618 stressors driving the coral-algae phase shift in Akumal, Mexico. Ocean & Coastal Management, 214,
619 p.105917. https://www.sciencedirect.com/science/article/abs/pii/S0964569121004002
620 RAPFISH (2019) *RAPFISH: A Rapid Appraisal for Fisheries*. http://www.rapfish.org/home. Accessed 15 July
621 2019
622 Roelfsema C, Kovacs EM, Markey K et al (2021) Benthic and coral reef community field data for Heron Reef,
623 Southern Great Barrier Reef, Australia, 2002–2018. Scientific Data, 8(1):1–7.
624 https://doi.org/10.1038/s41597-021-00871-5
625 Roelfsema C, Kovacs E, Ortiz JC et al (2018) Coral reef habitat mapping: A combination of object-based image
626 analysis and ecological modelling. Remote Sensing of Environment, 208:27–41.
627 Russell GC, Green K (2019) Assessing the Accuracy of Remotely Sensed Data Principles and Practices, Third
628 Edition. In *CRC Press* (Vol. 1, Issue 9).
629 Santavy DL, Horstmann CL, Sharpe LM, Yee SH, Ringold P (2021) What is it about coral reefs? Translation of
630 ecosystem goods and services relevant to people and their well-being. Ecosphere, 12(8).
631 https://doi.org/10.1002/ecs2.3639
632 Statistics of Muna Barat Regency. (2019) *Kabupaten Muna Barat Dalam Angka 2018*. Muna Barat Regency.
633 https://munbarkab.bps.go.id/. Accessed 28 May 2019
634 Susilo BS (2003) Small islands development sustainability: A case study in Kelurahan Pulau Panggang and Pulau
635 Pari, Kepulauan Seribu, DKI Jakarta. In Bogor Agriculture University 3(2).
636 Suwargana N (2014) *Analysis of Alos AVNIR-2 Imagery for coral reefs mapping (a case study: Banyuputih,*
637 *Situbondo) In: Proceedings of national seminar on remote sensing.* Indonesian National Institute of
638 Aeronautics and Space (LAPAN). http://sinasinderaja.lapan.go.id/wp-
639 content/uploads/2014/06/bukuprosiding_588-956.pdf. Accessed 20 June 2020
640 Thompson DR, Hochberg EJ, Asner GP et al (2017) Airborne mapping of benthic reflectance spectra with
641 Bayesian linear mixtures. Remote Sensing of Environment, 200:18–30.
642 US Geological Survey-USGS (2019) *Data sets*. The White House, Washington DC.
643 https://doi.org/DOI_Privacy@ios.doi.gov. Accessed 10 October 2019
644 Vaughan D, Agardy T (2020) Marine protected areas and marine spatial planning–allocation of resource use and
645 environmental protection. In Marine Protected Areas (pp. 13–35). Elsevier.
646 Yasir Haya LOM, Fujii M (2019) Assessing economic values of coral reefs in the Pangkajene and Kepulauan
647 Regency, Spermonde Archipelago, Indonesia. Journal of Coastal Conservation, 23(3):699–711.


Yasir Haya LOM, Fujii M (2017) Mapping the change of coral reefs using remote sensing and in situ measurements: a case study in Pangkajene and Kepulauan Regency, Spermonde Archipelago, Indonesia. Journal of Oceanography, 73(5):623–645. https://doi.org/10.1007/s10872-017-0422-4

Yasir Haya LOM, Fujii M (2020) Assessment of coral reef ecosystem status in the Pangkajene and Kepulauan Regency, Spermonde Archipelago, Indonesia, using the rapid appraisal for fisheries and the analytic hierarchy process. Marine Policy, 118. https://doi.org/10.1016/j.marpol.2020.104028

Zhang C (2015) Applying data fusion techniques for benthic habitat mapping and monitoring in a coral reef ecosystem. ISPRS Journal of Photogrammetry and Remote Sensing, 104:213–223.

Zhang C, Denka S, Cooper H, Mishra D R (2018) Quantification of sawgrass marsh aboveground biomass in the coastal Everglades using object-based ensemble analysis and Landsat data. Remote Sensing of Environment, 204:366–379.

Zunino S, Libralato S, Melaku CD, Prato G, Solidoro C (2021) Impact of Ocean Acidification on Ecosystem Functioning and Services in Habitat-Forming Species and Marine Ecosystems. Ecosystems, 24(7):1561–1575. https://doi.org/10.1007/s10021-021-00601-3



664 **Fig. 1** Study area located in the Tiworo Strait Conservation Area (TSCA) in the Muna Barat Regency of Southeast

665 Sulawesi in Indonesia.






Fig. 2 Spectral responses representing five types of seabed cover obtained by Landsat multi-sensor satellites in 1994 (Landsat 5 TM), 2003 (Landsat 7 ETM+), 2013 (Landsat 8 OLI/TIRS), and 2019 (Landsat 8 OLI/TIRS). Of the five types of seabed cover, it can be seen that the three classes with the lowest response values—live coral, dead coral, and seagrass—have object characteristics in which higher levels of electromagnetic radiation are absorbed compared to rubble and sand. This causes the digital number (DN) values of the three types to be lower than those of rubble and sand.



672 Fig. 3 Ratio of blue to green channels for multi-temporal Landsat imagery obtained in (a) 1994 (Landsat 5 TM),
673 (b) 2003 (Landsat 7 ETM+), (c) 2013 (Landsat 8 OLI/TIRS), and (d) 2019 (Landsat 8 OLI/TIRS). In the visible
674 light spectrum, because blue and green light can penetrate seawater further, the ratio between the blue and green
675 band is more informative for the sea-bottom compared to the red or other bands. In Landsat 5 TM and 7 ETM+,
676 band 1 is the blue band (0.45–0.52 μm) and band 2 is the green band (0.53–0.61μm), whereas in Landsat 8
677 OLI/TIR, band 2 is blue (0.450–0.515 μm) and band 3 is green (0.525–0.600 μm).



678 **Fig. 4** Distribution of changes in coral cover off Pulau Tiga within the TSCA according to Landsat multi-temporal

679 image data obtained over a 25-year period from 1994 to 2019.




Fig. 5 Composite of live coral change from 1994 to 2019 with four categories of change. In total, 89.77% of the coral reef area was transformed (left pie chart): 72.54% from live to dead coral, 19.33% from dead coral to rubble, 4.61% from rubble to sand, and unidentified changes (3.15%) (right pie chart).







Fig. 6 Two-dimensional Rapid Appraisal for Fisheries (RAPFISH) ordinations of the threats to coral reefs in the TSCA, based on (a) ecological, (b) economic, (c) social, (d) technological, and (e) legal and institutional dimensions. The horizontal axis represents sustainability [0% ("bad") to 100% ("good")] and the vertical axis represents changes in coral reef status (%).



687 **Fig. 7** Attribute leverage analysis of the RAPFISH ordinations for threats to coral reefs in the TSCA. From the
688 five dimensions analyzed (i.e., ecological, economic, social, technological, and legal and institutional), we
689 obtained a root squared correlation (RSQ) value of 0.97, indicating that the analysis was acceptable.
690



691 **Fig. 8** Kite diagram of the five threat dimensions to coral reef ecosystems in the TSCA. Based on the threshold
692 values listed in Table 4, the ecological dimension was regarded as under "very high threat" and the economic,
693 social, technological, and legal and institutional dimensions contributed to a "high threat" level. The overall value
694 for all threats was 30.13, indicating a "high threat" status of the coral reef ecosystems at the study site.



695 **Fig. 9** Causal loop diagram model in terms of threats to coral reef ecosystems and their sustainability strategies in
696 the Tiworo Strait Conservation Areas (TSCAs)
697

698 Table 1. Types and characteristics of Landsat image data

No.	Satellite	Sensor	Resolution (m)	Acquisition	Path/Row
1.	Landsat 5	TM	30	July 14, 1994	112/63
2.	Landsat 7	ETM	30	August 28, 2003	112/63
3.	Landsat 8	OLI/TIRS	30	April 29, 2013	112/63
4.	Landsat 8	OLI/TIRS	30	Sept. 21, 2019	112/63

699

700 Table 2. Descriptions and definitions of the bottom types surveyed (Suwargana, 2014; US Geological Survey-

701 USGS, 2019).

Classes	Visual description	Field defenitions
Live coral	Cyan to light green cloudy	Habitat dominated by a mix live scleratinian corals
Dead coral	Green with clear boundary	Substrate predominantly made of dead coral
Rubble	Dark orange with clear boundary	Pieces of broken corals, generally found in reef flats
Seagrass	Yellow to light orange with unclear boundary	Flowering plants, generally found intertidal areas and reef flats
Sand	Yellow and red	Sediment from carbonate origin (dead corals and skeleton for calcifying organisms)

702

703 Table 3. Attributes of threats to coral reef ecosystems and criteria for Rapid Appraisal for Fisheries (RAPFISH)
704 scoring. The scores ranged from bad (very high threat level; 0) to good (no threats), with the highest scores
705 varying among attributes.

Attribute	Score scale	Remarks	Source/ references
Ecological Dimension			
1. State of fishing exploitation	0,1,2,3,4	Collapsed (0); high (1); moderate (2); low (3); balanced or equal (4)	Pitcher (1999); Pitcher and Preikshot (2001)
2. Coral bleaching	0,1,2,3	% coral bleaching: ≥75-100% (0); 50-74.9% (1); 25-49.9% (2); <25% (3)	Modified from RAPFISH (2019)
3. State of coral reef destruction	0,1,2,3	% coral rubble: ≥75-100% (0); 50-74.9% (1); 25-49.9% (2); <25% (3)	Modified from Hill and Wilkinson (2004)
4. Change in the size of reef fish caught (for the last 10 years)	0,1,2	Declined substantially (0); decreased slightly (1); unchanged (2)	RAPFISH (2019)
5. Concentration of total suspended solid (TSS) in coral reefs	0,1	Threshold for coral growth: >10 mg/L (0); ≤10 mg/L (1)	Erftemeijer et al. (2012)
6. Predators on corals (i.e., crown-of-thorns starfish (COTs), starry puffer fish etc.)	0,1	COTs in coral reefs: >14 individuals/1000 m^2 (0); ≤14 individuals/ 1000 m^2 (1)	Endean and Stablum (1973); Hoeksema (2012)
Economic Dimension			
1. Profitability	0,1,2,3	High profit (0); marginal profit (1); without gain or loss (2); little loss (3); large loss (4)	RAPFISH (2019)
2. Fishermen's income level	0,1,2,3	Average relative income to the regional minimum wage (UMR): Far below (0); below (1); equal (2); higher (3); considerably higher (4)	Modified from RAPFISH (2019)
3. Accessibility to markets for swimming crabs	0,1,2	International market (0); national market (1); local market (2)	RAPFISH (2019)
4. Status of business capital	0,1,2,3	Loan capital (0); joint capital (1); grant capital (2); own capital (3)	Modified from RAPFISH (2019)
5. Marketing system of fish/swimming crabs	0,1,2,3	Monopoly or government buyer (0); semi-closed (1); partially open market (2); fully open market (3)	RAPFISH (2019)
6. Prices of raw materials for bombs &Trawl Net	0,1,2,3	Very affordable (0); affordable (1); expensive (2); very expensive (3)	Modified from RAPFISH (2019)
7. Prices of fish caught by destructive fishing	0,1,2,3,4	Prices compared to fish caught by non-destructive fishing: Very expensive (0); expensive (1); equal (2); cheap (3); Very cheap (4)	Modified from RAPFISH (2019)
8. Easy to get banking capital	0,1,2	Difficult (0); moderate (1); eazy (2)	Yasir Haya and Fujii (2019)
Social Dimension			
1. Knowledge level about marine conservation	0,1,2,3	None (0); low (1); moderate (2); high (3)	Modified from RAPFISH (2019)
2. Education level	0,1,2,3	The number of students who do not proceed to the university: 75-100% (0); 50-74.9% (1); 25-49.9% (2); <25% (3)	Statistics of Muna Barat Regency (2019)
3. Level of poverty and fixed income	0,1,2	High (0); moderate (1); low (2)	RAPFISH (2019)
4. Level of public	0,1,2	Low (0); moderate (1); high (2)	Modified from

awareness			RAPFISH (2019)
5. Habit of dumping wastes to the sea	0,1,2,3,4	Very high (0); high (1); moderate (2); low (3); very low (4)	Modified from RAPFISH (2019)
6. Frequency of conflicts (destructive versus non-destructive)	0,1,2	Frequent (0); seldom (1); none (2)	Modified from Susilo (2003)
7. Ability of making destructive fishing devices	0,1,2,3	High (0); moderate (1); low (2); none (3)	Modified from RAPFISH (2019)
Technological Dimension			
1. The number of vessels and fishing gears (for the last 10 years)	0,1,2	Increasing (0); unchanged (1); decreasing (2)	Statistics of Muna Barat Regency (2019)
2. Intensity of blast fishing for the last 10 years	0,1,2,3	High (0); medium (1); low (2); none (3)	RAPFISH (2019)
3. State of mangrove utilization (fire woods, construction, etc.)	0,1,2,3	High (0); medium (1); low (2); none (3)	RAPFISH (2019)
4. Selectivity of fishing gears	0,1,2,3	None (0); few (1); fair (2); many (3)	RAPFISH (2019)
5. Ratio of fishing grounds in coral reefs	0,1,2,3	75-100% (0); 50-74.9% (1); 25-49.9% (2); <25% (3)	Modified from RAPFISH (2019)
6. The number of vessels anchoring in coral reefs	0,1,2	Many (0); few (1); none (2)	Modified from RAPFISH (2019)
7. State of mangrove conversion (fishpond, settelement, buildings)	0,1,2,3	High (0); medium (1); low (2); none (3)	Modified from RAPFISH (2019)
8. Intensity of Trawl Net for the last 10 years	0,1,2,3	High (0); medium (1); low (2); none (3)	RAPFISH (2019)
9. Mariculture unfriendly	0,1,2,3	Very bad (0); bad (1); fair (2); good (3)	Modified from RAPFISH (2019)
Legal and Institutional Dimensions			
1. Lack of law enforcement	0,1,2,3	The level of legal violation: high (0); medium (1); low (2); none (3)	Modified from FAO (1995)
2. Overlapping policies between institutions	0,1,2,3	Numerous (1); fair (1); few (2); none (3)	Modified from FAO (1995)
3. Socialization and campaign of the public awareness	0,1,2	Never (0); fair (1); frequent (2)	Modified from RAPFISH (2019)
4. Involvement of law officers	0,1,2	Frequent (0); fair (1); never (2)	Modified from FAO (1995)
5. Lack of monitoring and supervision by community	0,1,2,3	Never (0); seldom (1); fair (2); frequent (3)	Modified from RAPFISH (2019)
6. Role models in the community	0,1,2	A public figure who understands the environmental issues: Few (0); fair (1); many (2)	Modified from RAPFISH (2019)
7. Knowledge and local wisdom	0,1,2	Little (0); fair (1); a lot (2)	Modified from Susilo (2003)

706

707 Table 4. Threshold index values for assessing the threats to coral reefs (modified from Pitcher and Preikshot 2001).

Threshold value of index	Status
0.00 - 25.00	Very high threats
25.01 - 50.00	High threats
50.01 - 75.00	Less threats
75.01 - 100.00	No threats

708

709 Table 5. Aerial estimates of major bottom types in 1994, 2003, 2013, and 2019 off Pulau Tiga within the Tiworo

710 Strait Conservation Area (TSCA).

Type of classes	Area of seabed cover (ha)			
	1994	2003	2013	2019
Live corals	78.30 (36.20%)	45.45 (21.02%)	22.32 (10.32%)	8.01 (3.70%)
Dead corals	15.43 (7.13%)	30.33 (14.02%)	61.35 (28.37%)	66.42 (30.71 %)
Rubbles	20.07 (9.28%)	29.79 (13.77%)	32.13 (14.86%)	33.66 (15.56 %)
Seagrass	53.06 (24.53%)	56.16 (25.97%)	54.15 (25.04%)	55.53 (25.68 %)
Sand	49.91 (22.85%)	54.54 (25.22%)	46.32 (21.42%)	52.65 (24.34 %)

711

712 Table 6. Changes in seabed cover from 1994 to 2003, 2003 to 2013, and 2013 to 2019.

Type of classes	Change of bottom types						Type of change
	1994 to 2003 (ha)	Ratio (%)	2003 to 2013 (ha)	Ratio (%)	2013 to 2019 (ha)	Ratio (%)	
Live corals	-32.85	-41.95	-23.13	-29.54	-14.31	-18.28	(-)
Dead corals	14.90	96.57	31.02	201.04	5.07	32.86	(+)
Rubbles	9.72	48.43	2.34	11.66	1.53	7.62	(+)
Seagrass	-3.10	-5.84	2.01	3.79	1.38	2.60	(-/+)
Sand	5.13	10.38	-8.22	-16.64	6.33	12.81	(-/+)

713

714 Table 7. Cumulative stress values and analysis of all dimensions. Values indicate the acceptability of the coral reef

715 status analysis [Stress (S) < 0.25, root squared correlation (RSQ) ≈ 1].

Dimension	Stress value	RSQ
Ecological	0.12	0.97
Economic	0.12	0.97
Social	0.11	0.97
Technological	0.12	0.97
Legal and institutional	0.12	0.97

716

717