



Title	Nickel-Catalyzed Defluorophosphonylation of Aryl Fluorides
Author(s)	You, Zhensheng; Masuda, Yusuke; Iwai, Tomohiro et al.
Citation	Journal of organic chemistry, 87(21), 14731-14737 https://doi.org/10.1021/acs.joc.2c02048
Issue Date	2023-11-09
Doc URL	https://hdl.handle.net/2115/90676
Rights	This document is the Accepted Manuscript version of a Published Work that appeared in final form in [JournalTitle], copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see https://pubs.acs.org/articlesonrequest/AOR-BEB28BHKFETDERJXCABA .
Type	journal article
File Information	J. Org. Chem._87(21)_14731-14737.pdf




Nickel-Catalyzed Defluorophosphonylation of Aryl Fluorides

Zhensheng You,² Yusuke Masuda,^{1,2} Tomohiro Iwai,^{2†} Kosuke Higashida,^{1,2*} and Masaya Sawamura^{1,2*}

¹ Institute for Chemical Reaction Design and Discovery (WPI-ICReDD), Hokkaido University, Sapporo, Hokkaido 001-0021, Japan.

² Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan.

† Present address: Graduate School of Arts and Sciences, The University of Tokyo, Meguro-ku, Tokyo 153-8902, Japan



ABSTRACT: A Ni-catalyzed cross-coupling reaction between aryl fluorides and dialkyl phosphonates [HP(O)(OR)₂] (R = secondary alkyl groups) in the presence of potassium *tert*-butoxide as a base is reported. The reaction converted various aryl fluorides into the corresponding aryl phosphonates even when electron-donating substituents were present on the aromatic ring. The combined experimental and computational studies suggested Ni–K⁺ cooperative action of a Ni(0) complex chelated with a strongly electron-donating ion-bridged dimeric phosphite ligand system [P(OR)₂O[−]K⁺]₂ that facilitates turnover-limiting C–F bond oxidative addition of aryl fluorides.

INTRODUCTION

The direct functionalization of C–F bonds has attracted increasing attention in recent years because of the increasing abundance of fluorinated compounds in pharmaceuticals¹ and other functional materials,² affording various opportunities for diversity-oriented synthesis through late-stage functionalization of complicated molecules. Whereas numerous methods for converting C–F bonds into C–C, C–H, C–O, and C–N bonds have been developed,³ few methods exist for forming weaker C–P bonds toward the synthesis of organophosphorus compounds. The classical approach to C–F to C–P bond transformation involves a nucleophilic aromatic substitution (S_NAr) reaction with phosphide anions,⁴ which are generated using hazardous alkali metals.⁵ Würthwein and coworkers have demonstrated that silylphosphines undergo S_NAr phosphination of aryl fluorides at high temperatures (>170 °C) (Scheme 1a).⁶ In these S_NAr-type reactions, the pronucleophiles are limited to secondary phosphines and phosphine oxides; the successful use of more electron-deficient phosphonic acid diesters [HP(O)(OR)₂] has not been reported. Organophosphonic acids [RP(O)(OH)₂] and their derivatives are widely used as drugs or pro-drugs,⁷ chelators of metallic salts,⁸ surface modifiers,⁹ and phosphoantigens.¹⁰ Thus, the development of a phosphonylation reaction for aryl fluorides is demanded. To the best of our knowledge,

however, the literature contains only one example of C(sp²)–F bond phosphonylation, which was achieved via a photoinduced single-electron-transfer (SET) process (Figure 1b)¹¹ that required irradiation with ultraviolet light (λ = 254 nm). We recently reported a nucleophile-dependent S_NAr reaction of non-activated aryl fluorides with potassium diorganophosphinites (R₂PO[−]K⁺) (Figure 1c).¹² In this reaction, the potassium cation plays a critical role, stabilizing the negative charge of the leaving fluoride anion. With this knowledge, we envisaged combining the unique property of the K⁺ cation with the well-established ability of a Ni catalyst to activate C–F bonds^{3,13} and thereby achieve phosphonylation of aryl fluorides (Figure 1d).¹⁴

Herein, we report a Ni-catalyzed cross-coupling reaction between aryl fluorides and dialkyl phosphonates [HP(O)(OR)₂] in the presence of potassium *tert*-butoxide (KO^tBu) as a stoichiometric base. The reaction uses commercially available Ni complexes as catalyst precursors and requires no exogenous ligand. Not only electron-deficient but also electron-neutral and even electron-rich aryl fluorides were successfully converted into the corresponding aryl phosphonates. Interestingly, the present coupling reaction proceeds specifically with di-*sec*-alkyl phosphonates; no reaction occurs with primary or tertiary alkyl phosphonates. Mechanistic studies by kinetic experiments and density functional theory (DFT) calculations suggested that the catalytic cycle involves turnover-limiting oxidative addition of

the aryl fluoride to a Ni(0) complex coordinated with potassium dialkyl phosphites [P(OR)₂O⁻K⁺], which occurs through cooperative action of a Ni(0)–K⁺ bimetallic system.




Figure 1. Defluorinative C–P bond formation reactions (KHMDS = potassium hexamethyldisilazide; KOtBu = potassium *tert*-butoxide).

RESULTS AND DISCUSSION

Specifically, the reaction between 4-fluorobiphenyl (**1a**, 0.125 mmol) and dicyclohexyl phosphonate (**2a**, 0.25 mmol) in the presence of NiBr₂·diglyme (5 mol%, 0.00625 mmol) and KOtBu (0.25 mmol) in toluene (0.5 mL) at 120 °C led to the clean and complete conversion of **2a** to the corresponding defluorinative phosphonylation product (**3a**) in quantitative yield (99% based on ¹H NMR spectroscopy) (Table 1, entry 1). When the amount of KOtBu was decreased to 0.125 mmol (1 equiv to **1a**, 0.5 equiv to **2a**), product **3a** was not obtained, suggesting the formation of an inactive Ni species through the direct reaction of a Ni species with nondeprotonated dialkyl phosphonate **2a** (entry 2).¹⁵ NiI₂ exhibited a catalytic performance similar to that of NiBr₂·diglyme (entry 3), whereas other Ni(II) complexes such as NiCl₂ and Ni(acac)₂ (acac = acetylacetonate) gave **3a** in moderate yields (entries 4–6). Notably, the Ni(0) complex Ni(cod)₂ (cod = 1,5-cyclooctadiene) also catalyzed the present reaction, suggesting that Ni(0) is an active species in the catalytic process (entry 7). No reaction occurred in the absence of a Ni catalyst (entry 8). The base strongly influenced the reaction efficiency. Specifically, the use of bases that have smaller cations (e.g., NaOtBu and LiOtBu) in place of KOtBu in entry 1 resulted in a substantial decrease in the product yield (entries 9 and 10). The addition of 18-crown-6 (2 equiv) inhibited the reaction completely, suggesting direct participation of a potassium cation in the catalysis (entry 11). The phosphonylation product **3a** was not obtained at all with less basic potassium salts such as K₂CO₃ and K₃PO₄ (entries 12 and 13). Potassium hexamethyldisilazide (KHMDS), which is sufficiently basic to

deprotonate dialkyl phosphonates, also failed to give **3a** (entry 14). By contrast, the use of KHMDS with Ni(cod)₂ instead of NiBr₂·diglyme afforded **3a** in a quantitative yield, suggesting that the reduction of a Ni(II) precatalyst occurred with *tert*-butoxide but not with KHMDS (entry 15). The addition of exogenous ligands did not substantially affect the product yield.¹⁶

Table 1. Ni-catalyzed phosphonylation of 4-fluorobiphenyl (**1a**) with dicyclohexyl phosphonate (**2a**).^a

Entry	[Ni]	Base	Yield [%] ^b
1	NiBr ₂ ·diglyme	KOtBu	99
2 ^c	NiBr ₂ ·diglyme	KOtBu	0
3	NiI ₂	KOtBu	99
4	NiBr ₂	KOtBu	55
5	NiCl ₂	KOtBu	62
6	Ni(acac) ₂	KOtBu	77
7	Ni(cod) ₂	KOtBu	70
8	none	KOtBu	0
9	NiBr ₂ ·diglyme	NaOtBu	33
10	NiBr ₂ ·diglyme	LiOtBu	2
11 ^d	NiBr ₂ ·diglyme	KOtBu	0
12	NiBr ₂ ·diglyme	K ₂ CO ₃	0
13	NiBr ₂ ·diglyme	K ₃ PO ₄	0
14	NiBr ₂ ·diglyme	KHMDS	0
15	Ni(cod) ₂	KHMDS	99

^a Reaction conditions: **1a** (0.125 mmol), **2a** (0.25 mmol), Ni complexes (0.00625 mmol, 5 mol% to **1a**), base (0.25 mmol), toluene (0.5 mL), 120 °C, 15 h. ^b ¹H NMR yield obtained using 1,3,5-trimethoxybenzene as an internal standard. ^c 0.125 mmol of base. ^d 18-Crown-6 (0.250 mmol, 2.0 equiv) was added.


acac = acetylacetonate, cod = 1,5-cyclooctadiene, HMDS = hexamethyldisilazide.

With the optimal conditions in hand, we investigated the substrate scope (Table 2). Aryl fluorides possessing electron-donating substituents such as methoxy (OMe), dimethylamino (NMe₂), diphenylamino (NPh₂), and methyl (Me) groups participated in the phosphonylation reaction to afford the corresponding aryl phosphonates in good yields (**3b–3e**). Simple fluorobenzene also exhibited excellent reactivity, forming **3f** in 90% yield. Electron-withdrawing amide groups were tolerated on the aromatic ring, and the corresponding products **3g** and **3h** were obtained in 86% and 74% yields, respectively. However, the aryl fluoride with a highly electron-withdrawing trifluoromethyl group resulted in a low yield of **3i** (15% yield). π -Extended aryl fluorides were suitable substrates for the present phosphonylation reaction, and the corresponding products were obtained in excellent yields (**3j–3l**). However, 2-fluorobiphenyl and 1-fluoro-2,6-dimethylbenzene did not give products **3m** and **3n**. Heteroaryl fluorides were successfully converted to the

corresponding phosphonates (**3o** and **3p**). For the synthesis of **3b,d,k**, the reaction was conducted with Ni(cod)₂ and KHMDS because the protocol with NiBr₂·diglyme and KO^tBu produced small amounts of phosphorus-containing byproducts, which hampered the isolation of the products in pure form.

Next, the scope of dialkyl phosphonates was examined. Diisopropyl phosphite (**2b**) afforded the corresponding phosphonate **3q** in 84% yield, whereas dialkyl phosphonates with tertiary (**2c**) or primary (**2d**) *O*-alkyl groups failed to give the products (**3r**, **3s**). Diphenyl phosphonate (**2e**) also exhibited no reactivity. Thus, the present reaction was specifically possible with phosphonates with two secondary *O*-alkyl groups. This unusual reactivity trend disfavoring the smallest pronucleophile [HP(O)(OEt)₂] can be explained by assuming the formation of inactive Ni species through over-coordination of the relatively small potassium diethyl phosphite ligand [P(OEt)₂O⁻K⁺] to a Ni


Table 2. Scope of aryl fluorides and dialkyl phosphonates.^a




^a Reaction conditions: **1** (0.125 mmol), **2** (0.25 mmol), NiBr₂·diglyme (0.00625 mmol, 5 mol% to **1**), KO^tBu (0.25 mmol), PhMe (0.5 mL), 120 °C, 20 h. Yields of isolated products are shown. ^b Ni(cod)₂ (0.00625 mmol) as catalyst, KHMDS (0.25 mmol) as base. ^c THF as a solvent, 80 °C, 20 h.

center or through aggregation of a Ni–P(OEt)₂OK complex. Racemic phosphinate **3u** was obtained in 53% yield from 4-fluorobiphenyl and cyclohexyl phenylphosphinate (**2f**) under slightly modified reaction conditions. The synthetic applications are summarized in Scheme 1. The reaction was applicable for a gram-scale synthesis. When the reaction of aryl fluoride **1b** with phosphonate **2b** was carried out on a 5.0 mmol scale, 1.1 g of arylphosphonate **3v** was isolated (84% yield, Scheme 1a). Defluorophosphonylation of *N*-methyl paroxetine, an antidepressant, afforded the corresponding arylphosphonate **3w** in 50% yield, demonstrating the potential of the present protocol for the synthesis of structurally complicated organophosphorus compounds using scaffolds of biologically functional molecules (Scheme 1b). Twofold defluorophosphonylation occurred with 1,4-difluorobenzene (**4**) under slightly modified reaction conditions, affording the corresponding phenylenediphosphonic acid ester (**5**) in 63% yield (Scheme 1c).


Scheme 1. Synthetic applications.



Scheme 2. Mechanistic experiments.



b) Reactions with a mixture of phosphites






Figure 2. (a) Calculated oxidative addition pathways. (b) Calculated reductive elimination pathways.

To gain insights into the mechanism, we performed kinetic studies by *in situ* infrared (IR) spectroscopy for the reaction of potassium salt $K[2a-H]$, which was formed *in situ* from **2a** and $KOtBu$, with 4-methoxyphenyl fluoride (**1b**) promoted by $Ni(cod)_2$ in toluene at 120 °C (Scheme 2a). The rate was found to be first order in both **1b** and $Ni(cod)_2$, suggesting that a reaction of the aryl fluoride with a monomeric Ni complex would be a turnover-limiting step. However, we failed to determine the reaction order for $K[2a-H]$ because of its low solubility.¹⁶

To explain the specific reactivity of di-*sec*-alkyl phosphonates, we conducted reactions with a mixture of phosphonates with different *O*-alkyl groups (Scheme 2b). When aryl fluoride **1b** was treated with a 1:1 mixture of diisopropyl and diethyl phosphonates (**1b** : **2b** : **2d** = 1:1:1) in the presence of $NiBr_2 \cdot diglyme$ (5 mol%) and $KOtBu$ (2 equiv), no C–P coupling product was obtained, indicating that diethyl phosphonate (**2d**) inhibited the reaction of diisopropyl phosphonate (**2b**). Thus, the irreversible formation of a catalytically inactive species from **2d** and $NiBr_2 \cdot diglyme$ is strongly suggested.

For computational investigations, we assume metal chelation by ion-bridged dimers of the potassium dialkyl phosphites $\{M[P(OR)_2O^-K^+]\}_2$ on the basis of analogy with the chelation by hydrogen-bonded phosphorus dimers $[(RO)_2POH \cdots OP(OR)_2]$ reported in the literature.^{15,17,18} Calculations were performed at the M06/SDD,6-311+G(d,p)/SMD//M06/lanl2dz,6-31G(d) level of theory using the Gaussian 16 package. To include solvent effects, the explicit coordination of a toluene molecule to each K^+ cation was considered in all the calculations. Energy profiles are given in Figure 2 for the oxidative addition of fluorobenzene (**1f**) to the Ni(0) complex $\{Ni^0[P(OiPr)_2OK]_2 \cdot (toluene)_2\}$ coordinated with two $P(OiPr)_2OK$ ligands (Figure 2a) and for reductive elimination of the arylphosphonates $[PhP(O)(OiPr)_2]$ from Ni(II) complexes $\{[P(OiPr)_2OK]_2Ni^{II}(Ph)[P(O)(OiPr)_2] \cdot (toluene)_2\}$ (Figure 2b).

As shown in Figure 2a, the C–F bond oxidative addition of fluorobenzene to the Ni center of Ni(0) intermediate **Int-1a** ($R = iPr$) to produce the corresponding pseudo-square-planar Ni(II) intermediate (**Int-2a**) occurs with Lewis acidic direct participation of one of the K^+ cations, as indicated by the increase of the $K \cdots F$ interaction (from 2.76 Å to 2.46 Å) as the reaction proceeds from **Int-1a** to transition state **TS1a-2a**. This process is 10.9 kcal·mol⁻¹ exergonic with a barrier of 14.1 kcal·mol⁻¹,


altering the P–Ni–P bite angle from 105° (for **Int-1a**) to 96° (for **Int-2a**). We reason that not only the push–pull effect of the Ni– K^+ bimetallic system but also strong electron donation by the two anionic phosphorus ligands $[P(OiPr)_2O^-]$ facilitate the C–F bond oxidative addition.¹⁹

We next attempted to identify a transition state for the direct reductive elimination of $PhP(O)(OiPr)_2$ from **Int-2a**; however, a reasonable transition-state structure was not found. This failure prompted us to investigate reductive elimination after ligand exchange at the Ni(II) center from the F^- anion to $P(OiPr)_2O^-$. The geometry-optimized Ni(II) complex (**Int-3a**) with three anionic P ligands adopts a pseudo-square-planar geometry. The P–Ni–P bite angle (96°) with the original two P ligands is unchanged upon this ligand exchange. The computational estimation of the relative energy between **Int-2a** and **Int-3a** is too challenging because of the insoluble natures of KF and $P(OiPr)_2OK$ in the reaction system and was therefore not pursued in the present study. Reductive elimination from **Int-3a** proceeds through **TS3a-4a** with an energy barrier (10.9 kcal·mol⁻¹) much lower than that for the oxidative addition process (14.1 kcal·mol⁻¹) to afford **Int-4a** with an η^2 -coordinated phosphonylbenzene via a 7.2 kcal·mol⁻¹ exergonic process.

We also conducted a computational study for the less favorable reaction with the bulkier phosphorus agent di-*tert*-butyl phosphonate ($R = tBu$). The corresponding energy diagrams are given in Figure 2 as blue lines. As in the case with diisopropyl phosphonate ($R = iPr$), the C–F bond oxidative addition (**Int-1b**–**TS1b-2b**–**Int-2b**) proceeds with Lewis acidic participation of the K^+ cation, with an energy barrier of 15.8 kcal·mol⁻¹, which is only 1.7 kcal·mol⁻¹ larger than that for the reaction with diisopropyl phosphonate ($R = iPr$) (Figure 2a). Thus, the oxidative addition step is likely not responsible for the lower reactivity of di-*tert*-butyl phosphonate compared with that of diisopropyl phosphonate. By contrast, the change of the phosphonate alkyl groups from *iPr* (**TS3a-4a**) to *tBu* (**TS3b-4b**) strongly influenced the ease of reductive elimination, increasing the energy barrier to 20.1 kcal·mol⁻¹ (Figure 2b). The energy barrier of **TS3b-4b** is 9.2 kcal·mol⁻¹ higher in energy than that for the reaction with the diisopropyl phosphonate, which is thereby deduced to be a reason for the experimentally observed inertness of $HP(O)(OtBu)_2$ (**2c**). Steric congestion in **TS3b-4b** is likely responsible for the increased energy barrier.²⁰

On the basis of the results of the experimental and theoretical studies, we propose the reaction mechanism shown in Scheme 3. The catalyst precursor $\text{NiBr}_2 \cdot \text{diglyme}$ is activated by $\text{P(OR)}_2\text{O}^- \text{K}^+$ ($\text{K}[\text{2-H}]$) generated by deprotonation of dialkyl phosphonate **2** with KOtBu to afford Ni(0) complexes $\{\text{Ni}[\text{P(OR)}_2\text{OK}]_n (n = 3, 4)\}$ that chelated with a strongly electron-donating ion-bridged dimeric phosphite ligand system $[\text{P(OR)}_2\text{O}^- \text{K}^+]_2$.²¹ Ligand exchange between one or two molecules of $\text{K}[\text{2-H}]$ and aryl fluoride **1** gives a Ni(0) complex (**A**) η^2 -coordinated with the aryl fluoride. The pronounced inhibitory effect by diethyl phosphonate **2d** is deduced to be attributable to the inertness of $\text{Ni}^0[\text{P(OEt)}_2\text{OK}]_n (n = 3, 4)$ toward ligand dissociation. Then, turnover-limiting oxidative addition of aryl fluoride **1** to the Ni center of **A** gives an aryl nickel(II) fluoride (**B**). Next, replacement of the F^- anion on the Ni center with a phosphonate anion $[\text{2a-H}]^-$ forms aryl(phosphonyl)nickel(II) complex **C**. Finally, reductive elimination of arylphosphonate **3** from **C** and re-coordination of aryl fluoride **1** regenerates Ni(0) complex **A** to complete the catalytic cycle.

Scheme 3. A proposed catalytic cycle.



CONCLUSIONS

In summary, Ni-catalyzed defluorinative phosphonylations of aryl fluorides with dialkyl phosphonates $[\text{HP(O)(OR)}_2]$ have been achieved using KOtBu as a base. The reaction required no exogenous ligands, and commercially available and bench-stable Ni(II) complexes exhibited high catalytic activities. Various aryl fluorides were successfully converted to the corresponding arylphosphonates irrespective of their electronic natures. The reaction proceeded specifically with di-*sec*-alkyl phosphonates. Experimental and computational mechanistic investigations suggested that Ni– K^+ cooperative action of a Ni(0) complex chelated with a strongly electron-donating ion-bridged dimeric phosphite ligand $[\text{P(OR)}_2\text{O}^- \text{K}^+]_2$ facilitates turnover-limiting C–F bond oxidative addition of aryl fluorides. Further reaction development with metal complexes with ion-bridged dimeric phosphonate ligands is underway in our laboratory.

EXPERIMENTAL SECTION

Typical Procedure for the Ni-catalyzed defluorophosphonylation of aryl fluorides

In a nitrogen-filled glovebox, $\text{NiBr}_2 \cdot \text{diglyme}$ (2.2 mg, 0.00625 mmol, 5 mol%) and PhMe (0.1 mL) were placed in an oven-dried 10 mL glass tube containing a magnetic stirring bar. Next, **1a** (21.5 mg, 0.125 mmol, 1 equiv), **2a** (61.6 mg, 0.25 mmol, 2 equiv), and PhMe (0.2 mL) were added to the mixture. After stirring for 5 min, KOtBu (28.1 mg, 0.25 mmol, 2 equiv) and PhMe (0.2 mL) were added, and the color of the mixture turned into orange upon stirring. The glass tube was sealed with a screw cap and was removed from the glove box. The mixture was stirred at 120 °C for 20 h. After cooling to room temperature, the dark-red reaction mixture was passed through a short plug of silica gel with a $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (9:1) eluent. Volatiles were removed by evaporation under reduced pressure. In order to remove recovered dialkylphosphonates, the crude reaction mixture was heated to 90 °C under high vacuum (140 Pa) for 12 h. After cooling to room temperature, a residue was purified by flash chromatography on silica gel with slow gradient elution ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 100:0-to-98:2) followed by preparative thin layer chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 98:2) to give **3a** as a light-brownish oil (48.4 mg, 0.12 mmol, 97% yield).

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures, details of mechanistic studies, and NMR spectra (PDF).

Cartesian coordinates of optimized geometries (XYZ).

AUTHOR INFORMATION

Corresponding Authors

* Kosuke Higashida – Institute for Chemical Reaction Design and Discovery (WPI-ICReDD), Hokkaido University, Sapporo, Hokkaido 001-0021, Japan; Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan; <https://orcid.org/0000-0002-0999-0763>; Email: higashida@icredd.hokudai.ac.jp

* Masaya Sawamura – Institute for Chemical Reaction Design and Discovery (WPI-ICReDD), Hokkaido University, Sapporo, Hokkaido 001-0021, Japan; Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan; <https://orcid.org/0000-0002-8770-2982>; Email: sawamura@sci.hokudai.ac.jp

Authors

Zhensheng You – Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan

Yusuke Masuda – Institute for Chemical Reaction Design and Discovery (WPI-ICReDD), Hokkaido University, Sapporo, Hokkaido 001-0021, Japan; Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan; <https://orcid.org/0000-0002-0386-9989>

Tomohiro Iwai – Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan; <https://orcid.org/0000-0002-5878-6339>

Funding Sources

This research was supported by JSPS KAKENHI Grant numbers JP21H04680 in Grant-in-Aid for Scientific Research (A) (MS), JP20K15268 in Grant-in-Aid for Early-Career Scientists (KH), JP21K14626 in Grant-in-Aid for Early-Career Scientists (YM),

JP22H05331 in Grant-in-Aid for Transformative Research Area (A) Digitalization-driven Transformative Organic Synthesis (Digi-TOS) (YM), JP20H04793 in Grant-in-Aid for Scientific Research on Innovative Areas, "Hybrid Catalysis" (TI), JP22J11710 in Grant-in-Aid for JSPS Research Fellow (ZY) and by the Naito Foundation (TI).

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

Z.Y. thanks the Program for Leading Graduate Schools (Hokkaido University "Ambitious Leader's Program") and the MEXT Doctoral program for Data-Related Innovation Expert Hokkaido University (D-DRIVE-HU).

REFERENCES

- (1) (a) Muller, K.; Faeh, C.; Diederich, F. Fluorine in pharmaceuticals: looking beyond intuition. *Science* **2007**, *317*, 1881–1886. (b) Wang, J.; Sánchez-Roselló, M.; Aceña, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. Fluorine in Pharmaceutical Industry: Fluorine-Containing Drugs Introduced to the Market in the Last Decade (2001–2011). *Chem. Rev.* **2014**, *114*, 2432–2506. (c) Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. Next Generation of Fluorine-Containing Pharmaceuticals, Compounds Currently in Phase II–III Clinical Trials of Major Pharmaceutical Companies: New Structural Trends and Therapeutic Areas. *Chem. Rev.* **2016**, *116*, 422–518. (d) Inoue, M.; Sumii, Y.; Shibata, N. Contribution of Organofluorine Compounds to Pharmaceuticals. *ACS Omega* **2020**, *5*, 10633–10640. (e) Sap, J. B. I.; Meyer, C. F.; Straathof, N. J. W.; Iwumene, N.; am Ende, C. W.; Trabanco, A. A.; Gouverneur, V. Late-stage difluoromethylation: concepts, developments and perspective. *Chem. Soc. Rev.* **2021**, *50*, 8214–8247.
- (2) (a) Guittard, F.; de Givenchy, E. T.; Geribaldi, S.; Cambon, A. Highly fluorinated thermotropic liquid crystals: an update. *J. Fluorine Chem.* **1999**, *100*, 85–96. (b) Babudri, F.; Farinola, G. M.; Naso, F.; Ragni, R. Fluorinated organic materials for electronic and optoelectronic applications: the role of the fluorine atom. *Chem. Commun.* **2007**, 1003–1022. (c) Berger, R.; Resnati, G.; Metrangolo, P.; Weber, E.; Hulliger, J. Organic fluorine compounds: a great opportunity for enhanced materials properties. *Chem. Soc. Rev.* **2011**, *40*, 3496–3508.
- (3) For selected reviews, see (a) Amii, H.; Uneyama, K. C–F Bond Activation in Organic Synthesis. *Chem. Rev.* **2009**, *109*, 2119–2183. (b) Lei, A.; Liu, W.; Liu, C.; Chen, M. Arylation of unactivated arenes. *Dalton Trans.* **2010**, *39*, 10352–10361. (c) Braun, T.; Wehmeier, F. C–F Bond Activation of Highly Fluorinated Molecules at Rhodium: From Model Reactions to Catalysis. *Eur. J. Inorg. Chem.* **2011**, *2011*, 613–625. (d) Rosen, B. M.; Quasdorf, K. W.; Wilson, D. A.; Zhang, N.; Resmerita, A.-M.; Garg, N. K.; Percec, V. Nickel-Catalyzed Cross-Couplings Involving Carbon–Oxygen Bonds. *Chem. Rev.* **2011**, *111*, 1346–1416. (e) Tasker, S. Z.; Standley, E. A.; Jamison, T. F. Recent advances in homogeneous nickel catalysis. *Nature* **2014**, *509*, 299–309. (f) Kakiuchi, F.; Kochi, T.; Murai, S. Chelation-Assisted Regioselective Catalytic Functionalization of C–H, C–O, C–N and C–F Bonds. *Synlett* **2014**, *25*, 2390–2414. (g) Ahrens, T.; Kohlmann, J.; Ahrens, M.; Braun, T. Functionalization of Fluorinated Molecules by Transition-Metal-Mediated C–F Bond Activation To Access Fluorinated Building Blocks. *Chem. Rev.* **2015**, *115*, 931–972. (h) Yan, G.; Qiu, K.; Guo, M. Recent advance in the C–F bond functionalization of trifluoromethyl-containing compounds. *Org. Chem. Front.* **2021**, *8*, 3915–3942. (i) Wang, Z.; Sun, Y.; Shen, L.-Y.; Yang, W.-C.; Meng, F.; Li, P. Photochemical and electrochemical strategies in C–F bond activation and functionalization. *Org. Chem. Front.* **2022**, *9*, 853–873.
- (4) (a) Coote, S. J.; Dawson, G. J.; Frost, C. G.; Williams, J. M. J. The Preparation of Functionalised Aryl Phosphines from Aryl Fluorides by Nucleophilic Aromatic Substitution with Potassium Diphenylphosphide. *Synlett* **1993**, *1993*, 509–510. (b) Hingst, M.; Tepper, M.; Stelzer, O. Nucleophilic Phosphorylation of Fluoroaromatic

Compounds with Carboxyl, Carboxymethyl, and Aminomethyl Functionalities – an Efficient Synthetic Route to Amphiphilic Arylphosphanes. *Eur. J. Inorg. Chem.* **1998**, 73–82. (c) Seipel, K. R.; Platt, Z. H.; Nguyen, M.; Holland, A. W. Microwave-Assisted Synthesis of Phenylene-Bridged Aminophosphine Ligands: Acceleration of N-Arylation and Aryl Fluoride Phosphorylation Reactions. *J. Org. Chem.* **2008**, *73*, 4291–4294.

(5) (a) Horner, L.; Beck, P.; Hoffmann, H. Phosphororganische Verbindungen, XIX. Reduktion von Phosphorverbindungen mit Alkalimetallen. *Chem. Ber.* **1959**, 2088–2094. (b) Nandi, P.; Dye, J. L.; Bentley, P.; Jackson, J. E. Preparation of Diphenyl Phosphide and Substituted Phosphines using Alkali Metal in Silica Gel (M–SG). *Org. Lett.* **2009**, *11*, 1689–1692. (c) Ye, J.; Zhang, J.-Q.; Saga, Y.; Onozawa, S.; Kobayashi, S.; Sato, K.; Fukaya, N.; Han, L.-B. Ready Approach to Organophosphines from ArCl via Selective Cleavage of C–P Bonds by Sodium. *Organometallics* **2020**, *39*, 2682–2694.

(6) (a) Goryunov, L. I.; Grobe, J.; Le Van, D.; Shteingarts, V. D.; Mews, R.; Lork, E.; Würthwein, E.-U. Di- and Trifluorobenzenes in Reactions with Me₂EM (E = P, N; M = SiMe₃, SnMe₃, Li) Reagents: Evidence for a Concerted Mechanism of Aromatic Nucleophilic Substitution. *Eur. J. Org. Chem.* **2010**, 1111–1123. (b) Zhivetyeva, S. I.; Goryunova, L. I.; Bagryanskaya, I. Y.; Grobe, J.; Shteingarts, V. D.; Würthwein, E.-U. Phosphinodifluorination of polyfluorobenzenes by silylphosphines Ph(R)PSiMe₃ (R = Me, Ph): Further experimental and computational evidences for the concerted A_ND_N mechanism of aromatic nucleophilic substitution. *J. Fluor. Chem.* **2014**, *164*, 58–69.

(7) (a) Verbrugghen, T.; Vandurm, P.; Pouyez, J.; Maes, L.; Wouters, J.; Van Calenbergh, S. Alpha-Heteroatom Derivatized Analogues of 3-(Acetylhydroxyamino)propyl Phosphonic Acid (FR900098) as Antimalarials. *J. Med. Chem.* **2013**, *56*, 376–380. (b) Kramer, G. J.; Mohd, A.; Schwager, S. L. U.; Masuyer, G.; Acharya, K. R.; Sturrock, E. D.; Bachmann, B. O. Interkingdom Pharmacology of Angiotensin-I Converting Enzyme Inhibitor Phosphonates Produced by Actinomycetes. *ACS Med. Chem. Lett.* **2014**, *5*, 346–351. (c) Wiemer, A. J.; Wiemer, D. F. Prodrugs of Phosphonates and Phosphates: Crossing the Membrane Barrier. *Top. Curr. Chem.* **2015**, *360*, 115–160. (d) Brücher, K.; Gräwert, T.; Konzuch, S.; Held, J.; Lienau, C.; Behrendt, C.; Illarionov, B.; Maes, L.; Bacher, A.; Wittlin, S.; Mordmüller, B.; Fischer, M.; Kurz, T. Prodrugs of Reverse Fosmidomycin Analogues. *J. Med. Chem.* **2015**, *58*, 2025–2035. (e) Desai, J.; Wang, Y.; Wang, K.; Malwal, S. R.; Oldfield, E. Isoprenoid Biosynthesis Inhibitors Targeting Bacterial Cell Growth. *ChemMedChem* **2016**, *11*, 2205–2215. (f) Falagas, M. E.; Vouloumanou, E. K.; Samonis, G.; Vardakas, K. Z. Fosfomycin. *Clin. Microbiol. Rev.* **2016**, *29*, 321–347.

(8) (a) Paurová, M.; Havlíčková, J.; Pospíšilová, A.; Vetrík, M.; Císařová, I.; Stephan, H.; Pietzsch, H.-J.; Hrubý, M.; Hermann, P.; Kotek, J. Bifunctional Cyclam-Based Ligands with Phosphorus Acid Pendant Moieties for Radiocopper Separation: Thermodynamic and Kinetic Studies. *Chem. Eur. J.* **2015**, *21*, 4671–4687. (b) Ševčík, R.; Vaněk, J.; Michalíková, R.; Lubal, P.; Hermann, P.; Santos, I. C.; Santos, I.; Campello, M. P. C. Formation and decomplexation kinetics of copper(II) complexes with cyclen derivatives having mixed carboxylate and phosphonate pendant arms. *Dalton Trans.* **2016**, *45*, 12723–12733. (c) Kostelnik, T. I.; Scheiber, H.; Cappai, R.; Choudhary, N.; Lindheimer, F.; Jaraquemada-Peláez, M.; Orvig, C. Phosphonate Chelators for Medicinal Metal Ions. *Inorg. Chem.* **2021**, *60*, 5343–5361.

(9) (a) Pujari, S. P.; Scheres, L.; Marcellis, A. T. M.; Zuilhof, H. Covalent Surface Modification of Oxide Surfaces. *Angew. Chem., Int. Ed.* **2014**, *53*, 6322–6356. (b) Boissezon, R.; Muller, J.; Beaugeard, V.; Monge, S.; Robin, J.-J. Organophosphonates as anchoring agents onto metal oxide-based materials: synthesis and applications. *RSC Adv.* **2014**, *4*, 35690–35707.

(10) (a) Morita, C. T.; Jin, C.; Sarikonda, G.; Wang, H. Nonpeptide antigens, presentation mechanisms, and immunological memory of human Vγ2Vδ2 T cells: discriminating friend from foe through the recognition of prenyl pyrophosphate antigens. *Immunol. Rev.* **2007**, *215*, 59–76. (b) Zhang, Y.; Cao, R.; Yin, F.; Lin, F.-Y.; Wang, H.; Krysiak, K.; No, J.-H.; Mulkamala, D.; Houlihan, K.; Li, J.; Morita, C. T.; Oldfield, E. Lipophilic Pyridinium Bisphosphonates: Potent γδ T Cell Stimulators. *Angew. Chem. Int. Ed.* **2010**, *49*, 1136–1138.

- (11) Dou, Q.; Lang, Y.; Zeng, H.; Li, C.-J. Photoinduced transition-metal and external photosensitizer free phosphonation of unactivated C(sp²)-F bond via SET process under mild conditions. *Fundam. Res.* **2021**, *1*, 742–746.
- (12) You, Z.; Higashida, K.; Iwai, T.; Sawamura, M. Phosphinylation of Non-activated Aryl Fluorides through Nucleophilic Aromatic Substitution at the Boundary of Concerted and Stepwise Mechanisms. *Angew. Chem. Int. Ed.* **2021**, *60*, 5778–5782.
- (13) For selected examples of Ni-catalyzed C(sp²)-F bond functionalizations of monofluoroarenes, see (a) Böhm, V. P. W.; Gstöttmayr, C. W. K.; Weskamp, T.; Herrmann, W. A. Catalytic C–C Bond Formation through Selective Activation of C–F Bonds. *Angew. Chem. Int. Ed.* **2001**, *40*, 3387–3389. (b) Yoshikai, N.; Mashima, H.; Nakamura, E. Nickel-Catalyzed Cross-Coupling Reaction of Aryl Fluorides and Chlorides with Grignard Reagents under Nickel/Magnesium Bimetallic Cooperation. *J. Am. Chem. Soc.* **2005**, *127*, 17978–17979. (c) Wang, J.-R.; Manabe, K. High Ortho Preference in Ni-Catalyzed Cross-Coupling of Halophenols with Alkyl Grignard Reagents. *Org. Lett.* **2009**, *11*, 741–744. (d) Tobisu, M.; Xu, T.; Shimasaki, T.; Chatani, N. Nickel-Catalyzed Suzuki–Miyaura Reaction of Aryl Fluorides. *J. Am. Chem. Soc.* **2011**, *133*, 19505–19511. (e) Nakamura, Y.; Yoshikai, N.; Iliés, L.; Nakamura, E. Nickel-Catalyzed Monosubstitution of Polyfluoroarenes with Organozinc Reagents Using Alkoxydiphosphine Ligand. *Org. Lett.* **2012**, *14*, 3316–3319. (f) Zhu, F.; Wang, Z.-X. Nickel-Catalyzed Coupling of Fluoroarenes and Amines. *Adv. Synth. Catal.* **2013**, *355*, 3694–3702. (g) Zhu, F.; Wang, Z.-X. Nickel-Catalyzed Cross-Coupling of Aryl Fluorides and Organozinc Reagents. *J. Org. Chem.* **2014**, *79*, 4285–4292. (h) Ogawa, H.; Yang, Z.-K.; Minami, H.; Kojima, K.; Saito, T.; Wang, C.; Uchiyama, M. Revisitation of Organoaluminum Reagents Affords a Versatile Protocol for C–X (X = N, O, F) Bond-Cleavage Cross-Coupling: A Systematic Study. *ACS Catal.* **2017**, *7*, 3988–3994. (i) Harada, T.; Ueda, Y.; Iwai, T.; Sawamura, M. Nickel-catalyzed amination of aryl fluorides with primary amines. *Chem. Commun.* **2018**, *54*, 1718–1721 and references cited therein.
- (14) For nickel-catalyzed phosphorylation of aryl chlorides, bromides, iodides, and pseudo halides, see (a) Balthazor, T. M.; Grabiak, R. C. Nickel-catalyzed Arbuzov reaction: mechanistic observations. *J. Org. Chem.* **1980**, *45*, 5425–5426. (b) Zhao, Y.-L.; Wu, G.-J.; Li, Y.; Gao, L.-X.; Han, F.-S. [NiCl₂(dppp)]-Catalyzed Cross-Coupling of Aryl Halides with Dialkyl Phosphite, Diphenylphosphine Oxide, and Diphenylphosphine. *Chem. Eur. J.* **2012**, *18*, 9622–9627. (c) Shen, C.; Yang, G.; Zhang, W. Nickel-catalyzed C–P coupling of aryl mesylates and tosylates with H(O)PR¹R². *Org. Biomol. Chem.* **2012**, *10*, 3500–3505. (d) Yang, J.; Xiao, J.; Chen, T.; Han, L.-B. Nickel-Catalyzed Phosphorylation of Phenol Derivatives via C–O/P–H Cross-Coupling. *J. Org. Chem.* **2016**, *81*, 3911–3916. (e) Li, C.-J. Nickel-Catalyzed Phosphorylation of Tosylates. *Russ. J. Gen. Chem.* **2020**, *90*, 725–730. (f) Zhu, D.-L.; Jiang, S.; Wu, Q.; Wang, H.; Chai, L.-L.; Li, H.-Y.; Li, H.-X. Visible-Light-Induced Nickel-Catalyzed P(O)–C(sp²) Coupling Using Thioxanthen-9-one as a Photoredox Catalysis. *Org. Lett.* **2021**, *23*, 160–165.
- (15) For selected reviews of transition-metal catalysts with phosphite ligands, see (a) Ackermann, L. Catalytic Arylations with Challenging Substrates: From Air-Stable HASPO Preligands to Indole Syntheses and C–H-Bond Functionalizations. *Synlett* **2007**, 507–526. (b) Ackermann, L. Air-Stable Bifunctional HASPO Preligands for Metal-Catalyzed Cross-Couplings and Direct C–H Bond Arylations. *Isr. J. Chem.* **2010**, *50*, 652–663. (c) Reek, J. N. H.; Bruin, B.; Pullen, S.; Mooibroek, T. J.; Kluwer, A. M.; Caumes, X. Transition Metal Catalysis Controlled by Hydrogen Bonding in the Second Coordination Sphere. *Chem. Rev.* **2022**, *122*, 12308–12369.
- (16) See the Supporting Information for details.
- (17) (a) Werner, H.; Khac, T. N. “Supersandwich”-Type Trinuclear Complexes with a Spiro Center. *Angew. Chem. Int. Ed.* **1977**, *16*, 324–325. (b) Werner, H.; Khac, T. N. Two-, four-, five- and six-coordinated metal complexes containing the organometallic chelate C₅H₅M[P(OMe)₂O]₂, M = Ni and Pd. *Inorg. Chim. Acta* **1978**, *30*, L347–L348. (c) Werner, H.; Khac, T. N. Metallabis(phosphonate) als Chelatliganden. I. Synthese einkerniger Nickelbis(phosphonat)-Komplexe mit OHO-Wasserstoffbrücke und entsprechender Alkali-metall-, Ammonium- und Thallium-Verbindungen. *Z. Anorg. Allg. Chem.* **1981**, *475*, 241–250. (d) Paine, R. T.; Duesler, E. N.; Moody, D. C. Synthesis and structure of a supersandwich pentanuclear uranium(IV) complex, U{Ni[P(O)(OMe)₂]₂(η -5-C₅H₅)₄}. *Organometallics* **1982**, *1*, 1097–1098. (e) Ackermann, L.; Born, R.; Spatz, J. H.; Meyer, D. Efficient Aryl–(Hetero)Aryl Coupling by Activation of C–Cl and C–F Bonds Using Nickel Complexes of Air-Stable Phosphine Oxides. *Angew. Chem. Int. Ed.* **2005**, *44*, 7216–7219.
- (18) A ³¹P NMR spectroscopy study of the reaction between Ni(cod)₂ and HP(O)(O*i*Pr)₂/KHMDS (1:1 to 1:5) in toluene-*d*₈ showed dose-dependent dissociation of the cod ligand and the formation of a mixture of Ni–P coordination compounds. See Supporting Information for details.
- (19) An S_NAr pathway from a π -Ni(0) complex was also considered, but we were unable to locate a reasonable transition state. S_NAr pathways were proposed for the Ni-catalyzed silylation reactions and the base-mediated defluorosilylation. See: (a) Cui, B.; Jia, S.; Tokunaga, E.; Shibata, N. Defluorosilylation of fluoroarenes and fluoroalkanes. *Nat. Commun.* **2018**, *9*, 4393. (b) Zarate, C.; Nakajima, M.; Martin, R. A Mild and Ligand-Free Ni-Catalyzed Silylation via C–OMe Cleavage. *J. Am. Chem. Soc.* **2017**, *139*, 1191–1197. (c) Liu, X.-W.; Zarate, C.; Martin, R. Base-Mediated Defluorosilylation of C(sp²)-F and C(sp³)-F Bonds. *Angew. Chem. Int. Ed.* **2019**, *58*, 2064–2068.
- (20) Further calculation results are discussed in Supporting Information.
- (21) DFT calculations suggested that the Ni(0) species could be generated through reductive elimination of trialkyl phosphate from a *t*-butoxonickel(II) phosphite complex. See Supporting Information for details (Figure S21).