HOKKAIDO UNIVERSITY

Title	Clinical features of anti-mitochondrial M2 antibody-positive myositis : case series of 17 patients
Author(s)	Nagai, Azusa; Nagai, Toshiyuki; Yaguchi, Hiroaki et al.
Citation	Journal of the neurological sciences, 442, 120391 https://doi.org/10.1016/j.jns.2022.120391
Issue Date	2022-11-15
Doc URL	https://hdl.handle.net/2115/90686
Rights	© 2022, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	Fig.1 A B C D E F.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Correlation of %VC and mRS score

$R = 0.67$, $p = 0.0034$

mRS score

0
1
2
3
4
5

40
60
80
100

%VC (%)

Correlation of BMI and %VC

$R = 0.68$, $p = 0.0025$

%VC (%)

100

75

50

25

16

20

24

28

BMI

Correlation of BMI and mRS score


$R = 0.56$, $p = 0.019$

mRS score


0
1
2
3
4
5


16
20
24
28

BMI


Correlation of LVEF and mRS score
R = −0.037, p = 0.89
mRS score
0
1
2
3
4
5
30
40
50
60
70
LVEF (%)

Correlation of disease duration before diagnosis and mRS score




Correlation of AMA-M2 index and mRS score
R = −0.026, p = 0.92
mRS score
0
1
2
3
4
5
0
100
200
300
400
AMA-M2 (index)