



HOKKAIDO UNIVERSITY

Title	The association of thyroglobulin single nucleotide polymorphism with miniature dachshunds-specific inflammatory colorectal polyps and its involvement in interleukin-6 amplifier induced chronic inflammation [an abstract of dissertation and a summary of dissertation review]
Author(s)	Teoh, Yong Bin; テオ, ヤン ビン
Degree Grantor	北海道大学
Degree Name	博士(獣医学)
Dissertation Number	甲第15655号
Issue Date	2023-09-25
Doc URL	https://hdl.handle.net/2115/90962
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Type	doctoral thesis
File Information	Teoh_Yong_Bin_abstract.pdf, 論文内容の要旨



学位論文内容の要旨
Abstract of the dissertation

博士の専攻分野の名称：博士（獣医学）

氏名：テオ ヤン ビン

Name : TEOH YONG BIN

学位論文題名
The title of the doctoral dissertation

The Association of Thyroglobulin Single Nucleotide Polymorphism with Miniature Dachshunds-Specific Inflammatory Colorectal Polyps and its Involvement in Interleukin-6 Amplifier Induced Chronic Inflammation

(ミニチュアダックスフンドに特異的な炎症性結直腸ポリープに関連するサイログロブリンの一塩基多型とインターロイキン6増幅回路に誘導される慢性炎症への関与)

Inflammatory colorectal polyp (ICRP) of Miniature Dachshund (MD) is a novel inflammatory bowel disease (IBD) characterized by granulomatous inflammation consisting neutrophil infiltration and goblet cell hyperplasia in the colorectal region with an unknown pathogenesis. Whole-exome sequencing revealed that *TG* c.4567C>T and *FBN1* c.1205C>T (p.P402L) are MD specific single nucleotide polymorphisms (SNP), but only thyroglobulin (*TG*) was found to be significantly different between disease-control groups in our cohort ($P = 0.0221$). Mechanistic analysis showed that *TG* increased the activation of the IL-6 amplifier, which is an enhanced-*NF-κB* activation in nonimmune cells in the presence of a simultaneous activation of *NF-κB* and *STAT3* and is a fundamental machinery of various inflammatory diseases in patients and animal models. Indeed, *TG* treatment increased *NF-κB*-mediated IL-6 expression, while *TG* deficient reduced it at least in vitro. Moreover, we showed that *NF-κB* and *STAT3* were more activated in intestine samples with risk allele comparing to those with non-risk allele. Although systemic expression of *TG* was unable to be detected, local expression of *TG* was increased in normal intestine samples with risk allele. Thus, these results suggest that regional increase of *TG* expression in intestine induces chronic inflammation via the activation of the IL-6 amplifier in MDs, which may also be important in human IBD patients with concurrent thyroidal disease.