



Title	Synergistic effect of standardized extract of <i>Asparagus officinalis</i> stem and heat shock on progesterone synthesis with lipid droplets and mitochondrial function in bovine granulosa cells
Author(s)	Ho, Khoi Thieu; Balboula, Ahmed Zaky; Homma, Kohei et al.
Citation	Journal of steroid biochemistry and molecular biology, 225, 106181 https://doi.org/10.1016/j.jsbmb.2022.106181
Issue Date	2023-01
Doc URL	https://hdl.handle.net/2115/91027
Rights	© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	J Steroid Biochem Mol Biol._225_106181.pdf



Synergistic effect of standardized extract of *Asparagus officinalis* stem and heat shock on progesterone synthesis with lipid droplets and mitochondrial function in bovine granulosa cells

Khoi Thieu Ho ^{1,4}, Ahmed Zaky Balboula², Kohei Homma³, Jun Takanari³, Hanako Bai¹,
Manabu Kawahara¹, Khang Thi Kim Nguyen ⁴, Masashi Takahashi⁵

¹Graduate School of Agriculture, Hokkaido University, Sapporo, Hokkaido, 060-8589, Japan

²University of Missouri Columbia, MO 65211, USA

³AMINO UP Co. Ltd., Sapporo, Hokkaido, 004-0839, Japan

⁴College of Agriculture, Can Tho University, Can Tho city, Viet Nam

⁵Research Faculty of Agriculture, Hokkaido University, Sapporo, Hokkaido, 060-8589, Japan

***Corresponding author:**

Masashi Takahashi, Graduate School of Agriculture, Hokkaido University, Sapporo,
Hokkaido, 060-8589, Japan

email address: mmasashi@anim.agr.hokudai.ac.jp

Abstract

Progesterone (P4) is a well-known steroid hormone that plays a key role in oocyte growth and the maintenance of pregnancy in mammals, including cattle. Heat stress (HS) has an adverse effect on P4 synthesis through an imbalance in the cellular redox status. We have recently revealed that a standardized extract of *Asparagus officinalis* stem (EAS) increases P4 through non-HS induction of heat shock protein 70 (HSP70) and a synergistic increase of HSP70 by enhancing the intracellular redox balance, which was adversely affected by HS in bovine granulosa cells (GCs). Bovine GCs collected from bovine ovarian follicles were cultured at 38.5°C and 41°C for 12 h with or without 5 mg/mL EAS. After treatment, cells and culture supernatant were collected for the analysis. Enzyme-linked immunosorbent assay (ELISA) was performed to detect in P4 level. Quantitative reverse-transcription polymerase chain reaction (RT-qPCR) was used to detect expression of steroidogenesis related genes. Fluorescence staining was used to detect mitochondrial activity and lipid droplet. P4 level was increased by EAS treatment in association with increase in steroidogenic acute regulatory protein (STAR), 3 β -hydroxysteroid dehydrogenase (3 β -HSD), mitochondrial membrane activity and lipid droplet both under non-HS and HS conditions. Notably, synergistic effect of EAS with HS co-treatment was observed to show a greater increase in P4 synthesis when comparison with EAS treatment under non-HS condition. Furthermore, inhibition of HSP70 significantly reduced EAS-induced P4 synthesis, mitochondrial activity and synthesis of lipid droplets. These results suggest that P4 synthesis by EAS is mediated

41 by the steroidogenesis pathway via HSP70-regulated activation of *STAR* and *3 β -HSD*,
42 together with improved mitochondrial activity and lipid metabolism in bovine GCs. Moreover,
43 effect of EAS has a synergistic effect of with HSP70-regulated steroidogenesis pathway.

44

45 **Keywords:** Bovine granulosa cells, EAS, HSP70, Progesterone, synergistic effect

46

1. Introduction

Progesterone (P4) is an important steroid hormone that plays a pivotal role in the establishment of uterine receptivity, oocyte maturation, and the maintenance of pregnancy [1,2]. In mammals, granulosa cells (GCs) play a fundamental role in the maturation and acquisition of developmental competence in bovine oocytes [3]. GCs differentiate into luteal cells that are responsible for P4 production [4]. Therefore, GCs are good models for inducing steroidogenesis under heat stress (HS) conditions to evaluate the importance of maintaining reproductive functions.

P4 synthesis in GCs is regulated by cholesterol transport into the mitochondria by steroidogenesis enzymes, such as STAR, cytochrome P450 family 11 subfamily a member 1 (CYP11A1), and 3 β -HSD [5,6]. Importantly, mitochondria play a central role in steroidogenesis and are abundant in GCs [6,7]. Moreover, P4 synthesis in bovine GCs is associated with lipid metabolism [8]. In addition, cholesterol from lipid droplets is one of the available sources for P4 biosynthesis [7]. One of the main transcription factors that control lipid metabolism by regulating enzymes required for cholesterol, fatty acid, triacylglycerol, and phospholipid synthesis is *SREBP-1* [9]. Thus, P4 synthesis in bovine GCs is regulated by steroidogenesis, mitochondrial function, and lipid metabolism. However, a critical issue that affects dairy farming is the reduction of reproductive performance in high-yielding dairy cows [10]. The reproductive performance of dairy cows is decreased by increased body

temperature [11,12] which decreases ovarian function, follicular growth, and corpus luteum function in summer.

High ambient temperature produces reactive oxygen species (ROS) with reduced P4 production in bovine GCs [13]. In addition, a negative effect of ROS on P4 levels has been reported in human follicular fluid [14]. Follicular fluid contains soluble factors secreted from GCs that maintain the intrafollicular redox status [15,16]. ROS induces redox status imbalance, which leads to cellular damage, alteration of cellular function, and physiological processes [17,18]. Thus, HS has an adverse impact on P4 synthesis owing to HS-induced toxic factors that disturb the redox imbalance in GCs.

Under stress conditions, cellular conditions are restored by the induction of heat shock proteins (HSPs), which act as molecular chaperones by folding and unfolding proteins while maintaining homeostasis of the cells [19]. Therefore, strengthening chaperone systems by inducing HSPs is one of the keys to regulating their functions in various biological pathways in ovarian functions, including P4 synthesis. However, as described above, HS is accompanied by the induction of ROS with harmful effects.

Recently, a standardized extract of *Asparagus officinalis* stem (EAS) was shown to induce HS-independent HSP70 in human promyelocytic leukemia cells [20]. In addition, we have revealed that EAS has a beneficial effect in inducing HSP70 under non-HS conditions and maintaining redox balance through ROS reduction and glutathione (GSH) synthesis in bovine GCs [21]. Moreover, EAS showed a synergistic increase in HSP70 expression and an

improvement in the intracellular redox status of bovine GCs, which was adversely affected by HS [21]. The abundance regulation of EAS-induced HSP70 which affects redox balance under both HS and non-HS conditions may compromise P4 synthesis in bovine GCs. In addition, the effects of EAS on P4 biosynthesis under HS and non-HS conditions in bovine GCs cells have not yet been investigated. In the present study, we investigated the effects of EAS on P4 levels, steroidogenesis genes, mitochondrial function, and lipid metabolism under non-HS and HS conditions.

2. Materials and Methods

Extract of *Asparagus officinalis* stem (EAS), which was extracted from asparagus (*A. officinalis* L.), was produced following a previously described method [22] and commercially available as a supplement (ETAS®).

2.1 Collection and culture of bovine GCs

Bovine ovaries were obtained from a local abattoir and were transported within 4 hours to the laboratory at 20°C. The ovaries were washed several times with a sterile saline solution. Oocytes were collected from follicles (2-8 mm in diameter) without signs of atresia, using a disposable 18-gauge needle attached to a 10-mL syringe. After cumulus-oocyte complexes were picked up, the remaining follicular fluid with floating GCs cells was cultured in

Dulbecco's modified Eagle's medium (high glucose) (DMEM) (Wako, Osaka, Japan) containing 5% fetal bovine serum (FBS), 0.06 g/L penicillin G potassium (Nacalai Tesque, Kyoto, Japan), and 0.1 g/L streptomycin sulfate (Nacalai Tesque, Kyoto, Japan) at 38.5°C under 5% CO₂ in air.

After overnight culture, the theca cell cluster layer was gently removed from the bottom of the cell culture dishes. The cells remaining at the bottom of the dish were bovine GCs, which were then washed with calcium- and magnesium-free phosphate-buffered saline (PBS) (–), and cultured in 5% FBS in DMEM at 38.5°C under 5% CO₂ in air. When the cells reached confluence, they were washed with PBS (–) and dissociated from the substratum with PBS (–) containing 0.05% trypsin and 0.53 mM Ethylenediaminetetraacetic acid (EDTA) for 2 min at 38.5°C in a CO₂ incubator. After the addition of 5% FBS in DMEM to inactivate trypsin activity, the separated cell suspension was centrifuged at 1,200 × *g* for 3 min. To culture these bovine GCs, 1 × 10⁵ cells/mL viable cells were seeded in each well of a 4-well dish (Thermo Fisher Scientific) or 8-well slide chamber (Thermo Fisher Scientific) and cultured at 38.5°C under 5% CO₂ in air. After the cells reached 70% confluence, the medium was replaced with 0.9 mL of DMEM in 5% FBS together with 0.1 mL of EAS solution in PBS (–), and 0.1 mL of PBS (–) was added to the control group. The cells were then cultured at 38.5°C or 41°C under 5% CO₂ atmosphere.

2.2 Experimental design

To investigate the effect of EAS on P4 production, the cells were cultured at 38.5°C for 12 h with 0.5, 1, 5, and 10 mg/mL of EAS. In addition to evaluating the effects of EAS on P4 synthesis under non-HS and HS conditions, cells were cultured at 38.5°C and 41°C for 12 h with or without 5 mg/mL EAS. To confirm the role of EAS-induced HSP70 in P4 synthesis, EAS-treated cells were exposed to 10 μ M pifithrin- μ (2-phenylethynesulfonamide:PES) (StressMarq Biosciences Inc., Victoria, Canada) under non-HS and HS conditions. After culture, cells and culture medium were used for mRNA expression analysis, P4 measurement, and detection of lipid droplets and mitochondria.

2.3 P4 measurement

The concentrations of P4 in culture supernatants of bovine CGs were measured by enzyme-linked immunosorbent assay (ELISA) using P4 measurement kits (ADI- 900– 011, Enzo Life Sciences, USA). The intra- and inter-assay coefficients of variation for P4 were less than 10%. The aforementioned experiments were performed in duplicate.

2.4 RNA extraction and quantitative reverse-transcription polymerase chain reaction (RT-qPCR)

Total RNA was extracted from the cells using ISOGEN II (Nippon Gene, Toyama, Japan) according to the manufacturer's instructions. The RNA concentration was measured using spectrophotometry (NanoDrop ND-2000; Thermo Fisher Scientific). All RNA samples were

stored at -80°C until use. Total RNA was reverse transcribed using ReverTra Ace qPCR RT Master Mix with gDNA remover (TOYOBO Life Science, Osaka, Japan) with a thermal cycler (Astec GeneAtlas Type G Thermal Cycler; ASTEC, Fukuoka, Japan). All cDNA samples were stored at -30°C until use for Quantitative PCR. Specific primers (Table S1) were designed using Primer-BLAST (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>). qPCR was performed using THUNDERBIRD SYBR qPCR Mix (Toyobo Life Science) at a final primer concentration of $0.5\ \mu\text{M}$. To quantify mRNA expression levels, PCR was performed under the following conditions with the LightCycler Nano (Roche Diagnostics, Basel, Switzerland): 1 cycle at 95°C for 30 s (denaturation), 45 cycles at 95°C for 10 s (denaturation), 55°C for 15 s (primer annealing), and 72°C for 30 s (extension). Relative mRNA abundance was calculated using the $\Delta\Delta\text{Ct}$ method with H2AFZ as a reference gene.

2.5 Immunodetection of HSP70

HSP70 immunofluorescence analysis was performed as previously described [21]. After GCs were cultured in 8-well slide chambers, they were fixed with 4% paraformaldehyde diluted with PBS (–), permeabilized with 0.2% Triton X-100 in PBS (–), blocked with 2% (w/v) BSA (Sigma-Aldrich) in PBS (–), and incubated at 4°C overnight with rabbit anti-human HSP70 polyclonal primary antibody (SPC-103; StressMarq Biosciences Inc., Victoria, Canada). After washing three times with PBS, the cells were incubated for 30 min with the secondary antibody Alexa Fluor 488 donkey anti-rabbit IgG (A21206; Thermo Fisher

Scientific). The 8-well slide chambers were then mounted with mounting solution (Fluoro-KEEPER Anti fade Reagent, Non-Hardening Type with DAPI, Nacalai Tesque) and observed under a fluorescence microscope (Leica Camera AG, Wetzlar, Germany).

2.6 Detection of Lipid droplet and mitochondrial activity

For the detection of lipid droplets, cells were washed with PBS (–) after culture and fixed with 4% PFA in PBS (–) for 10 min at room temperature. After washing the cells three times with PBS (–) for 5 min, they were treated with 1 µg/mL Nile Red (Wako, Osaka, Japan) for 30 min at 38.5°C in a 5% CO₂ atmosphere.

To detect mitochondrial activity, cells were washed with PBS (–) after culture and treated with 200 nM MitoTracker™ Green FM (Thermo Fischer Scientific) for 30 min at 38.5°C in a 5% CO₂ atmosphere.

Cells from both detections were then washed with PBS (–), and 10 µL of mounting solution (Nacalai Tesque) was added to the cells and covered with a cover glass. Fluorescence images were obtained using a fluorescence microscope (Leica, Wetzlar, Germany).

Fluorescence intensity was quantified using ImageJ software v1.52A (National Institutes of Health; <http://imagej.nih.gov/ij/>). Corrected total cell fluorescence (CTCF) was carried out following a previously described formula [23]:

1. CTCF = integrated density (total area of selected cells × mean fluorescence of background readings).

2. $\text{CTCF per cell} = \text{CTCF} / \text{Ncells}$

where “integrated density” is the integrated intensity of pixels for all cells in the image, total cell area is the number of pixels of all cells, background fluorescence is the average mean gray value of nearby regions containing no cells, and Ncells is the number of cells that was measured by counting fluorescently labeled nuclei from images.

Images were deconvolved using Huygens Professional software (Hilversum, Netherlands)

2.7 Statistical analysis

Experimental data are shown as mean $\pm$ standard error of the mean (S.E.M.). Analysis of variance (ANOVA) and Tukey’s test were performed using R software (version 3.5.3; <https://www.r-project.org/>). Probabilities of less than 5% ($P < 0.05$) indicate statistically significant differences.

3. Results

3.1 Effects of EAS on P4 production

We investigated the effects of various doses of EAS on P4 secretion in bovine GCs. As shown in Figure 1, we found that EAS significantly ($p < 0.001$) promoted P4 levels peaking at 5 mg/mL. Collectively, EAS showed a dose-dependent induction of P4 production.

3.2 Effects of EAS on P4 production and steroidogenic and lipid metabolism gene expression in bovine GCs under non-HS and HS conditions

In our previous study, we showed that EAS induces HSP70 and improves redox balance in both non-HS and HS conditions [21]. In the present study, we investigated whether EAS affects P4 synthesis in both environments, together with the expression of steroidogenesis-related genes in EAS-treated bovine GCs, under non-HS and HS conditions. As shown in Figure 2A, P4 levels were significantly ($p < 0.001$) increased by EAS compared with the untreated control. However, P4 levels in HS-treated cells tended to be lower than that for the control cells ($p < 0.1$) (Figure 2A). Interestingly, P4 levels were significantly increased ($p < 0.001$) by EAS under HS conditions (Figure 2A).

Furthermore, *STAR* and *3 β -HSD* mRNA expression levels were similar to P4 levels in the EAS and EAS+HS groups ($p < 0.001$) (Figure 2B, 2C). In contrast, EAS did not affect the expression of *CYP11A1*, *progesterone receptor*, or *estrogen receptor* under either non-HS or HS conditions ($p > 0.05$) (Figure 2D and Figure S1).

Moreover, *SREBP-1* expression was significantly increased in both the EAS and EAS+HS groups, similar to the pattern of P4 levels and expression of steroidogenic genes ($p < 0.05$) (Figure 2E).

These results confirmed that EAS-induced P4 production is affected by the enhancement of *STAR*, *3 β -HSD*, and *SREBP-1*, and it has the greatest effect on P4 synthesis in bovine GCs

under HS conditions. Notably, HS treatment had a synergistic effect with EAS on the increase in *STAR*, *3 β -HSD*, and *SREBP-1*.

3.3 Effects of EAS on lipid metabolism and mitochondrial activity in bovine GCs under non-HS and HS conditions

To clarify the effects of EAS on P4 levels, steroidogenesis, and lipid metabolism gene expression, we examined lipid synthesis and mitochondrial function in EAS-treated cells under non-HS and HS conditions. As shown in Figure 3A, EAS increased lipid droplets and mitochondrial activity under non-HS and HS conditions. Remarkably, the EAS-induced HSP70 protein was expressed in both the nucleus and cytoplasm, but HS-induced HSP70 was localized mainly in the nucleus (Figure 3A).

Next, we deconvoluted the acquired lipid metabolism and mitochondrial activity images. Deconvolution of the images efficiently removes background noise pixels in lipid droplets and mitochondrial images of African green monkey kidney fibroblast cells [24]. After deconvolution, the lipid droplets and mitochondrial structure appeared to be assembled from multiple particles (Figure 3A). Notably, the localization of lipid droplets and mitochondrial function were mainly observed in the cytosol around the nucleus (Figure 3A). The results show that EAS increased lipid droplets and mitochondrial particles under non-HS and HS conditions.

Similar to P4 synthesis, the fluorescence levels of lipid droplets and mitochondria in the EAS+HS group were significantly ($p < 0.001$) higher than those in the other treatment groups (Figure 3B, C). HS significantly ($p < 0.05$) increased the number of lipid droplets but significantly ($p < 0.01$) reduced mitochondrial activity (Figure 3B, C). These results show that EAS and its synergistic effect with HS increased lipid metabolism and mitochondrial activity, and these results may be related to EAS-induced HSP70.

3.4 Inhibition of HSP70 attenuates P4 synthesis, lipid metabolism, and mitochondrial activity in EAS-treated bovine GCs under non-HS and HS conditions

We hypothesized that EAS-induced HSP70 may be involved in the effects of EAS on P4 synthesis, lipid droplet formation, and mitochondrial function in bovine GCs under non-HS and HS conditions. To test this hypothesis, we evaluated the effects of HSP70 inhibition on P4 synthesis, steroidogenic gene expression, lipid synthesis, and mitochondrial activity in EAS-treated cells under both non-HS and HS conditions. As expected, PES significantly decreased EAS-induced P4 synthesis related to reduced P4, *STAR*, *3 β -HSD*, and *SREBP-1* levels under both non-HS and HS conditions ($p < 0.05$) (Figure 4A, B, C, E). However, an HSP70 inhibitor did not affect *CYP11A1* expression ($p > 0.05$) (Figure 4D).

In addition, HSP70 blockage inhibited the EAS-mediated upregulation of fluorescent levels of lipid droplets and mitochondria, whereas HSP70 expression was unchanged (Figure 5). Consistently, PES suppressed lipid synthesis and mitochondrial activity in EAS-treated

bovine GCs under non-HS and HS conditions ($p < 0.05$) (Figure 6). Taken together, these data suggest that EAS-induced P4 synthesis is regulated by EAS-induced HSP70.

4. Discussion

In the present study, we confirmed that EAS induces HSP70-mediated P4 synthesis through steroidogenic genes associated with the induction of lipid metabolism and mitochondrial activity in bovine GCs under both non-HS and HS conditions. Notably, EAS treatment under HS treatment synergistically increased P4 synthesis with increased expression of *STAR*, *3 β -HSD*, and *SREBP-1*, lipid metabolism, and mitochondrial activity.

The present results show that P4 levels in bovine GCs were highest when treated with 5 mg/mL EAS. This result was similar to that of our previous study in which bovine GCs were treated with or without 5 mg/mL EAS [21]. EAS exerts beneficial effects during stress and improves sleep quality in humans [25]. Other studies have revealed beneficial effects of EAS on brain cell function [26]. Recent evidence has suggested that the central nervous system can synthesize steroid hormones from cholesterol [27]. P4 synthesis is initiated by cholesterol import into the mitochondria by *STAR*, after which *CYP11A1* transforms cholesterol to pregnenolone, which is converted into P4 by *3 β -HSD* [28]. In GCs, P4 binds to the progesterone receptor and facilitates its activity in gene transcription and cellular responses [29]. Under *in vitro* conditions, bovine GCs are good models for P4 studies because of luteinization of this cell line, which involves the switch from the production of

estrogens to P4 [30]. In the present study, EAS strongly increased P4 levels and the expression of *STAR* and *3 β -HSD*, but did not affect the expression of *CYP11A1*, *progesterone receptor*, and *estrogen receptor*. These results suggest that EAS regulates P4 synthesis through *STAR* and *3 β -HSD* pathways in the mitochondria of bovine GCs. In addition, EAS enhanced P4 levels but did not affect P4 related signal pathway of GCs through the progesterone receptor in an autocrine or paracrine manner. Furthermore, HS can synergistically promote this pathway with EAS treatment, resulting in promotion of P4 synthesis. Importantly, mitochondria play a central role in steroidogenesis enzymatic activity [6,7]. Surprisingly, in our results, EAS increased mitochondrial activity in response to HS, despite the fact that HS itself decreased mitochondrial activity.

Generally, HS in the summer season is known to cause a decrease in ovarian function, with reduced P4 levels and corpus luteum size [31–33]. In mammalian cells, lipid droplets are the cholesterol preservation sites for steroid hormones [34]. In the fat tissue of Japanese black cattle, *SREBP-1* affects lipid metabolism [35]. Our present results show that lipid droplets and *SREBP-1* were induced by both EAS and HS at the same level, and the highest expression levels were observed after co-treatment with EAS and HS. Collectively, EAS-induced P4 synthesis through enhancement of *STAR*, *3 β -HSD* enzyme activity, mitochondrial activity, and lipid metabolism in bovine GCs. In addition, significant effect of EAS and HS co-treatment on P4 synthesis was also found in the present study.

In our previous study, EAS was shown to enhance redox balance through the reduction of ROS levels under ordinary non-HS conditions, and, surprisingly, EAS had a significant effect on HSP70 induction and ROS reduction in bovine GCs even under HS conditions compared with the increased ROS and decreased GSH under HS conditions without EAS [21].

In a previous study, STAR, a key enzyme in P4 synthesis, was shown to be inhibited by ROS in rat Leydig cells [36]. ROS also blocks P4 synthesis through CYP11A1 and 3 β -HSD in human granulosa luteal cells [37]. In bovine GCs, HS-induced ROS reduce *STAR* and *CYP11A1* gene expression [13]. Hydrogen peroxide, the most popular ROS, decreases P4 levels, *STAR*, and *3 β -HSD* gene expression in bovine GCs [38]. In addition, high concentrations of ROS negatively affect mitochondria in mouse GCs cells [39]. Therefore, redox balance might also be important in regulating steroidogenesis-related genes and mitochondrial activity in bovine GCs.

In addition, lipid droplets are markers of cellular stress, and they play an important role in maintaining redox status under stress conditions [40]. Under stress conditions, lipid droplets sequester toxic lipids and delay the release of lipids, and this lipid droplet biosynthesis maintains redox balance [40]. Thus, EAS-treated GCs in this study, while maintaining redox balance, possibly caused the stabilization of lipid droplets as a source of cholesterol compared to non-EAS-treated cells. Collectively, the enhancement of P4 synthesis by EAS treatment while maintaining redox status depends on the ROS levels in bovine GCs.

In the present study, HS induced lipid droplet formation and reduced mitochondrial activity in bovine GCs. Our previous study suggested that ROS and ROS-induced DNA damage are detected mainly in the nuclei of bovine GCs under HS conditions [21]. In bovine somatic cell nuclear transfer embryos, mitochondrial and DNA damage are induced by an increase in ROS [41]. Thus, impairment of DNA damage in the nucleus reduces mitochondrial function in HS bovine GCs and affects gene expression. Previous studies have shown that lipid droplets have dynamic functions in cellular metabolism [42]. In mice, mitochondrial damage is caused by hypoxia-induced formation of lipid droplets [43]. Thus, the high levels of lipid droplets in bovine GCs induced by HS can be explained by mitochondrial dysfunction. Collectively, in our results, HS-induced ROS damage in the nucleus impaired mitochondrial function, resulting in the accumulation of lipid droplets in bovine GCs. EAS and EAS plus HS treatment also induced lipid droplet formation, and these results are related with expression of HSP70 that will be discussed in more details below. In our previous study, overall data confirmed that the beneficial effects of EAS-induced HSP70 are mediated by regulating redox balance and protecting cells from HS-related harmful effects in bovine GCs [21].

To show the relationship between HSP70, the steroidogenesis pathway, and mitochondrial function, we inhibited HSP70 activity. Inhibition of HSP70 by PES did not affect the expression of HSP70 protein; however, HSP70 inhibition clearly reduced mitochondrial activity and lipid droplet synthesis. Previous studies have shown that PES disrupts the co-chaperone and substrate proteins of HSP70 without affecting HSP70 expression; thus, PES

inhibits the function of HSP70 protein in multiple cell signaling pathways [44]. In the present study, PES strongly inhibited P4 synthesis, expression of *STAR*, *3 β -HSD*, and *SREBP-1*, mitochondrial activity, and lipid droplet synthesis in bovine GCs, even in the presence of EAS, under both non-HS and HS conditions. Therefore, these data strongly support the hypothesis that an increase in EAS-induced HSP70 promotes P4 synthesis. In addition, HSP70 localization was detected in both the nucleus and cytoplasm of EAS-treated cells, particularly under HS conditions. In mammalian cells, HSP70 has chaperone functions in both the nucleus and cytoplasm [45,46].

In our previous study, EAS-induced HSP70 reduces DNA damage by reducing ROS, which are mainly generated in the nucleus in bovine GCs [21]. Modulation of redox balance-regulated mitochondrial activity and lipid droplets reduces ROS damage in the nuclei of bovine GCs. In addition, the transcription of steroidogenesis enzymes mainly occurs in the nucleus [47]. Therefore, EAS-induced HSP70 balances the redox status to improve steroidogenesis enzyme activity, mitochondrial function, and lipid synthesis in bovine GCs.

On the other hand, synthetic chemical chaperones increase 3 β -HSD metabolic activity in the inner mitochondrial membrane by promoting the folding speed of 3 β -HSD from the native to the active state [48]. In rat adipocyte lipid droplets, HSP70 stimulation is involved in stabilizing the droplet monolayer, protein folding, and mobilization of nascent proteins to the lipid droplets [49]. The size of lipid droplets and fat accumulation are increased by HSP70 through the enhancement of *SREBP-1* in mouse liver cells [50]. Therefore, EAS-induced HSP70

enhances the accumulation of lipid droplets and 3 β -HSD activity in the mitochondria in the cytoplasm with the chaperone function of bovine GCs. In the present study, promotion effect of EAS on P4 synthesis via steroidogenesis pathway, mitochondrial functions, lipid synthesis with synergistic augmentation with HS. These results suggest the EAS is a potential tool for improving the reproductive functions by regulating of intracellular redox and molecular chaperone. Recent study has revealed that the HSP inducer in the EAS was purified and identified as a mixture of Asparagus-Derived Proline-Containing 3-Alkyldiketopiperazines (Asparaprolines) [20]. However, evaluation of such components on the detailed cellular functions is not clarified. Further study will clarify the beneficial effect on cellular functions and regulation of reproductive performance.

5.Conclusion

P4 synthesis by EAS is mediated by the steroidogenesis pathway via HSP70-regulated activation of STAR and 3 β -HSD, together with improved mitochondrial activity and lipid metabolism in bovine GCs. Moreover, EAS has synergistic effect of with HSP70-regulated steroidogenesis pathway. These findings provide a basis for the application of EAS to improve reproductive functions in mammals, including cattle, in combination with HS treatment.

Supplementary materials:

Table S1: Primer sequences used for qRT-PCR

Figure S1: Effects of EAS on progesterone receptor and estrogen receptor in bovine GCs cells under non-HS and HS conditions

Acknowledgements

We would like to thank Editage (www.editage.com) for English language editing.

Funding

This study was supported by the Grant-in-Aid for Scientific Research (2021) from the Tojuro Iijima Foundation for Food Science and Technology.

Availability of data and materials: All data generated through this study are included in this article.

Ethics approval and consent to participate: Not Applicable . All experiments were carried out by cultured cells collected from ovaries at slaughter house.

Competing interests: The authors declare no conflicts of interest.

5. References

- [1] T. Fair, P. Lonergan, The role of progesterone in oocyte acquisition of developmental competence, *Reprod Domest Anim.* 47 Suppl 4 (2012) 142–147.
<https://doi.org/10.1111/j.1439-0531.2012.02068.x>.
- [2] F.W. Bazer, T.E. Spencer, G.A. Johnson, R.C. Burghardt, Uterine receptivity to implantation of blastocysts in mammals, *Front Biosci (Schol Ed)*. 3 (2011) 745–767.
<https://doi.org/10.2741/s184>.
- [3] R. Fatehi A., Roelen, B., Colenbrander, B., Schoevers, E., Gadella, B., Bevers, M., & Van den Hurk, Presence of cumulus cells during in vitro fertilization protects the bovine oocyte against oxidative stress and improves first cleavage but does not affect further development, *Zygote*. 13 (2005) 177–185.
<https://doi.org/doi:10.1017/S0967199405003126>.
- [4] M.A. Abedel-Majed, S.M. Romereim, J.S. Davis, A.S. Cupp, Perturbations in Lineage Specification of Granulosa and Theca Cells May Alter Corpus Luteum Formation and Function, *Frontiers in Endocrinology*. 10 (2019).
<https://doi.org/10.3389/fendo.2019.00832>.
- [5] J.C. Chapman, J.R. Polanco, S. Min, S.D. Michael, Mitochondrial 3 beta-hydroxysteroid dehydrogenase (HSD) is essential for the synthesis of progesterone by corpora lutea: An hypothesis, *Reproductive Biology and Endocrinology*. 3 (2005) 11.
<https://doi.org/10.1186/1477-7827-3-11>.

- 419 [6] G. Bassi, S.K. Sidhu, S. Mishra, The Expanding Role of Mitochondria, Autophagy and
420 Lipophagy in Steroidogenesis, *Cells*. 10 (2021) 1851.
421 <https://doi.org/10.3390/cells10081851>.
- 422 [7] M.B. Rone, J. Fan, V. Papadopoulos, Cholesterol transport in steroid biosynthesis: role
423 of protein-protein interactions and implications in disease states, *Biochim Biophys Acta*.
424 1791 (2009) 646–658. <https://doi.org/10.1016/j.bbalip.2009.03.001>.
- 425 [8] S. Elis, A. Desmarchais, V. Maillard, S. Uzbekova, P. Monget, J. Dupont, Cell
426 proliferation and progesterone synthesis depend on lipid metabolism in bovine
427 granulosa cells, *Theriogenology*. 83 (2015) 840–853.
428 <https://doi.org/10.1016/j.theriogenology.2014.11.019>.
- 429 [9] R. Bertolio, F. Napoletano, M. Mano, S. Maurer-Stroh, M. Fantuz, A. Zannini, S. Bicciato,
430 G. Sorrentino, G. Del Sal, Sterol regulatory element binding protein 1 couples
431 mechanical cues and lipid metabolism, *Nat Commun*. 10 (2019) 1326.
432 <https://doi.org/10.1038/s41467-019-09152-7>.
- 433 [10] M.C. Lucy, ADSA Foundation Scholar Award - Reproductive loss in high-producing dairy
434 cattle: Where will it end?, *Journal of Dairy Science*. 84 (2001) 1277–1293.
435 [https://doi.org/10.3168/jds.S0022-0302\(01\)70158-0](https://doi.org/10.3168/jds.S0022-0302(01)70158-0).
- 436 [11] S.J. Wilson, R.S. Marion, J.N. Spain, D.E. Spiers, D.H. Keisler, M.C. Lucy, Effects of
437 controlled heat stress on ovarian function of dairy cattle. 1. Lactating cows, *Journal of*

438 Dairy Science. 81 (1998) 2124–2131. <https://doi.org/10.3168/jds.S0022->
439 0302(98)75788-1.

440 [12] B.U. Wakayo, P.S. Brar, S. Prabhakar, Review on mechanisms of dairy summer
441 infertility and implications for hormonal intervention, *Open Vet J.* 5 (2015) 6–10.

442 [13] A. Khan, J.H. Dou, Y.C. Wang, X.L. Jiang, M.Z. Khan, H.P. Luo, T. Usman, H.B. Zhu,
443 Evaluation of heat stress effects on cellular and transcriptional adaptation of bovine
444 granulosa cells, *Journal of Animal Science and Biotechnology.* 11 (2020).
445 <https://doi.org/10.1186/s40104-019-0408-8>.

446 [14] B.M. Pizarro, A. Cordeiro, M.W. Reginatto, S.P.C. Campos, A.C.A. Mancebo, P.C.F.
447 Areas, R.A. Antunes, M. do C.B. Souza, K.J. Oliveira, F.F. Bloise, E. Bloise, R.S.
448 Fortunato, T.M. Ortega-Carvalho, Estradiol and Progesterone Levels are Related to
449 Redox Status in the Follicular Fluid During In Vitro Fertilization, *Journal of the Endocrine*
450 *Society.* 4 (2020). <https://doi.org/10.1210/jendso/bvaa064>.

451 [15] M.J. de los Santos, V. García-Láez, D. Beltrán-Torregrosa, J.A. Horcajadas, J.A.
452 Martínez-Conejero, F.J. Esteban, A. Pellicer, E. Labarta, Hormonal and molecular
453 characterization of follicular fluid, cumulus cells and oocytes from pre-ovulatory follicles
454 in stimulated and unstimulated cycles, *Hum Reprod.* 27 (2012) 1596–1605.
455 <https://doi.org/10.1093/humrep/des082>.

- 456 [16] R.J. Rodgers, H.F. Irving-Rodgers, Formation of the ovarian follicular antrum and
 457 follicular fluid, Biol Reprod. 82 (2010) 1021–1029.
 458 <https://doi.org/10.1095/biolreprod.109.082941>.
- 459 [17] W.J. Murdoch, Inhibition by oestradiol of oxidative stress-induced apoptosis in pig
 460 ovarian tissues, Journal of Reproduction and Fertility. 114 (1998) 127–130.
- 461 [18] B. Shi, D. Feng, M. Sagnelli, J. Jiao, X. Sun, X. Wang, D. Li, Fructose levels are elevated
 462 in women with polycystic ovary syndrome with obesity and hyperinsulinemia, Hum
 463 Reprod. 35 (2020) 187–194. <https://doi.org/10.1093/humrep/dez239>.
- 464 [19] H. Saibil, Chaperone machines for protein folding, unfolding and disaggregation, Nature
 465 Reviews Molecular Cell Biology. 14 (2013) 630–642. <https://doi.org/10.1038/nrm3658>.
- 466 [20] S. Inoue, J. Takanari, K. Abe, A. Nagayama, Y. Ikeya, N. Kohda, Isolation and Structure
 467 Determination of a Heat Shock Protein Inducer, Asparagus-Derived Proline-Containing
 468 3-Alkyldiketopiperazines (Asparaprolines), From a Standardized Extract of Asparagus
 469 officinalis Stem, Natural Product Communications. 15 (2020).
 470 <https://doi.org/10.1177/1934578x20914681>.
- 471 [21] K.T. Ho, K. Homma, J. Takanari, H. Bai, M. Kawahara, K.T.K. Nguyen, M. Takahashi, A
 472 standardized extract of Asparagus officinalis stem improves HSP70-mediated redox
 473 balance and cell functions in bovine cumulus-granulosa cells, Sci Rep. 11 (2021) 18175.
 474 <https://doi.org/10.1038/s41598-021-97632-6>.

- 475 [22] T. Ito, A. Sato, T. Ono, K. Goto, T. Maeda, J. Takanari, H. Nishioka, K. Komatsu, H.
 476 Matsuura, Isolation, Structural Elucidation, and Biological Evaluation of a 5-
 477 Hydroxymethyl-2-furfural Derivative, Asfural, from Enzyme-Treated Asparagus Extract,
 478 Journal of Agricultural and Food Chemistry. 61 (2013) 9155–9159.
 479 <https://doi.org/10.1021/jf402010c>.
- 480 [23] O. Gavet, J. Pines, Activation of cyclin B1-Cdk1 synchronizes events in the nucleus and
 481 the cytoplasm at mitosis, Journal of Cell Biology. 189 (2010) 247–259.
 482 <https://doi.org/10.1083/jcb.200909144>.
- 483 [24] M. Pribasnig, B. Kien, L. Pusch, G. Haemmerle, R. Zimmermann, H. Wolinski, Extended-
 484 resolution imaging of the interaction of lipid droplets and mitochondria, Biochimica et
 485 Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids. 1863 (2018) 1285–1296.
 486 <https://doi.org/10.1016/j.bbalip.2018.07.008>.
- 487 [25] T. Ito, K. Goto, J. Takanari, T. Miura, K. Wakame, H. Nishioka, A. Tanaka, J. Nishihira,
 488 Effects of enzyme-treated asparagus extract on heat shock protein 70, stress indices,
 489 and sleep in healthy adult men, J Nutr Sci Vitaminol (Tokyo). 60 (2014) 283–290.
 490 <https://doi.org/10.3177/jnsv.60.283>.
- 491 [26] M. Takahashi, T. Sakurai, T. Hisajima, Effect of Enzyme-Treated Asparagus Extract on
 492 brain function, The FASEB Journal. 30 (2016) 1176.10-1176.10.
 493 https://doi.org/10.1096/fasebj.30.1_supplement.1176.10.

- 494 [27] K.A. Bates, A.R. Harvey, M. Carruthers, R.N. Martins, Androgens, andropause and
495 neurodegeneration: exploring the link between steroidogenesis, androgens and
496 Alzheimer's disease, *Cell Mol Life Sci.* 62 (2005) 281–292.
497 <https://doi.org/10.1007/s00018-004-4383-1>.
- 498 [28] A.J. Conley, I.M. Bird, The role of cytochrome P450 17 alpha-hydroxylase and 3 beta-
499 hydroxysteroid dehydrogenase in the integration of gonadal and adrenal
500 steroidogenesis via the delta 5 and delta 4 pathways of steroidogenesis in mammals,
501 *Biol Reprod.* 56 (1997) 789–799. <https://doi.org/10.1095/biolreprod56.4.789>.
- 502 [29] J.J. Peluso, Progesterone as a regulator of granulosa cell viability, *The Journal of Steroid*
503 *Biochemistry and Molecular Biology.* 85 (2003) 167–173. [https://doi.org/10.1016/S0960-](https://doi.org/10.1016/S0960-0760(03)00192-4)
504 [0760\(03\)00192-4](https://doi.org/10.1016/S0960-0760(03)00192-4).
- 505 [30] E. Montaña, M. Olivera, Z.T. Ruiz-Cortés, Association between leptin, LH and its
506 receptor and luteinization and progesterone accumulation (P4) in bovine granulosa cell
507 in vitro, *Reprod Domest Anim.* 44 (2009) 699–704. [https://doi.org/10.1111/j.1439-](https://doi.org/10.1111/j.1439-0531.2007.01053.x)
508 [0531.2007.01053.x](https://doi.org/10.1111/j.1439-0531.2007.01053.x).
- 509 [31] J.L. Howell, J.W. Fuquay, A.E. Smith, Corpus Luteum Growth and Function in Lactating
510 Holstein Cows During Spring and Summer, *Journal of Dairy Science.* 77 (1994) 735–
511 739. [https://doi.org/10.3168/jds.S0022-0302\(94\)77007-7](https://doi.org/10.3168/jds.S0022-0302(94)77007-7).
- 512 [32] J.M. Burke, D.E. Spiers, F.N. Kojima, G.A. Perry, B.E. Salfen, S.L. Wood, D.J. Patterson,
513 M.F. Smith, M.C. Lucy, W.G. Jackson, E.L. Piper, Interaction of endophyte-infected

514 fescue and heat stress on ovarian function in the beef heifer, *Biol Reprod.* 65 (2001)
515 260–268. <https://doi.org/10.1095/biolreprod65.1.260>.

516 [33] D. Wolfenson, H. Sonego, A. Bloch, A. Shaham-Albalancy, M. Kaim, Y. Folman, R.
517 Meidan, Seasonal differences in progesterone production by luteinized bovine thecal
518 and granulosa cells, *Domest Anim Endocrinol.* 22 (2002) 81–90.
519 [https://doi.org/10.1016/s0739-7240\(01\)00127-8](https://doi.org/10.1016/s0739-7240(01)00127-8).

520 [34] D. Zweytick, K. Athenstaedt, G. Daum, Intracellular lipid particles of eukaryotic cells,
521 *Biochim Biophys Acta.* 1469 (2000) 101–120. [https://doi.org/10.1016/s0005-](https://doi.org/10.1016/s0005-2736(00)00294-7)
522 [2736\(00\)00294-7](https://doi.org/10.1016/s0005-2736(00)00294-7).

523 [35] S. Hoashi, N. Ashida, H. Ohsaki, T. Utsugi, S. Sasazaki, M. Taniguchi, K. Oyama, F.
524 Mukai, H. Mannen, Genotype of bovine sterol regulatory element binding protein-1
525 (SREBP-1) is associated with fatty acid composition in Japanese Black cattle, *Mamm*
526 *Genome.* 18 (2007) 880–886. <https://doi.org/10.1007/s00335-007-9072-y>.

527 [36] S.-C. Tsai, C.-C. Lu, C.-S. Lin, P.S. Wang, Antisteroidogenic actions of hydrogen
528 peroxide on rat Leydig cells, *J Cell Biochem.* 90 (2003) 1276–1286.
529 <https://doi.org/10.1002/jcb.10738>.

530 [37] T. Endo, R.F. Aten, L. Leykin, H.R. Behrman, Hydrogen peroxide evokes
531 antisteroidogenic and antigonadotropic actions in human granulosa luteal cells, *The*
532 *Journal of Clinical Endocrinology & Metabolism.* 76 (1993) 337–342.
533 <https://doi.org/10.1210/jcem.76.2.7679398>.

534 [38] M. Wang, Y. Li, A. Molenaar, Q. Li, Y. Cao, Y. Shen, P. Chen, J. Yan, Y. Gao, J. Li,
535 Vitamin E and selenium supplementation synergistically alleviate the injury induced by
536 hydrogen peroxide in bovine granulosa cells, *Theriogenology*. 170 (2021) 91–106.
537 <https://doi.org/10.1016/j.theriogenology.2021.04.015>.

538 [39] S.A.M. Hoque, T. Umehara, T. Kawai, M. Shimada, Adverse effect of superoxide-
539 induced mitochondrial damage in granulosa cells on follicular development in mouse
540 ovaries, *Free Radic Biol Med*. 163 (2021) 344–355.
541 <https://doi.org/10.1016/j.freeradbiomed.2020.12.434>.

542 [40] E. Jarc, T. Petan, Lipid Droplets and the Management of Cellular Stress, *Yale J Biol*
543 *Med*. 92 (2019) 435–452.

544 [41] I.-S. Hwang, H.-K. Bae, H.-T. Cheong, Mitochondrial and DNA damage in bovine
545 somatic cell nuclear transfer embryos, *J Vet Sci*. 14 (2013) 235–240.
546 <https://doi.org/10.4142/jvs.2013.14.3.235>.

547 [42] J.A. Olzmann, P. Carvalho, Dynamics and functions of lipid droplets, *Nat Rev Mol Cell*
548 *Biol*. 20 (2019) 137–155. <https://doi.org/10.1038/s41580-018-0085-z>.

549 [43] S.-J. Lee, J. Zhang, A.M.K. Choi, H.P. Kim, Mitochondrial dysfunction induces formation
550 of lipid droplets as a generalized response to stress, *Oxid Med Cell Longev*. 2013 (2013)
551 327167. <https://doi.org/10.1155/2013/327167>.

552 [44] J.I.-J. Leu, J. Pimkina, A. Frank, M.E. Murphy, D.L. George, A small molecule inhibitor
 553 of inducible heat shock protein 70, *Mol Cell.* 36 (2009) 15–27.
 554 <https://doi.org/10.1016/j.molcel.2009.09.023>.

555 [45] A.A. Michels, B. Kanon, A.W. Konings, K. Ohtsuka, O. Bensaude, H.H. Kampinga,
 556 Hsp70 and Hsp40 chaperone activities in the cytoplasm and the nucleus of mammalian
 557 cells, *J Biol Chem.* 272 (1997) 33283–33289. <https://doi.org/10.1074/jbc.272.52.33283>.

558 [46] N. Imamoto, Heat stress-induced nuclear transport mediated by Hsp70s confers nuclear
 559 function of Hsp70s, *Current Opinion in Cell Biology.* 52 (2018) 82–87.
 560 <https://doi.org/10.1016/j.ceb.2018.02.010>.

561 [47] T. Mizutani, S. Ishikane, S. Kawabe, A. Umezawa, K. Miyamoto, Transcriptional
 562 regulation of genes related to progesterone production [Review], *Endocrine Journal.* 62
 563 (2015) 757–763. <https://doi.org/10.1507/endocrj.EJ15-0260>.

564 [48] J.L. Thomas, H.S. Bose, Regulation of human 3-beta-hydroxysteroid dehydrogenase
 565 type-2 (3βHSD2) by molecular chaperones and the mitochondrial environment affects
 566 steroidogenesis, *J Steroid Biochem Mol Biol.* 151 (2015) 74–84.
 567 <https://doi.org/10.1016/j.jsbmb.2014.11.018>.

568 [49] H. Jiang, J. He, S. Pu, C. Tang, G. Xu, Heat shock protein 70 is translocated to lipid
 569 droplets in rat adipocytes upon heat stimulation, *Biochim Biophys Acta.* 1771 (2007) 66–
 570 74. <https://doi.org/10.1016/j.bbalip.2006.10.004>.

[50] J. Zhang, N. Fan, Y. Peng, Heat shock protein 70 promotes lipogenesis in HepG2 cells, Lipids in Health and Disease. 17 (2018) 73. <https://doi.org/10.1186/s12944-018-0722-8>.

Author contributions

Masashi Takahashi: Conceptualization, Supervision, Formal analysis, Writing-review & editing, Project administration. **Khoi Thieu Ho:** Conceptualization, Validation, Formal analysis, Investigation, Visualization, Writing-original draft, Writing-review & editing. **Hanako Bai:** Methodology, Writing-review & editing. **Manabu Kawahara:** Methodology, Writing-review & editing. **Kohei Homma:** Resources, Writing-review & editing. **Jun Takanari:** Resources, Writing-review & editing. **Ahmed Zaky Balboula:** Formal analysis, Writing-review & editing. **Khang Thi Kim Nguyen:** Formal analysis, Writing-review & editing.

Figure legends

Figure 1. Effect of various concentrations of EAS on P4 production in bovine GCs.

Cells were subjected to various concentrations of EAS (0.5, 1, 5, and 10 mg/mL) for 12 h at 38.5°C (n = 5). Detection of P4 secretion was performed with ELISA in cell supernatants. Data are shown as the mean $\pm$ S.E.M, a vs. b (p < 0.001), a vs. c (p < 0.05), and b vs. c (p < 0.05).

Figure 2. Effect of EAS on P4 synthesis and expression of steroidogenic genes in bovine GCs under non-HS and HS conditions.

Cells were treated with or without 5 mg/mL of EAS at 38.5°C (control, EAS) or 41°C (HS, HS + EAS) for 12 h. Detection of P4 secretion was performed with ELISA in cell supernatants (A, n = 3). The mRNA expression levels of STAR, 3 β -HSD, CYP11A1, and SREBP-1 (B–E, respectively, n = 5) are presented relative to H2AFZ as a reference gene. Data are shown as the mean $\pm$ S.E.M, (A) a vs. b (p < 0.001), a vs. c (p < 0.001), b vs. c (p < 0.05), (B) a vs. b (p < 0.05), a vs. c (p < 0.001), b vs. c (p < 0.05), (C) a vs. b (p < 0.01), a vs. c (p < 0.001), and b vs. c (p < 0.001), and (E) a vs. b (p < 0.05), a vs. c (p < 0.001), b vs. c (p < 0.05).

Figure 3. Effect of EAS on the synthesis of lipid droplets and mitochondrial activity in bovine GCs under non-HS and HS conditions.

Cells were treated with or without 5 mg/mL of EAS at 38.5°C (control, EAS) or at 41°C (HS, HS + EAS) for 12 h. Fluorescence staining and enlarged image of mitochondria, lipid droplets, and HSP70 are shown (A). Scale bar = 50 µm; original magnification = 63×. The fluorescence intensity of mitochondria and lipid droplets was analyzed using CTCF (B and C, respectively, n = 5). Data are shown as the mean ± S.E.M, (B) a vs. b (p < 0.05), a vs. c (p < 0.001), b vs. c (p < 0.001), (C) a vs. b (p < 0.01), a vs. c (p < 0.01), a vs. d (p < 0.001), b vs. c (p < 0.001), b vs. d (p < 0.001), c vs. d (p < 0.001).

Figure 4. Effect of HSP70 inhibition on EAS-induced P4 synthesis and expression of steroidogenic genes in bovine GCs under non-HS and HS conditions.

Cells were treated with or without 10 µM PES to 38.5°C (control, EAS, EAS + PES) or at 41°C (HS, HS + EAS, HS + EAS + PES) for 12 h. Detection of P4 secretion was performed with ELISA in cell supernatants (A, n = 3). The mRNA expression levels of STAR, 3β-HSD, CYP11A1, and SREBP-1 (B–E, respectively, n = 5) are presented relative to H2AFZ as a reference gene. Data are shown as the mean ± S.E.M, (A) a vs. b (p < 0.05), a vs. c (p < 0.001), and b vs. c (p < 0.01), (B) a vs. b (p < 0.001), a vs. c (p < 0.001), and b vs. c (p < 0.05); (C) a vs. b (p < 0.05); a vs. c (p < 0.001), b vs. c (p < 0.001), and (E) a vs. b (p < 0.001), a vs. c (p < 0.001), b vs. c (p < 0.01).

Figure 5. HSP70 inhibitor reversed EAS-induced lipid metabolism and mitochondrial activity in bovine GCs under non-HS and HS conditions.

Cells were treated with or without 10 μ M PES at 38.5°C (control, EAS, EAS + PES) or at 41°C (HS, HS + EAS, HS + EAS + PES) for 12 h. Fluorescence staining images of mitochondria, lipid droplets, and HSP70 are shown. Scale bar = 50 μ m; original magnification = 63 $\times$.

Figure 6. Fluorescence intensity of lipid droplets and mitochondrial activity in bovine GCs treated with EAS under non-HS and HS conditions.

The fluorescence intensity of lipid droplets and mitochondrial activity in cells treated with EAS under non-HS and HS conditions was analyzed by CTCF (A and B, respectively, n = 5). Data are shown as the mean $\pm$ S.E.M, (A) a vs. b ($p < 0.05$), a vs. c ($p < 0.001$), and b vs. c ($p < 0.05$), (B) a vs. b ($p < 0.05$), a vs. c ($p < 0.05$), a vs. d ($p < 0.001$), b vs. c ($p < 0.001$), b vs. d ($p < 0.05$), c vs. d ($p < 0.001$).

Highlights

- EAS increases progesterone synthesis by increasing of steroidogenesis-related genes
- EAS increases lipid synthesis and mitochondrial activity
- Synergistic effect of EAS and heat shock on progesterone synthesis
- HSP70 inhibition decreases EAS-stimulated progesterone synthesis pathway




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

