



HOKKAIDO UNIVERSITY

Title	New Species of Insects from Japan described during the Year 1926
Citation	Insecta matsumurana, 1(4), 207-210
Issue Date	1927-05
Doc URL	https://hdl.handle.net/2115/9126
Type	departmental bulletin paper
File Information	1(4)_p207-210.pdf



NEW SPECIES OF INSECTS FROM JAPAN DESCRIBED
DURING THE YEAR 1926.

RHYNCHOTA

FUKUI T.:—Naichi-san Shokuchu-kamemushi-ka Zusetsu (1) [Kontyu (Insecta), Tokyo, 1, pp. 7-18].

Gardena fusca, p. 13 (Reduviidae).

KATO M.:—The Japanese Cicadidae, with Descriptions of 4 new Species, one new Subspecies and two new aberrant Forms (Trans. Nat. Hist. Soc. Formosa, XIII, pp. 23-29).

Tanna sozanensis, p. 23; *T. sayurie*, p. 25; *T. horiensis* p. 26; *T. viridis* KATO var. *niitakaensis*; *Leptosemia conica*, p. 27; *Mogannia formosana* MATS. subsp. *elegans*, p. 28; *M. formosana* MATS. ab. *interrupta*, p. 29; *Scieroptera formosana* SCHMIEDT ab. *trigutta*; *S. formosana* SCHMIEDT ab. *albifascia*.

KATO M.:—Japanese Cicadidae, with Descriptions of 4 new Species (Trans. Nat. Hist. Soc. Formosa, XVI, pp. 171-175).

Cicada kyushyuensis, p. 171; *Nipponosemia virescens*, p. 172; *Euterpnosia varicolor*, p. 174; *Melampsalta isshikii*, p. 175.

KUWANA I.:—The Diaspine Coccidae of Japan, IV (Bull. Imp. Plant Quar. Yokohama, Japan, pp. 1-44).

Aulacaspis yabunikki, p. 32; *A. wakayamaensis* p. 33.

MATSUMURA S.:—On the five Species of Dendrolimus, injurious to Conifers in Japan, with their parasitic and predaceous Insects (Jour. Coll. Agr. Hokkaido Imp. Univ. XVIII, pp. 1-42).

Nabis kurilensis, p. 41 (Reduviidae).

SHINJI O.:—On new Species *Thoracaphis linderæ* SHINJI (Zool. Mag. Tokyo, XXXVIII, pp. 359-361).

Thoracaphis linderæ, p. 359 (Pemphigidae).

TAKAHASHI R.:—The Aphids of Myzocallis infesting the Bamboo (Proc. Ent. Soc. Washington, 28, pp. 159-162).

Myzocallis taiwanus, p. 160.

HORI M.:—Jagaimo Higenaga-aburamushi ni kansuru Chōsa (Rep. No. 17, Hakkaido Agr. Exp. Station, 51-83).

Macrosiphum matsumuraeanum, p. 52 (Aphididae).

LEPIDOPTERA

HORI H.:—A new Psychid from Japan [Kontyu (Insecta), Tokyo, 1, pp. 28-29].

Eumeca niphonica, p. 28.

HORI H.:—A new Geometrid from Japan [Kontyu (Insecta), 1, pp. 83-85].

Phalaena kiushiuana, p. 83.

HORI H.:—Description of a new Variety of Geometridae (Zool. Mag. Tokyo, XXXVIII, pp. 61-62).

Cidaria corydalaria GRAES. var. *japonica*, p. 61.

KUWAYAMA S.:—Two new Aberrations of Japanese Butterflies (Entomologist, London, Vol. XIX, pp. 180-181).

Lycæna lycornas BUTL. ab. *paucimaculosus*, p. 180; *Chrysophanus phlaeas daimio* SEITZ ab. *distalba*, p. 181.

KÔNO H.:—New aberrant Forms of "*Argynnis rutilana* MOTSCH." and "*Argynnis adippe pallescens* BUTL." (Insect World, Gifu, XXX, pp. 228-230).

Argynnis rutilana MOTSCH. ab. *sapporensis*, p. 228; *A. adippe pallescens* BUTL. ab. *shūriyana*, p. 229.

MATSUMURA S.:—On the five Species of Dendrolimus, injurious to Conifers in Japan, with their

parasitic and predaceous Insects (Jour. Coll. Agr. Hokkaido Imp. Univ. XVIII, pp. 1-48).

Dendrolimus albolineatus MATS. ab. *frequens*, p. 9; ab. *kiminensis*; ab. *nigriscens*; ab. *kurilensis*; ab. *centropallida*, p. 10; ab. *submarginata*; ab. *albata*; ab. *ichinosawana*; ab. *centro-zonalis*; *D. superans* BTLR. ab. *scribae*, p. 13; *D. spectabilis* BTLR. ab. *griseomarginalis*, p. 17; ab. *tokyonis*; ab. *hirosimiana*, p. 18; ab. *koonis*; ab. *shakojiana*; ab. *scribae*; ab. *ryuzana*; ab. *ferruginea*, p. 19; ab. *fallax*; ab. *coreana*; ab. *tricolor*; *D. punctatus* WK. ab. *baibarana*, p. 21; ab. *kantozana*; ab. *pallidiola*.

NAKAHARA W.:—Some Saghalien Butterflies: Notes and Descriptions (Insc. Inscit. Mens., Washington, XIV, pp. 45-48).

Parnassius stubbendorfi esakii, p. 46; *Pieris napi saghalienis*, p. 47.

NAKAHARA W.:—Butterflies from southern Kurile Islands (Insc. Inscit. Mens. pp. 49-51).

Lethe callipteris obscura, p. 50; *Erebia sedakobii doii*; *Pamphila silvius shikotanus*, p. 51.

SONAN J.:—Notes on a few Butterflies from Formosa (Trans. Nat. Hist. Soc. Formosa, XVI, pp. 177-183).

Ixias pyrene insignis BTLR. ab. *breviptera*, p. 177; *Cupha erymanthis* DRUR. ab. *decolorata*; *Phenagris atroguttata matsumurai*, p. 178; *P. atroguttata daiiozana* WILEM. f. *marginata*, p. 180.

SONAN J.:—Notes on a few Butterflies from Formosa and Okinawa (Trans. Nat. Hist. Soc. Formosa, XVI, pp. 230-235).

Euploea leucostictos GMEL. var. *okinawanis*, p. 230; *Hypolimnas bolina priscilla* BUTL. ab. *formosana*, p. 233; *Sphisa chandora androdamas* FRUH. ab. *albifasciata*.

COLEOPTERA

EGGERS H.:—Japanische Borkenkäfer, I (Ent. Blätt. Berlin, 22, pp. 133-138).

Hyorrhynchus niishimai, p. 133; *Sphaerotrypes imitans*, p. 134; *S. carpini*; *Polygraphus sachalinensis*, p. 135; *P. laticollis*; *Cryphalus piceus*, p. 136; *Ernoporus longus*; *Pityogenes foveolatus*, p. 137; *Dryocoetes rugicollis*; *D. picipennis*, p. 138.

KLEINE K.:—Die Lyciden Formosas (Stett. Ent. Zeit. 87, pp. 97-109).

Lycostomus crassus, p. 97; *Plateros purus*, p. 99; *Ditomeces griseostriatus*, p. 101; *Cladophorus certus*, p. 103; *Parapyropterus* (n. g.) *nigrostriatus*, p. 104; *Procautires socius*.

KÔNO H.:—Einige Coleopteren aus Korea, mit Beschreibung einer neuen Art (Trans. Nat. Hist. Soc. Formosa, XVII, pp. 87-91).

Rhynchites koreanus, p. 89.

KÔNO H.:—Beiträge zum Lebenslauf der Japanischen-Attelabiden, *Apoderus fulvus* ROELOFS und *Europs aceri*, n. sp. (Zool. Mag. Tokyo, XXXVIII, pp. 217-224).

Euopus aceri, p. 223.

KÔNO H. et TAMANUKI K.:—Die Käfer-Ausbeute aus Nordsachalin im Jahre 1922 (Zool. Mag. Tokyo, XXXVIII, pp. 276-296).

Corymbites bozberi GERM. ab. *fuscus* p. 281; *Oberea rufomaculata* p. 286; *Chryptocephalus karafutonis*, p. 287; *Phytodecta viminalis* L. ab. *karafutensis*.

VOSS E.:—Die Unterfamilien Attelabinae und Apoderinae (Stett. Ent. Zeit. 87, pp. 1-88).

Paroplapoderus (n. g.) *japonicus*, p. 62; *P. turbidus*, p. 63; f. *funerula*, p. 64; *Phymatopoderus* (n. g.) *pavens*, p. 74; *Tomapoderus* (n. g.) *ruficollis* VOSS f. *ruficeps*, p. 79.

DIPTERA

ALEXANDER C. P.:—New or little known Tipulidae from eastern Asia, pt. 1 (Philipp. Journ. Scien., Vol. 31, No. 3, pp. 363-381).

Erioptera (*Erioptera*) *harukawai*, p. 368; *Gonomyia* (*Ptilostena*) *kuwayamai*; *Cladura bidens*, p. 369; *C. megacauda*, p. 370; *Oropeza shirakiella*, p. 376; *Tipula yamamuriana*, p. 379;

MATSUMURA S.:—On the five Species of Dendrolimus, injurious to Conifers in Japan, with their parasitic and predaceous Insects (Journ. Coll. Agr. Hokkaido Imp. Univ., Vol. XVIII, pp. 1-42).

Echynomyia dendrolimusi, p. 39 (Tachinidae).

HYMENOPTERA

COCKERELL D. A.:—Some Bees in the Collection of the California Academy of Sciences (The Pan-Pacific Entomologist, San Francisco, III, pp. 80-90).

Colletes kobensis, p. 81; *Anthrophora tsushimensis*, p. 84; *Megachile remotissima*, p. 86; *Nomada koreana*, p. 88; *Epeolus tsushimensis*, p. 89.

GAHAN A. B.:—Some Braconid and Chalcid Flies from Formosa, parasitic on Aphids (Proc. U. S. Nat. Hist. 70, pp. 1-7).

Aphidius unilachni, p. 1; *A. laticeps*, p. 2; *A. commodus*, p. 3; *Trioxys communis*, p. 4; *T. struma*, p. 5.

MATSUMURA S.:—On the five Species of Dendrolimus, injurious to Conifers in Japan, with their parasitic and predaceous Insects (Jour. Coll. Agr. Hokkaido Imp. Univ. XVIII, pp. 22-38).

Mesostenus matsukemushii, p. 23; *Hemiteles dendrolimusi*; *Pezomachus dendrolimusi*, p. 24; *Exochilum dendrolimusi*, p. 25; *E. laricis*; *Habronyx matsukemushii*, p. 26; *Mesochorus kuwayamae*, p. 27; *Casinaria takagai*, p. 28; *Pimpla (Apechthis) dendrolimusi*, p. 29; *P. (Eremochila) jezoensis*, p. 30; *P. (Pimpla) spectabilis*; *Chorinaeus dendrolimusi*, p. 32; *Phanomeris dendrolimusi*; *P. spectabilis*, p. 33; *Monodontomerus spectabilis*; *Pteromalus dendrolimusi*, p. 34; *P. matsuyadorii*, p. 35; *P. matsukemushii*; *P. kuwayamae*, p. 36; *Holcaerus (?) dendrolimusi*; *Encyrtus pinicolus*, p. 37; *Trichogramma dendrolimusi*.

MASI L. H.:—Sauter's Formosa-Ausbeute (Konowia, V, pp. 1-20, 170-178, 264-279, 325-381).

Diomorus orientalis, p. 2; *Ecdamua mirabilis*, p. 5; *Dimeromicrus affinis*, p. 7; *Podagrion epibulum*, p. 10; *P. prionomerum*, p. 12; *P. opisthacanthus*, p. 14; *P. tananicum*, p. 16; *P. epichiron*, p. 18; *P. dispar*, p. 19; *P. viduum*, p. 170; *Pachytomoidella helictoscela*, p. 173; *Pachytomoides hornianus*, p. 176; *P. orchestica*, p. 260; *P. megarhopalus*, p. 266; *Microterys mercetii*, p. 268; *Neocladia howardi* PERK. var. *violascens*, p. 272; *Metaplopa* (n. g.) *grallaria*, p. 276; *Eupelmus fuscus*, p. 325; *E. anpingensis*, p. 328; *Calosota (Hylephila)* (n. subg.) *stenozastra*, p. 330; *Sauteria* (n. g.) *eximia*, p. 334; *S. elegans*, p. 338; *Paracleonymus angustatus*, p. 340; *Glyptosticha* (n. g.) *flavipes*, p. 349; *G. nigricans*, p. 353; *Pycnetron curculionides*, p. 356; *Picroscytus ruschkae*, p. 359; *Dinarinus sauteri*, p. 360; *D. pyrromerus*, p. 362; *Eurydinotomorpha sinica*, p. 363; *Schizaspidia vicina*; *S. scutellaris*, p. 370; *Philomides frator*, p. 372; *Perilampus obsoletus*, p. 375; *Euperilampoides concinus*, p. 378 (Chalcididae).

SATO K.:—On the Genus *Janus* STEPHENS, with Descriptions of two new Species from Japan [Kon-yu (Insecta), Tokyo, 1, pp. 16-27].

Janus japonicus, p. 22; *J. micromaculatus*, p. 23 (Tenthredinidae).

TAKEUCHI K.:—A new Sawfly from Okinawa (Trans. Nat. Hist. Soc. Formosa, XVI, pp. 228-229). *Macrophya iuquinana*, p. 228.

UCHIDA T.:—Ueber eine Psithyrus-Art von Sachalin (Zool. Mag. Tokio, XXVIII, pp. 152-155).

Psithyrus naiphtchianus MATS. var. *candidus*, p. 155.

UCHIDA T.:—Ueber die Bethyilden Japans (Zool. Mag. Tokio, XXXVIII, pp. 181-186).

Cleptes galloisi, p. 183.

UCHIDA T.:—Erster Beitrag zur Ichneumoniden-Fauna Japans (Jour. Coll. Agr. Hokkaido Imp. Univ. XVIII, pp. 43-173).

Protichneumon ventralis MATS. var. *segmentalis*, p. 55; *Coelichneumon coxalis*, p. 59; *Coreojoppa* (n. g.) *flavomaculata*, p. 64; *Cobunus* (n. g.) p. 65; *Hoplismenus coreanus*, p. 69; *H. speciosus*; *Chas-mias agitata*, p. 72; do var. *rufipes*; *C. fuscus*, p. 73; *Exephanes apicalis*, p. 74; *E. tibialis*, p. 75;

var. *4-punctata*, p. 76; *Ichneumon nigrifemoratus*, p. 78; *I. toyoharensis*, p. 79; *I. longicellus*, p. 80; *I. magnus*, p. 81; *I. ohtaniensis*; *I. inquinatus* WESM. var. *niphonica*, p. 83; *I. nigricoxalis*, p. 88; *I. longicrus*; *I. enayamensis*, p. 89; *I. rishiriensis*, p. 90; *I. odomariensis*; *I. shisukensis*, p. 91; *Stenichneumon flavolineatus*, p. 94; *S. kamuensis*, p. 95; *S. nigristigmatus*; *S. iwatensis*, p. 96; *S. naganonis*, p. 97; *S. naironis*, p. 98; *S. posticalis* MATS. var. *albomaculata*, p. 99; *Barichneumon bifasciatus*, p. 100; *Cratichneumon quadratus*, p. 101; *Melanichneumon sambonis*, p. 103; *M. femoratus*, p. 104; *M. tiia'i's*, p. 105; *M. marginalis*; *M. albivatus*, p. 106; *M. leucomelas* GM. var. *nipponica*, p. 107; *M. nigricrus*, p. 108; *M. teshionis*, p. 109; *Togea* (n. g.) *albofasciata*, p. 110; *T. nigra*, p. 112; *T. formosana*; *Eupalamus coreanus*, p. 113; *E. flaviventralis*, p. 114; *E. trentepohlii* WESM. var. *japonica*, p. 115; *Pseudochasmodon* (n. g.) *major*, p. 116; *Amblyteles kawakamiensis*, p. 118; *A. kiminaiensis*, p. 119; *A. bifasciatus*, p. 120; *A. flavomaculatus*, p. 123; *A. septentrionis*; *A. diminutus*, p. 124; *A. macilentus* BERTH. var. *japonica*; *A. pentalineatus*, p. 125; *Spilichneumon sachalinensis*, p. 127; *T. nipponicus*; *S. rubrocinctus* TOSQ. var. *taiyudongus*, p. 128; *S. oratorius* F. var. *nigrifrons*, p. 130; *S. jezoensis*; *S. longicornis*, p. 131; *S. unipunctatus*; *S. flavicornis*, p. 132; *S. bipunctatus*; *Ctenichneumon tappanus*, p. 134; *C. kawamurae*, p. 135; *C. okunii*, p. 136; *C. australis*; *C. medius*, p. 137; *C. furumakiensis*, p. 138; *C. kongosanus*, p. 139; *C. omiensis*; *C. infumatus*, p. 140; *C. formosanus*; *C. nigricans*, p. 141; *C. apicalis*, p. 142; *C. urticae*, p. 143; *C. issikii*; *C. sambonis*, p. 145; *Otohimea* (n. g.) p. 147; *O. coreana*, p. 148; *Hepioplumus sachalinensis*, p. 149; *Probolus slaviceki* KRIECH. var. *sachalinensis*, p. 150; *Eurylabus quadratus*, p. 151; *Platylabus iwatensis*, p. 152; *P. nigricornis*, p. 153; *P. hakonensis*; *P. spiraculus*, p. 154; *P. coreanus*, p. 155; *Pristiceros pallidipes*; *P. apicalis*, p. 156; *Phaeogenes yezoensis*, p. 159; *P. similimus*; *P. flavescens*, p. 161; *Colpognathus sapporensis*, p. 162; *Takanona* (n. g.) *ishiyamana*, p. 164; *Monontos* (n. g.) *niphonicus*, p. 165; *Rhexidermus apicalis*, p. 166.

WATERTON J.:—A new Species of Saw-fly (Hym. Tenthredinoidea) from Japan (Entomologist, London, Vol. LIX, pp. 206-209).

Dolerus harukawai, p. 206.

LITERATURE (SYSTEMATIC) PUBLISHED DURING THE YEAR 1926, WHICH CONTAINS JAPANESE INSECTS.

O. BANG-HAAS—Novitates Macrolepidopterologicae, I.

S. SCHENKELING—Colcopterorum Catalogus, Pars 82 (M. BERNHAUER et O. SCHEERPELTZ—Staphylinidae, VI).

„ —Pars 83 (A. HETSCHKO—Thorictidae, Catopochrotidae, Monoedidae, Synteliidae, Cossyphodidae).

„ —Pars 84 (J. OHENBERGER—Buprestidae, I).

„ —Pars 85 (A. HETSCHKO—Lathridiidae).

„ —Pars 86 (W. HORN—Carabidae, Cicindelinae).

E. STRAND—Lepidopterorum Catalogus, pars 33 (H. BURGEFF—Zygaenidae).
