



HOKKAIDO UNIVERSITY

Title	Nitrosophilus kaionis sp. nov., a hydrogen-, sulfur- and thiosulfate-oxidizing chemolithoautotroph within Campylobacteria isolated from a deep-sea hydrothermal vent in the Mid-Okinawa Trough
Author(s)	Fukazawa, So; Mino, Sayaka; Tsuchiya, Jiro et al.
Citation	Archives of Microbiology, 205(1), 12 https://doi.org/10.1007/s00203-022-03350-6
Issue Date	2023-01
Doc URL	https://hdl.handle.net/2115/91292
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: http://dx.doi.org/10.1007/s00203-022-03350-6
Type	journal article
File Information	Nitrosophilus kaionis sp. nov., a hydrogen-, sulfur- and thiosulfate-oxidizing chemolithoautotroph within "Campylobacteria" isolated from a deep-sea hydrothermal vent in the Mid-Okinawa Trough.pdf



***Nitrosophilus kaiyonis* sp. nov., a hydrogen-, sulfur- and thiosulfate-oxidizing chemolithoautotroph within “*Campylobacteria*” isolated from a deep-sea hydrothermal vent in the Mid-Okinawa Trough**

So Fukazawa¹, Sayaka Mino^{1*}, Jiro Tsuchiya¹, Satoshi Nakagawa^{2,3}, Ken Takai³, Tomoo Sawabe¹

¹Laboratory of Microbiology, Faculty of Fisheries Sciences, Hokkaido University, Hakodate, Japan

²Laboratory of Marine Environmental Microbiology, Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University, Kyoto, Japan

³Institute for Extra-cutting-edge Science and Technology Avant-garde Research (X-star), Japan Agency for Marine-Earth Science & Technology (JAMSTEC), Yokosuka, Japan

Running title: *Nitrosophilus kaiyonis* sp. nov.

Key words; deep-sea hydrothermal vents, chemolithoautotroph, thermophile, Mid-Okinawa Trough, *Campylobacteria*

*Correspondence:

Sayaka Mino

Laboratory of Microbiology, Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate 041-8611, Japan

Phone: +81-138-40-5570; Fax: +81-138-40-5570

Email: sayaka.mino@fish.hokudai.ac.jp

Abstract

A novel bacterium, strain MOT50^T, was isolated from the chimney structure at the Iheya North field in the Mid-Okinawa Trough. The cells were motile short rods with single polar flagellum. Growth was observed between 40 and 65°C (optimum, 52°C), at pH values between 5.0 and 7.1 (optimum, pH 6.1) and in the presence of 2.0–4.0% NaCl (optimum, 2.5%). The isolates utilized molecular hydrogen, thiosulfate, or elemental sulfur as the sole electron donor. Thiosulfate, elemental sulfur, nitrate, and molecular oxygen are utilized as the sole electron acceptor. Ammonium is required as a nitrogen source. Thiosulfate, elemental sulfur, sulfate, or sulfite serves as a sulfur source for growth. The G+C content of the genomic DNA was 28.9%. Phylogenetic analysis based on the 16S rRNA gene sequences indicated that strain MOT50^T belonged to the genus *Nitrosophilus* of the class “*Campylobacteria*”, and its closest relative was *Nitrosophilus labii* HRV44^T (97.20%). On the basis of the phylogenetic, phylogenetical and molecular characteristics, it is proposed that the organism represents a novel species within the genus *Nitrosophilus*, *Nitrosophilus kaiyonis* sp. nov. The type strain is MOT50^T (=JCM 39187^T =KCTC 25251^T).

Introduction

The class “*Campylobacteria*” (homotypic synonym *Epsilonproteobacteria* Garrity *et al.*, 2006) is known as a dominant bacterial group in hydrothermal environments that can account for more than 80% of microbial communities in the chimney structures (Nakagawa *et al.*, 2006; Muto *et al.*, 2017). Members of this group play an important ecological role as primary producers in the deep-sea hydrothermal ecosystems (Nakagawa *et al.*, 2006, Muto *et al.*, 2017). After the first cultivation of thermophilic “*Campylobacteria*” from deep-sea hydrothermal vents (Campbell *et al.*, 2001), taxonomically diverse isolates have been successfully obtained with the development of cultivation techniques, and their physiological features have been described (Zhang *et al.*, 2018). In addition, recent *in situ* incubation experiment has revealed genus specific primary production traits among hydrothermal vent *Campylobacteria* (McNichol *et al.*, 2022). Further efforts to isolate and characterize novel campylobacterial strains from deep-sea hydrothermal environments could help to better elucidate the ecophysiological traits of taxa among *Campylobacteria*.

The class “*Campylobacteria*” consists of 225 validly described species, including subspecies, 18 of which are thermophilic species according to the List of Prokaryotic Names with Standing in Nomenclature (LPSN) at the time of writing (21 June 2022). Although thermophilic species have been detected in global deep-sea hydrothermal environments (Nakagawa *et al.*, 2005b; Takai *et al.*, 2004; Voordeckers *et al.*, 2005), 4 out of 6 thermophilic genera (e.g., *Nitratiruptor*, *Nitrosophilus*, *Hydrogenimonas*, and *Lebetimonas*) have been described with two or single species. To better

understand the evolution and physiological diversity of the thermophilic “*Campylobacteria*”, it is necessary to obtain thermophilic “*Campylobacteria*” and investigate their physiological and genomic characteristics. In this study, we report the newly thermophilic campylobacterium, strain MOT50^T, belonging to the genus *Nitrosophilus* based on its physiological and genomic characterization.

Materials and Methods

Sample collection, enrichment, and purification

Strain MOT50^T was isolated from the C0016B chimney structure at the Iheya North field in the Mid-Okinawa Trough (27° 47' 26.9" N, 126° 53' 47.4" E), at a depth of 993 m, by *ROV Hyper Dolphin* during the KY14-01 scientific cruise aboard *R/V Kaiyo* in January 2014. The interior part of the chimney sample was mixed anaerobically with 25 ml sterilized seawater containing 0.05% (w/v) neutralized sodium sulfide in 100 ml glass bottles (Schott Glaswerke, Mainz, Germany) soon after the vehicle was recovered. The bottle was then tightly sealed with a butyl-rubber stopper under a 100% N₂ gas phase (0.2 MPa) and stored at 4°C until use. For enrichment, 100 µl of the resultant slurry was inoculated into 15 ml test tubes containing 3 ml MMJHS (Takai *et al.*, 2003). MMJHS medium contained 1 g NaNO₃, 1 g NaHCO₃, 1 g Na₂S₂O₃·5H₂O, 10 g S⁰ per liter MJ synthetic seawater. The gas phase (0.2 MPa) of MMJHS medium was H₂/CO₂ (80:20, v/v). Growth of thermophiles was observed after one day at 50°C. Strain MOT50^T was isolated using the dilution-to-extinction technique (Takai & Horikoshi, 2000) with MMJHS medium at 50°C. The purity was confirmed with a routine microscopic examination and by repeated partial sequencing of 16S rRNA gene using several PCR primers (Lane, 1991).

Cell morphology and growth characteristics

Cells were observed with the ZEISS Axiophot microscope (Carl Zeiss Co., Oberkochen, Germany). For transmission electron microscope, cells grown in MMJHS medium at 52°C in the late exponential phase were stained with EM Stainer (Nisshin EM Co., Ltd, Tokyo, Japan). Electron micrographs were obtained using JEM-1011 transmission electron microscope (JEOL, Tokyo, Japan).

Growth of the novel isolates was measured by direct cell counts after staining with 4', 6-diamidino-2-phenylindole (DAPI) (Porter & Feig, 1980). To determine optimum temperature, pH and NaCl concentrations, cultures were grown in 3 ml MMJHS medium in various conditions.

The isolate was tested for the ability to grow on combinations of a single electron donor and acceptor. MJ synthetic seawater containing 0.1% (w/v) NaHCO₃ was used as the basal medium. To examine the growth on hydrogen as an electron donor, H₂/CO₂ (80:20) was used as the gas phase. In an attempt to examine the growth on thiosulfate (0.1%, w/v) or elemental sulfur (S⁰) (1%, w/v) as an electron donor, N₂/CO₂ (80/20) was used as the gas phase. Nitrate (0.1%, w/v), thiosulfate (0.1%,

w/v), nitrous oxide (33%, v/v), sulfate (0.1%, w/v), elemental sulfur (1%, w/v) and molecular oxygen (0.1, 1 and 2%, v/v) were tested for potential electron acceptors. The growth was confirmed by measuring the OD₆₂₀ (Microplate Reader, Infinite F200, Tecan, Mannheim, Switzerland) or cell increase via microscopy.

Heterotrophic growth was tested in MMJHS medium without NaHCO₃ under a gas phase of 100% H₂ (0.3% MPa). Each of the following potential carbon sources was tested: yeast extract, peptone, tryptone, casamino acids, D(+)-glucose, galactose, sucrose, fructose, lactose, maltose, starch (all 0.1%, w/v), formate, acetate, glycerol, tartrate, malate, succinate, propionate, lactate, pyruvate (all 10 mM), methanol (0.05%, w/v), ethanol (0.1%, v/v) and 2-propanol (0.2%, v/v). In addition, to assess the utilization of these organic compounds as an energy source, substrates were added to medium excluded thiosulfate (0.1%, w/v) and elemental sulfur (1%, w/v) from MMJHS medium under a N₂/CO₂ (80/20) gas phase (0.3 MPa).

Potential nitrogen and sulfur sources required for growth were tested. To determine the nitrogen sources utilization, NH₄Cl (0.025%, w/v), NaNO₃ (0.1%, w/v) and NaNO₂ (0.1% w/v) were examined in MMJHS medium lacking all nitrogen sources, under a H₂/CO₂ (80:20) gas phase (0.3 MPa). To determine the sulfur sources utilization, sulfate (0.34%, w/v), thiosulfate (0.1%, w/v), sulfite (0.1 and 1%, w/v) and elemental sulfur (1%, w/v) were examined in MMJHS medium in which sulfur compounds were removed and replaced with the chloride salts under an H₂/CO₂ (80/20) gas phase (0.3 MPa).

Molecular phylogenetic analysis based on 16S rRNA gene sequences

The 16S rRNA gene was amplified by PCR using primers Eubac 27F and 1492R (Lane, 1991). The nearly complete 16S rRNA gene sequence (1,420 bp) of strain MOT50^T was obtained by direct sequencing of both strands. The 16S rRNA gene sequence was analyzed using the BLAST search algorithm with all nucleotides (Altshul *et al.*, 1997). In order to determine the phylogenetic position of the strain, the other thermophilic campylobacterial sequences were retrieved from the Silva database (Quast *et al.*, 2013), and the NCBI RefSeq database (<https://www.ncbi.nlm.nih.gov/>) and then aligned using Silva Incremental Aligner v1.2.11 (Pruesse *et al.*, 2012). A phylogenetic tree was constructed using the neighbor-joining method (Saitou *et al.*, 1987) with the MEGAX software (Kumar *et al.*, 2018) using 1,134 homologous sequence positions for each organism. Bootstrap analysis was conducted using 1,000 replications to provide a confidence result for the phylogenetic tree topology.

Genome sequencing and assembly

Genomic DNA of strain MOT50^T was extracted from the cells grown in MMJHS medium with Wizard genomic DNA purification kit (Promega, Madison, Wisconsin, USA) according to the

127 manufacturer's protocol. The genome sequencing was performed using both Oxford Nanopore
 128 Technology (ONT) MinION and Illumina MiSeq platforms. For the ONT sequencing, the library was
 129 prepared using the Rapid Barcoding Sequence kit (Oxford Nanopore Technologies, Oxford, UK)
 130 according to the standard protocol provided by the manufacturer. The library was made by mixing
 131 DNA samples with Fragment Mix RB01-12. The constructed library was loaded into the Flow Cell
 132 (R9.4.1) on a MinION device and performed a 48-hour sequencing run with MinKNOW3.6.0
 133 software. Basecalling for ONT reads was performed with Guppy v3.4.4 (Oxford Nanopore
 134 Technologies). Illumina reads were combined with ONT reads for hybrid assembly with Unicycler
 135 v0.4.7 (Wick *et al.*, 2017), with default parameters. The whole genome sequence was annotated with
 136 DFAST (Tanizawa *et al.*, 2018). Metabolic pathways were confirmed using BlastKOALA (Kanehisa
 137 *et al.*, 2016) and sulfur oxidizing pathway was visualized by KEGG mapper (Kanehisa & Sato, 2020).
 138 In order to survey genes responsible for sulfur reducing metabolisms (*psrABC* and *sud*), homology
 139 search equipped with in silico MolecularCloning (In Silico Biology, Inc., Yokohama, Japan) was
 140 performed with default setting using amino acid sequences (WS0116-0118 and WS1629) from
 141 *Wolinella succinogenes* DSM 1740^T (BX571656.1) as reference sequences (Yamamoto *et al.*, 2010).
 142 To compare the related strains of strain MOT50^T, complete genome sequences of *Nitratiruptor*
 143 *tergarcus* MI55-1^T (Nakagawa *et al.*, 2005b) and *Hydrogenimonas thermophila* EP1-55-1%^T (Takai
 144 *et al.*, 2004) were also obtained in the same way.

145 **Calculation of genome sequence similarities**

146 To determine the taxonomic position of strain MOT50^T, genome-based taxonomic indices
 147 were calculated with related strains. *In silico* DNA-DNA hybridization (DDH) values of strain
 148 MOT50^T against *Nitrosophilus alvini* EPR55-1^T (Shiotani *et al.*, 2020), *Nitrosophilus labii* HRV44^T
 149 (Fukushi *et al.*, 2020), *Nitratiruptor tergarcus* MI55-1^T (Nakagawa *et al.*, 2005b), *Nitratiruptor* sp.
 150 SB155-2 (Nakagawa *et al.*, 2007) and *Hydrogenimonas thermophila* EP1-55-1%^T (Takai *et al.*, 2004)
 151 were calculated using the Genome-to-Genome Distance Calculator version (Meier-Kolthoff *et al.*,
 152 2013) with the BLAST+ alignment tool. Results were based on recommended formula2, which is
 153 independent of genome length and is robust against using draft genomes. Average nucleotide identity
 154 (ANI) and average amino acid identity (AAI) values were calculated by using the EZ BioCloud (Yoon
 155 *et al.*, 2017) and the aai.rb script (<https://github.com/lmrodriguezz/enveomics>) (Rodriguez-R &
 156 Konstantinidis, 2016), respectively. The genome sequences of *Nitrosophilus alvini* EPR55-1^T,
 157 *Nitrosophilus labii* HRV44^T and *Nitratiruptor* sp. SB155-2, were retrieved from NCBI RefSeq
 158 database (<https://www.ncbi.nlm.nih.gov/>).

159 In order to determine the phylogenetic position of the strain, the other campylobacterial
 160 sequences were retrieved from the NCBI RefSeq database (<https://www.ncbi.nlm.nih.gov/>) and

constructed a phylogenetic tree based on Campbell's 139 single-copy genes (Campbell *et al.*, 2013) using the Anvi'o v5.5.0 (Eren *et al.*, 2015). The workflow of this analysis was based on the workflow (<http://merenlab.org/2017/06/07/phylogenomics/>).

Pan-genomic analysis

Pan-genomic analysis was conducted with 22 deep-sea campylobacterial genomes using the Anvi'o v5.5.0 (Eren *et al.*, 2015). The workflow of this analysis was based on the workflow (<http://merenlab.org/2017/06/07/phylogenomics/>). COG-annotated species-specific gene clusters were exported from pangenome (anvi-summarize) for the gene functional analysis. Genomes used in this study is shown in supplementary Table S1.

Nucleotide sequence accession number

The nucleotide sequence of the 16S rRNA gene and the complete genome sequence of strain MOT50^T have been deposited in the GenBank/ EMBL/DDBJ under accession numbers LC716068 and AP025696, respectively. Complete genome sequences of *Nitratiruptor tergarcus* MI55-1^T and *Hydrogenimonas thermophila* EP1-55-1%^T have been deposited in the GenBank/ EMBL/DDBJ under accession numbers AP026671-AP026672 and AP026673-AP026676, respectively.

Results

Cell morphology and growth characteristics

Cells of MOT50^T were Gram-negative rods (2.0 µm long and 1.0 µm in wide) (Fig. 1). The cells were motile by means of the polar flagellum. Spore formation was not observed under any laboratory conditions.

Strain MOT50^T grew at temperatures between 40 and 65°C, showing optimum growth at 52°C (doubling time, 2 h). No growth was observed below 35°C or above 70°C. Growth occurred between pH 5.0 and 7.1, with optimum growth at pH 6.1. No growth was observed below pH 4.6 or above 7.8. The isolate grew at concentrations in the range 2.0–4.0% (w/v) NaCl, with optimum growth at 2.5% NaCl. No growth was observed in concentrations below 1.5% NaCl or above 5.0% NaCl (Supplementary Fig. S1). Growth characteristics of strain MOT50^T were similar to those of *Nitrosophilus labii* HRV44^T (Table 1).

Strain MOT50^T was able to utilize thiosulfate (0.1%, w/v), elemental sulfur (1%, w/v), or H₂ as the sole electron donor. Thiosulfate (0.1%, w/v), elemental sulfur (1%, w/v), nitrate (0.1%, w/v), nitrous oxide (33%, v/v) or molecular oxygen (0.1%, 1% and 2%, v/v) could serve as the sole electron acceptor. N₂O consumption was observed only when H₂ was provided as the electron donor. The combination of H₂ and nitrate resulted in the highest cell growth. The isolate was unable to use any organic compounds examined in this study as energy or carbon sources. MOT50^T was able to utilize

195 ammonium (0.025%, w/v) and nitrate (0.1%, w/v) as a nitrogen source. The isolate was able to utilize
196 thiosulfate (0.1%, w/v) and elemental sulfur (1%, w/v) as a sulfur source.

197 **Phylogenetic analysis based on 16S rRNA gene sequences**

198 The 16S rRNA gene sequence of MOT50^T shared the highest similarity with that of
199 *Nitrosophilus labii* HRV44^T (97.20%), followed by that of *Nitrosophilus alvini* EPR55-1^T (95.07%).
200 These values were below the threshold for species definition (98.7%) based on 16S rRNA gene
201 sequences (Yarza *et al.*, 2014). The phylogenetic analysis also showed that the isolate was a member
202 of the genus *Nitrosophilus* (Fig. 2).

203 **Genome properties**

204 A single complete circular contig of MOT50^T with a length of 1,769,550 bp was obtained by
205 Unicycler, and 1,798 contig sequences (CDSs) were predicted (Supplementary Fig. S2). This genome
206 size was the smallest among previously sequenced genomes of *Nitrosophilus* spp., i.e., *Nitrosophilus*
207 *alvini* EPR55-1^T (1,807,889 bp and 1,797 CDSs) (Shiotani *et al.*, 2020), and *Nitrosophilus labii*
208 HRV44^T (chromosome; 1,990,315 bp and 2,050 CDSs, plasmid; 102,672 bp and 128 CDSs) (Fukushi
209 *et al.*, 2020). The G+C content of MOT50^T genome was 28.9%, which is similar to that of
210 *Nitrosophilus labii* HRV44^T (33.4%) (Table 1). MOT50^T had a complete set of Sox genes
211 (*soxABXYZ*) responsible for sulfur oxidation. Strain MOT50^T had homologues of *psrABC* encoding
212 polysulfide reductase and of *sud* encoding sulfide dehydrogenase (Klimmek *et al.*, 1998).

214 **Taxonomic placement of MOT50^T based on the genomic analyses**

215 *In silico* DDH values of strain MOT50^T against *Nitrosophilus alvini* EPR55-1^T,
216 *Nitrosophilus labii* HRV44^T, *Nitratiruptor tergarcus* MI55-1^T, *Nitratiruptor* sp. SB155-2 and *H.*
217 *thermophila* EP1-55-1%^T were 14.5%, 20.0%, 19.8%, 19.7%, and 17.6%, respectively, which were
218 below the threshold of bacterial species boundary (Meier-Kolthoff *et al.*, 2013). In addition, ANI
219 values of strain MOT50^T against *Nitrosophilus alvini* EPR55-1^T, *Nitrosophilus labii* HRV44^T,
220 *Nitratiruptor tergarcus* MI55-1^T, *Nitratiruptor* sp. SB155-2 and *H. thermophila* EP1-55-1%^T were
221 71.6%, 75.9%, 72.8%, 72.3% and 70.3%, respectively. These values are also below a threshold of
222 95.0% ANI similarity for the definition of bacterial species (Richter & Rosselló-Mora, 2009). These
223 results also support the proposal that strain MOT50^T is a new species within the class
224 “*Campylobacteria*”. AAI values of the novel isolate against *Nitrosophilus alvini* EPR55-1^T,
225 *Nitrosophilus labii* HRV44^T, *Nitratiruptor tergarcus* MI55-1^T, *Nitratiruptor* sp. SB155-2 and *H.*
226 *thermophila* EP1-55-1%^T were 69.9%, 75.7%, 68.7%, 67.7% and 58.8%, respectively. The values
227 against *Nitrosophilus* and *Nitratiruptor* species were above the threshold of the genus-level boundary
228 of “*Campylobacteria*” (60–62%) (Shiotani *et al.*, 2020). The phylogenetic tree based on nucleotide

sequences of Campbell's 139 SCGs showed that the isolate was a member of the genus *Nitrosophilus* (Fig. 3).

Pan-genomic analysis

The pangenome of members of “*Campylobacter*” isolated from deep-sea hydrothermal environments revealed a total of 13,859 gene clusters comprising 46,679 genes in which 424 gene clusters with 9,328 genes were represented SCGs, and 38 gene clusters with 943 genes were core genome (Fig. 4). The species-unique genes were also found among the genus *Nitrosophilus*; *N. labii* HRV44^T had approximately 1.6 times higher number of unique gene clusters (337 gene clusters with 367 genes) than other two species, *N. kaiyonis* MOT50^T (199 gene clusters with 201 genes) and *N. alvini* EPR55-1^T (197 gene clusters with 201 genes). COG functional annotation of species-unique gene clusters among the genus *Nitrosophilus* showed that a relative abundance of genes involved in cell wall/membrane/envelope biogenesis (M) in strain MOT50^T was higher (18.8% in without uncharacterized categories) than those in *N. labii* HRV44^T (6.4%) and *N. alvini* EPR55-1^T (10.1%) (Supplementary Fig. S3). Genes related to defense mechanisms (V) (9.0%), mobilome: prophages, transposons (X) (19.4%), and replication, recombination and repair (L) (28.6%) in strain HRV44^T were higher than those in strain MOT50^T (2.4%, 9.8%, and 13.8%, respectively) and strain EPR55-1^T (3.4%, 0%, and 17.9%, respectively).

Discussion

Although more than 20 years have passed since the first report on thermophilic members of “*Campylobacter*”, novel thermophilic campylobacterial species, i.e., *Lebetimonas natsushimae* (Nagata *et al.*, 2017), *Nitrosophilus labii* (Fukushi *et al.*, 2020), *Nitrosophilus pacificus* (Shiotani *et al.*, 2020), *Hydrogenimons urashimensis* (Mino *et al.*, 2021), still have been isolated from deep-sea hydrothermal environments, and their characterizations update physiological traits of these genera. For example, *L. natsushimae* is able to utilize formate as its electron donor, that trait has not been observed in other *Lebetimonas* species. *Nitrosophilus labii* is the first described species in “*Campylobacter*” that can reduce exogenous N₂O to N₂. Continued efforts to isolate and cultivate novel species are great worth not for elucidating microbial ecological roles in the deep-sea hydrothermal vent ecosystem but also for finding microbial resources that could contribute to application studies. In this study, we successfully isolated a novel thermophilic strain, MOT50^T, from a chimney structure collected at the Iheya North field in the Mid-Okinawa Trough, and characterized physiological and genomic features of the strain, and propose it as a novel species belonging to the genus *Nitrosophilus*.

Physiological characterization and advanced polyphasic taxonomy including genome comparison revealed strain MOT50^T is a novel bacterial species of the genus *Nitrosophilus*. Temperature, pH, and NaCl concentration ranges for growth of strain MOT50^T are similar to those of *Nitrosophilus labii* HRV44^T, whereas energy metabolic traits of strain MOT50^T are similar to those of *Nitrosophilus alvini* EPR55-1^T. Strain MOT50^T is able to utilize elemental sulfur and thiosulfate as both electron donors and acceptors. This energy metabolic trait has never been reported in any other thermophilic “*Campylobacteria*”. Key genes responsible for sulfur oxidation and reduction were found in the genome of MOT50^T, supporting the ability to utilize reduced sulfur compound as both electron donors and acceptors. In comparison with other species within the genus *Nitrosophilus*, strain MOT50^T is able to utilize a relatively wide variety of inorganic compounds (i.e., molecular hydrogen, thiosulfate, or elemental sulfur) as the sole energy source. Although we did not evaluate which combination of substances strain MOT50^T prefers when those are present at the same time, this characteristic might help strain MOT50^T to adapt to fluctuating environmental conditions surrounding deep-sea hydrothermal vents.

Species-specific differences in gene content and its function commonly related to a specific metabolism, virulence, antibiotic resistance mechanisms, or other environmental adaptation (Costa *et al.*, 2020). COG functional annotation of gene clusters unique to each *Nitrosophilus* species showed that genes involved in the category M (cell wall/membrane/envelope biogenesis) accounted for higher proportion in strain MOT50^T than in strain HRV44^T or strain EPR55-1^T, suggesting an advantage role against physical and chemical stresses (Jiang *et al.*, 2019) for strain MOT50^T. Also, genes involved in the categories V (defense mechanisms), X (mobilome: prophages, transposons), and L (replication, recombination and repair) in strain HRV44^T were more abundant than those in strain MOT50^T and strain EPR55-1^T, suggesting that gene exchange and recombination are important in strain HRV44^T and these functions are responsible for strategies in response to the environment factors (Jiang *et al.*, 2019).

287

288 **Description of *Nitrosophilus kaiyonis* sp. nov.**

289 *Nitrosophilus kaiyonis* (ka.i.yo'nis. N.L. gen. n. *kaiyonis*, of *Kaiyo*, the name of the Japanese
290 retired research ship used for the exploration of deep ocean).

291 Cells are Gram-negative, motile and short-rod shape and a mean length of 2.0 μm and width
292 of 1.0 μm. The temperature range for the growth is 40–65°C (optimum, 52°C; doubling time, 2h). The
293 pH range for the growth is 5.0–7.1 (optimum, pH 6.1). NaCl in the concentration range for the growth
294 is 2.0–4.0% (optimum, 2.5%). Strictly chemolithoautotrophic growth occurs with molecular
295 hydrogen, thiosulfate, or elemental sulfur as the sole electron donor. Thiosulfate, elemental sulfur,

296 nitrate, and molecular oxygen are utilized as the sole electron acceptor. Ammonium is required as a
297 nitrogen source. Thiosulfate, elemental sulfur, sulfate, or sulfite serves as a sulfur source for growth.
298 The size of the complete genome is 1,769,550 bp. The GC content of DNA is 28.9%. The type strain,
299 MOT50^T (=JCM 39187^T =KCTC 25251^T) was isolated from a deep-sea hydrothermal vent in the
300 Mid-Okinawa Trough.

301

302 **Acknowledgements**

303 We gratefully acknowledge the R/V *Kaiyo* crew and the *ROV Hyper Dolphin* operation team
304 for helping us to collect samples from deep-sea hydrothermal vents in the Mid-Okinawa Trough. This
305 work was partially supported by JSPS KAKENHI Grant Numbers JP 17K15301 and 21K14913.

306

References

- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA et al (2008) The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 9: 75. <https://doi.org/10.1186/1471-2164-9-75>
- Campbell BJ, Jeanthon C, Kostka JE, Luther GW 3rd, Cary SC (2001) Growth and phylogenetic properties of novel bacteria belonging to the Epsilon subdivision of the *Proteobacteria* enriched from *Alvinella pompejana* and deep-sea hydrothermal vents. *Appl Environ Microbiol* 67: 4566–4572. <https://doi.org/10.1128/aem.67.10.4566-4572.2001>
- Campbell JH, O'Donoghue P, Campbell AG, Schwientek P, Sczyrba A, Woyke T et al (2013) UGA is an additional glycine codon in uncultured SR1 bacteria from the human microbiota. *Proc Natl Acad Sci U S A* 110:5540-5545. <https://doi.org/10.1073/pnas.1303090110>
- Costa SS, Guimarães LC, Silva A, Soares SC, Baraúna RA (2020) First steps in the analysis of prokaryotic pan-genomes. *Bioinform Biol Insights* 14:1177932220938064. <https://doi.org/10.1177/1177932220938064>
- Eren MA, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML et al (2015) Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* 3: e1319. <https://doi.org/10.7717/peerj.1319>
- Fukushi M, Mino S, Tanaka H, Nakagawa S, Tanaka K, Sawabe T (2020) Biogeochemical implications of N₂O-reducing thermophilic “*Campylobacteria*” in deep-sea vent fields, and the description of *Nitratiruptor labii* sp. nov. *iScience* 23: 101462. <https://doi.org/10.1016/j.isci.2020.101462>
- Grosche A, Sekaran H, Pérez-Rodríguez I, Starovoytov V, Vetriani C (2015) *Cetia pacifica* gen. nov., sp. nov., a chemolithoautotrophic, thermophilic, nitrate-ammonifying bacterium from a deep-sea hydrothermal vent. *Int J Syst Evol Microbiol* 65: 1144–1150. <https://doi.org/10.1099/ij.s.0.000070>
- Inagaki F, Takai K, Nealson KH, Horikoshi K (2003) *Sulfurimonas autotrophica* gen. nov., sp. nov., a sulfur and thiosulfate oxidizing ϵ -proteobacterium isolated from the Okinawa Trough. *Int J Syst Evol Microbiol* 53: 1801–1805. <https://doi.org/10.1099/ij.s.0.02682-0>
- Inagaki F, Takai K, Nealson KH, Horikoshi K (2004) *Sulfurovum lithotrophicum* gen. nov., sp. nov., a novel sulfur-oxidizing chemolithoautotroph within the ϵ -*Proteobacteria* isolated from Okinawa Trough hydrothermal sediments. *Int J Syst Evol Microbiol* 54: 1477–1482. <https://doi.org/10.1099/ij.s.0.03042-0>
- Jiang X, Li X, Yang L, Liu C, Wang Q, Chi W et al (2019) How microbes shape their communities? a microbial community model based on functional genes. *Genom Proteom Bioinform* 17: 91-105. <https://doi.org/10.1016/j.gpb.2018.09.003>

350
351 Kanehisa M, Sato Y, Morishima K (2016) BlastKOALA and GhostKOALA: KEGG tools for
352 functional characterization of genome and metagenome sequences. J Mol Biol 428: 726–731.
353 <https://doi.org/10.1016/j.jmb.2015.11.006>
354
355 Kanehisa M, Sato Y (2020) KEGG Mapper for inferring cellular functions from protein sequences
356 Protein Sci 29: 28-35. <https://doi.org/10.1002/pro.3711>
357
358 Klimmek O, Kreis V, Klein C, Simon J, Wittershagen A, Kroger A (1998) The function of the
359 periplasmic Sud protein in polysulfide respiration of *Wolinella succinogenes*. Eur J Biochem 253:
360 263–269. <https://doi.org/10.1046/j.1432-1327.1998.2530263.x>
361
362 Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: molecular evolutionary genetics
363 analysis across computing platforms. Mol Biol Evol 35:1547–1549.
364 <https://doi.org/10.1093/molbev/msy096>
365
366 Lane DJ. 16S/23S rRNA sequencing. In: Stackebrandt, E. and Goodfellow, M., Eds., Nucleic Acid
367 Techniques in Bacterial Systematic, John Wiley and Sons, New York. 1991. 115–175.
368
369 McNichol J, Dykema S, Mußmann M, Seewald JS, Sylva SP, Sievert SM (2022) Genus-specific
370 carbon fixation activity measurements reveal distinct responses to oxygen among hydrothermal vent
371 *Campylobacteria*. Appl Environ Microbiol 88:e0208321. <https://doi.org/10.1128/aem.02083-21>
372
373 Meier-Kolthoff JP, Auch AF, Klenk HP, Goëker M (2013) Genome sequence-based species
374 delimitation with confidence intervals and improved distance functions. BMC Bioinform 14: 60.
375 <https://doi.org/10.1186/1471-2105-14-60>
376
377 Mino S, Shiotani T, Nakagawa S, Takai K, Sawabe T (2021) *Hydrogenimonas urashimensis* sp. nov.,
378 a hydrogen-oxidizing chemolithoautotroph isolated from a deep-sea hydrothermal vent in the
379 Southern Mariana Trough. Syst Appl Microbiol 44: 126170.
380 <https://doi.org/10.1016/j.syapm.2020.126170>
381
382 Muto H, Takaki Y, Hirai M, Mino S, Sawayama S, Takai K et al (2017) S. A simple and efficient RNA
383 extraction method from deep-sea hydrothermal vent chimney structures. Microbes Environ 32: 330–
384 335. <https://doi.org/10.1264/jsme2.me17048>
385
386 Nagata R, Takaki Y, Tame A, Nunoura T, Muto H, Mino S et al (2017) *Lebetimonas natsushimae* sp.
387 nov., a novel strictly anaerobic, moderately thermophilic chemoautotroph isolated from a deep-sea
388 hydrothermal vent polychaete nest in the Mid-Okinawa Trough. Syst Appl Microbiol 40: 352–356.
389 <https://doi.org/10.1016/j.syapm.2017.06.002>
390
391 Nakagawa S, Takai K, Inagaki F, Hirayama H, Nunoura T, Horikoshi K et al (2005a) Distribution,
392 phylogenetic diversity and physiological characteristics of epsilon-*Proteobacteria* in a deep-sea
393 hydrothermal field. Appl Environ Microbiol 71: 1619–1632. [12](https://doi.org/10.1111/j.1462-</p>
</div>
<div data-bbox=)

2920.2005.00856.x

Nakagawa S, Takai K, Inagaki F, Horikoshi K, Sako Y (2005b) *Nitratiruptor tergarcus* gen. nov., sp. nov. and *Nitratifactor salsuginis* gen. nov., sp. nov., nitrate-reducing chemolithoautotrophs of the ϵ -*Proteobacteria* isolated from a deep-sea hydrothermal system in the Mid-Okinawa Trough. *Int J Syst Evol Microbiol* 55: 925–933. <https://doi.org/10.1099/ijs.0.63480-0>

Nakagawa S, Takaki Y, Shimamura S, Reysenbach AL, Takai K, Horikoshi K (2007) Deep-sea vent ϵ -proteobacterial genomes provide insights into emergence of pathogens. *Proc Natl Acad Sci U S A* 104: 12146–12150. <https://doi.org/10.1073/pnas.0700687104>

Nakagawa T, Takai K, Suzuki Y, Hirayama H, Konno U, Tsunogai U et al (2006) Geomicrobiological exploration and characterization of a novel deep-sea hydrothermal system at the TOTO caldera in the Mariana Volcanic Arc. *Appl Environ Microbiol* 8: 37–49. <https://doi.org/10.1128/aem.70.8.4648-4657.2004>

Porter K, Feig Y (1980) The use of DAPI for identifying and counting aquatic microflora. *Limnol Oceanogr* 25: 943–948. <https://doi.org/10.4319/lo.1980.25.5.0943>

Pruesse E, Peplies J, Glöckner FO (2012) SINA: accurate high-throughput multiple sequence alignment of ribosomal RNA genes. *Bioinformatics* 28: 1823–1829. <https://doi.org/10.1093/bioinformatics/bts252>

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P et al (2013) The SILVA ribosomal RNA gene 64 database project: improved data processing and web-based tools. *Nucleic Acids Res* 41: D590–D596. <https://doi.org/10.1093/nar/gks1219>

Richter M, Rosselló-Mora R (2009) Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* 106: 19126–19131. <https://doi.org/10.1073/pnas.0906412106>

Rodriguez-R LM, Konstantinidis KT (2016) The enveomics collection: a toolbox for specialized analyses of microbial genomes and metagenomes. *PeerJ Preprints* 4: e1900v1. <https://doi.org/10.7287/peerj.preprints.1900v1>

Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4: 406–425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>

Shiotani T, Mino S, Sato W, Nishikawa S, Yonezawa M, Sievert MS, Sawabe T (2020) *Nitrosophilus alvini* gen. nov., sp. nov., a hydrogen-oxidizing chemolithoautotroph isolated from a deep-sea hydrothermal vent in the East Pacific Rise, inferred by a genome-based taxonomy of the phylum “*Campylobacterota*”. *PLoS ONE* 15: e0241366. <https://doi.org/10.1371/journal.pone.0241366>

Smith JL, Campbell BJ, Hanson TE, Zhang CL, Cary SC (2008) *Nautilia profundicola* sp. nov., a thermophilic, sulfur-reducing epsilonproteobacterium from deep-sea hydrothermal vents. *Int J Syst*

438 Evol Microbiol 58: 1598–1602. <https://doi.org/10.1099/ijs.0.65435-0>

439

440 Takai K, Horikoshi K (2000) *Thermosipho japonicus* sp. nov., an extremely thermophilic bacterium

441 isolated from a deep-sea hydrothermal vent in Japan. Extremophiles 4: 9–17.

442 <https://doi.org/10.1007/s007920050002>

443

444 Takai K, Inagaki F, Nakagawa S, Hirayama H, Nunoura T, Sako Y et al (2003) Isolation and

445 phylogenetic diversity of members of previously uncultivated ϵ -*Proteobacteria* in deep-sea

446 hydrothermal fields. FEMS Microbiol Lett 218: 167–174. [https://doi.org/10.1111/j.1574-](https://doi.org/10.1111/j.1574-6968.2003.tb11514.x)

447 [6968.2003.tb11514.x](https://doi.org/10.1111/j.1574-6968.2003.tb11514.x)

448

449 Takai K, Nealson KH, Horikoshi K (2004) *Hydrogenimonas thermophila* gen. nov., sp. nov., a novel

450 thermophilic, hydrogen-oxidizing chemolithoautotroph within the ϵ -*Proteobacteria*, isolated from a

451 black smoker in a Central Indian Ridge hydrothermal field. Int J Syst Evol Microbiol 54: 25–32.

452 <https://doi.org/10.1099/ijs.0.02787-0>

453

454 Tanizawa Y, Fujisawa T, Nakamura Y (2018) DFAST: a flexible prokaryotic genome annotation

455 pipeline for faster genome publication. Bioinformatics 34: 1037–1039.

456 <https://doi.org/10.1093/bioinformatics/btx713>

457

458 Voordeckers JW, Starovoytov V, Vetriani C (2005) *Caminibacter mediatlanticus* sp. nov., a

459 thermophilic, chemolithoautotrophic, nitrate-ammonifying bacterium isolated from a deep-sea

460 hydrothermal vent on the Mid-Atlantic Ridge. Int J Syst Evol Microbiol 55: 773–779.

461 <https://doi.org/10.1099/ijs.0.63430-0>

462

463 Wick RR, Judd LM, Gorrie CL, Holt KE (2017) Unicycler: resolving bacterial genome assemblies

464 from short and long sequencing reads. PLoS Comput Biol 13: e1005595.

465 <https://doi.org/10.1371/journal.pcbi.1005595>

466

467 Yamamoto M, Nakagawa S, Shimamura S, Takai K, Horikoshi K (2010) Molecular characterization

468 of inorganic sulfur-compound metabolism in the deep-sea epsilonproteobacterium *Sulfurovum* sp.

469 NBC37-1. Environ Microbiol 12:1144–1153. <https://doi.org/10.1111/j.1462-2920.2010.02155.x>

470

471 Yamamoto M, Takai K (2011) Sulfur metabolisms in epsilon- and gamma-Proteobacteria in deep-sea

472 hydrothermal fields. Front in Microbiol 192. <https://doi.org/10.3389/fmicb.2011.00192>

473

474 Yarza P, Yilmaz P, Pruesse E, Glöckner FQ, Ludwig W, Schleifer KH et al (2014) Uniting the

475 classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. Nat

476 Rev Microbiol 12: 635–645. <https://doi.org/10.1038/nrmicro3330>

477

478 Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H et al (2017) Introducing EzBioCloud: a

479 taxonomically united database of 16S rRNA and whole genome assemblies. Int J Syst Evol Microbiol

480 67:1613–1617. <https://doi.org/10.1099/ijsem.0.001755>

481

482 Zhang Z, Wua Y, Zhang X (2018) Cultivation of microbes from the deep-sea environments. Deep
483 Sea Res Part II 155: 34–43. <https://doi.org/10.1016/j.dsr2.2017.07.008>

484 **Figure Legends**

485 **Fig. 1** Electron micrograph of a negatively stained cell of strain MOT50^T. Scale bar represents 1.0
486 μm .

487

488 **Fig. 2** Phylogenetic tree based on 16S rRNA gene sequences. Phylogenetic tree of the members of
489 “*Campylobacteria*” isolated from deep-sea hydrothermal environments by the neighbor-joining
490 method using 1,134 homologous sequence positions with TN93+G+I model. Numbers at branches
491 are bootstrap values (%) based on 1,000 replications.

492

493 **Fig. 3** Phylogenetic tree of the members of “*Campylobacteria*” isolated from deep-sea hydrothermal
494 environments based on nucleotide sequences of Campbell’s 139 SCGs.

495

496 **Fig. 4** The pan-genomic analysis of members of “*Campylobacteria*” isolated from deep-sea
497 hydrothermal environments. Bars represent the occurrence of gene clusters in a given genome. The
498 heatmap matrix in the upper right corner indicates the average nucleotide identity (ANI) over 70%.
499 The tree above the heatmap was constructed based on the presence/absence of gene clusters for each
500 strain. Genomes are ordered as layers using a tree based on ANI values matrix. The species names of
501 thermophiles and mesophiles are distinguished by red and black colors, respectively.

502

503 **Table 1.** Comparison of physiological characteristics of MOT50^T with species of “*Campylobacteria*” isolated from deep-sea hydrothermal environments

Characteristics	1	2	3	4	5	6	7	8	9	10	11
Origin	Mid-Okinawa Trough	East Pacific Rise	Mid-Okinawa Trough	Mid-Okinawa Trough	Mid-Okinawa Trough	Central Indian Ridge	East Pacific Rise	Mid-Atlantic Ridge	Mid-Okinawa Trough	Mid-Okinawa Trough	Mid-Okinawa Trough
Temperature range (°C)	40-65	50-60	45-60	40-55	37-65	35-65	30-55	45-70	28-40	10-40	10-40
Temperature optimum (°C)	52	60	53	55	55	55	40	55	37	28-30	25
pH range	5.0-7.1	5.4-8.6	5.4-6.4	5.4-6.9	ND	4.9-7.2	6.0-9.0	4.5-7.5	5.6-7.6	5.0-9.0	4.5-9.0
pH optimum	6.1	6.6	6.0	6.4	ND	5.9	7.0	5.5	7.0	6.5-7.0	6.5
NaCl range (% w/v)	2.0-4.0	2.4-3.2	2.0-4.0	1.5-4.0	ND	1.6-5.6	2.0-5.0	1.0-4.0	1.5-3.5	1.0-6.0	1.6-6.0
NaCl optimum (% w/v)	2.5	2.4	2.5	2.5	ND	3.2	3.0	3.0	3.0	4.0	4.0
Electron donor	H ₂ , S ₂ O ₃ ²⁻ , S ⁰	H ₂	H ₂	H ₂	H ₂ , S ⁰ , S ₂ O ₃ ²⁻	H ₂	H ₂ , formate	H ₂	H ₂	S ₂ O ₃ ²⁻ , S ⁰	S ₂ O ₃ ²⁻ , S ⁰ , S ²⁻
Electron acceptor	NO ₃ ⁻ , S ⁰ , S ₂ O ₃ ²⁻ , O ₂ (up to 1.0%), N ₂ O	NO ₃ , S ₂ O ₃ ²⁻ , O ₂ , S ⁰ , N ₂ O	NO ₃ ⁻ , N ₂ O, S ⁰ , O ₂ (up to 1.0%)	NO ₃ ⁻ , S ⁰ , O ₂ (up to 0.7%)	NO ₃ ⁻ , O ₂	NO ₃ ⁻ , O ₂ , S ⁰	S ⁰	NO ₃ ⁻ , S ⁰	NO ₃ ⁻ , O ₂	NO ₃ ⁻ , O ₂	O ₂
Carbon sources other than CO ₂	-	-	-	-	ND	-	formate	-	-	-	-
Nitrogen sources	NH ₄ ⁺	NH ₄ ⁺	NH ₄ ⁺ , NO ₃ ⁻	NH ₄ ⁺ , NO ₃ ⁻	ND	NH ₄ ⁺ , NO ₃ ⁻	NO ₃ ⁻ , NH ₄ ⁺ , peptone, yeast extract	ND	NH ₄ ⁺ , NO ₃ ⁻	NH ₄ ⁺	ND
Sulfur sources	S ₂ O ₃ ²⁻ , S ⁰ , SO ₄ ²⁻	S ₂ O ₃ ²⁻ , S ⁰ , SO ₃ ²⁻	S ⁰ , SO ₄ ²⁻	S ⁰	ND	ND	ND	ND	S ⁰	ND	ND
DNA G+C content (mol%)	28.9	37.7	33.4	36.9	39.7	33.6	33.5	25.6	35.5	48.0	35.2

504 ND, not determined; -, negative.

505 1, Strain MOT50^T; 2, *Nitrosophilus alvini* EPR55-1^T (Shiotani *et al*, 2020); 3, *Nitrosophilus labii* HRV44^T (Fukushi *et al*, 2020); 4, *Nitratiruptor tergarcus*
506 MI55-1^T (Nakagawa *et al.*, 2005b); 5, *Nitratiruptor* sp. SB155-2 (Nakagawa *et al.*, 2007); 6, *Hydrogenimonas thermophila* EP1-55-1%^T (Takai *et al.*,
507 2004); 7, *Nautilia profundicola* AmH^T (Smith *et al.*, 2008); 8, *Caminibacter mediatlanticus* TB-2^T (Voordeckers *et al.*, 2005); 9, *Nitratifractor salsuginis*
508 E9I37-1^T (Nakagawa *et al.*, 2005b); 10, *Sulfurovum lithotrophicum* 42BKT^T (Inagaki *et al.*, 2004); 11, *Sulfurimonas autotrophica* OK10^T (Inagaki *et al.*,
509 2003).

Supplementary Information

***Nitrosophilus kaiyonis* sp. nov., a hydrogen-, sulfur- and thiosulfate-oxidizing chemolithoautotroph within “*Campylobacteria*” isolated from a deep-sea hydrothermal vent in the Mid-Okinawa Trough**

So Fukazawa¹, Sayaka Mino^{1*}, Jiro Tsuchiya¹, Satoshi Nakagawa^{2,3}, Ken Takai³, Tomoo Sawabe¹

¹Laboratory of Microbiology, Faculty of Fisheries Sciences, Hokkaido University, Hakodate, Japan

²Laboratory of Marine Environmental Microbiology, Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University, Kyoto, Japan


³Institute for Extra-cutting-edge Science and Technology Avant-garde Research (X-star), Japan Agency for Marine-Earth Science & Technology (JAMSTEC), Yokosuka, Japan

Figure legends


Supplementary Fig. S1. The effect of temperature (a), pH (b), and NaCl concentration (c) on the growth of strain MOT50^T.

Supplementary Fig. S2. Graphical circular map of strain MOT50^T genome. Tracks from inside to outside indicate G+C skew, G+C content, rRNA, reverse strand CDS, and forward strand CDS, respectively.


Supplementary Fig. S3. Functional categories of species-specific gene clusters in *Nitrosophilus* species. Category abbreviations are as follows: C, energy production and conversion; E, amino acid transport and metabolism; F, nucleotide transport and metabolism; G, carbohydrate transport and metabolism; H, coenzyme transport and metabolism; I, lipid transport and metabolism; P, inorganic ion transport and metabolism; Q, secondary metabolites biosynthesis, transport and catabolism; X, mobilome: prophages, transposons; A, RNA processing and modification; B, chromatin structure and dynamics; J, translation, ribosomal structure and biogenesis; K, transcription; L, replication, recombination and repair; D, cell cycle control, cell division, chromosomal partitioning; M, cell wall/membrane/envelope biogenesis; N, cell motility; O, posttranslational modification, protein turnover, chaperones; T, signal transduction mechanisms; U, intracellular trafficking, secretion, and vesicular transport; V, defense mechanisms; W, extracellular structures; Z, cytoskeleton; R, general function predicted only; S, function unknown.



Supplementary Fig. S1.



Supplmentary Fig. S2.



Supplementary Fig. S3.

Supplementary Table S1. List of genomes used in this study

Genus	Species	Strain	Assembly level	# of contigs	# of chromosomes and plasmids	Genome size (bp)	G+C content (%)	Accession number
<i>Nitrosophilus</i>	<i>kaiyonis</i>	MOT50	Complete genome	1	1	1,769,550	28.9	AP025696
<i>Nitrosophilus</i>	<i>labii</i>	HRV44	Complete genome	2	2	2,092,987	33.4	GCF_014466985.1
<i>Nitrosophilus</i>	<i>alvini</i>	EPR55-1	Complete genome	1	1	1,807,889	37.7	GCF_015100395.1
<i>Nitratiruptor</i>	<i>tergarcus</i>	DSM 16512	Complete genome	2	2	1,940,884	36.9	AP026671-AP026672
<i>Nitratiruptor</i>	sp.	SB155-2	Complete genome	1	1	1,877,931	39.7	GCF_000010325.1
<i>Hydrogenimonas</i>	<i>thermophila</i>	EP1-55-1%	Complete genome	4	4*	2,551,102	33.6	AP026673-AP026676
<i>Hydrogenimonas</i>	<i>urashimensis</i>	SSM-sur55	Complete genome	2	2	2,286,687	52.8	GCF_016593255.1
<i>Lebetimonas</i>	<i>natsushimae</i>	HS1857	Contig	8	-	1,639,064	30.4	GCF_002335445.1
<i>Caminibacter</i>	<i>pacificus</i>	TB6	Complete genome	2	2	1,871,566	34.0	GCF_005083985.2
<i>Caminibacter</i>	<i>mediatlanticus</i>	TB-2	Complete genome	1	1	1,685,887	25.6	GCF_005843985.1
<i>Nautilia</i>	<i>profundicola</i>	AmH	Complete genome	1	1	1,676,444	33.5	GCF_000021725.1
<i>Nitratifractor</i>	<i>salsuginis</i>	DSM 16511	Complete genome	1	1	2,101,285	53.9	GCF_000186245.1
<i>Sulfurimonas</i>	<i>denitrificans</i>	DSM 1251	Complete genome	1	1	2,201,561	34.5	GCF_000012965.1
<i>Sulfurimonas</i>	<i>autotrophica</i>	DSM 16294	Complete genome	1	1	2,153,198	35.2	GCF_000147355.1
<i>Sulfurimonas</i>	<i>paralvinellae</i>	GO25	Complete genome	2	2	2,078,871	38.5	GCF_014905135.1
<i>Sulfurimonas</i>	<i>sediminis</i>	S2-6	Complete genome	1	1	2,320,257	37.3	GCF_014905115.1
<i>Sulfurimonas</i>	<i>hydrogeniphila</i>	NW10	Complete genome	1	1	2,342,011	37.3	GCF_009068765.1
<i>Sulfurimonas</i>	<i>indica</i>	NW8N	Contig	52	-	2,093,483	36.8	GCF_009192995.1
<i>Sulfurovum</i>	sp.	NBC37-1	Complete genome	1	1	2,562,277	37.1	GCF_000010345.1
<i>Sulfurovum</i>	<i>lithotrophicum</i>	ATCC BAA-797	Complete genome	1	1	2,217,891	44.3	GCF_000987835.1
<i>Sulfurovum</i>	<i>indicum</i>	ST-419	Complete genome	1	1	2,209,694	42.5	GCF_014931715.1
<i>Sulfurovum</i>	<i>riftiae</i>	1812E	Contig	72	-	2,374,692	45.7	GCF_001595645.1
<i>Persephonella</i>	<i>hydrogeniphila</i>	DSM15103	Contig	19	-	1,998,302	35.1	GCF_900215515.1



