HOKKAIDO UNIVERSITY

Title	Noninstrumented Clinical Assessment of Static Postural Stability in Chronic Ankle Instability : A Systematic Review and Meta-Analysis
Author(s)	Koshino, Yuta; Kobayashi, Takumi
Citation	Journal of sport rehabilitation, 1-11 https://doi.org/10.1123/jsr.2023-0437
Issue Date	2024-08-19
Doc URL	https://hdl.handle.net/2115/93074
Rights	Accepted author manuscript version reprinted, by permission, from Journal of Sport Rehabilitation, 2024, https://doi.org/10.1123/jsr.2023-0437. © Human Kinetics, Inc.
Type	journal article
File Information	R2_Manuscript.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Non-instrumented Clinical Assessment of Static Postural Stability in Chronic Ankle Instability: A Systematic Review and Meta-analysis

Yuta Koshino, PT, PhD[1], Takumi Kobayashi, PT, PhD[2]

1. Faculty of Health Sciences, Hokkaido University, Sapporo, Japan
2. Graduate School of Health Science, Gunma University, Maebashi, Japan

Yuta Koshino

e-mail: y-t-1-6@hs.hokudai.ac.jp, Twitter: @koshino_yuta

Takumi Kobayashi

e-mail: kobatakku@gmail.com, Twitter: @kobatakku

Corresponding author

Yuta Koshino, Ph.D., PT

Faculty of Health Sciences, Hokkaido University, Kita 12, Nishi 5, Kita-ku, Sapporo, 060-0812, Japan

Phone: +81-11-706-3531

E-mail: y-t-1-6@hs.hokudai.ac.jp

Funding: None.

Conflicts of interest: None.

Acknowledgments: None.

The protocol for this review was pre-registered with PROSPERO (CRD42023395396):

https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023395396

Highlights

- The single-leg and tandem stance in the Balance Error Scoring System can detect decreased static postural stability in chronic ankle instability.
- The use of foam surfaces in the Balance Error Scoring System could potentially detect greater static postural stability deficits in chronic ankle instability than firm surfaces.
- Foot-lift and time-in-balance tests may also be useful for detecting static postural stability deficits in chronic ankle instability.

Non-instrumented Clinical Assessment of Static Postural Stability in Chronic Ankle Instability: A Systematic Review and Meta-analysis

Abstract

Context: Several clinical tests are available to assess static postural stability in individuals with chronic ankle instability (CAI); however, it is unclear which test should be used.

Objective: To determine which non-instrumented clinical tests should be used to detect static postural stability deficits in individuals with CAI.

Evidence Acquisition: We searched four databases from their inception to February 2023, and included studies comparing static postural stability in individuals with CAI and healthy controls using non-instrumented assessments. Two reviewers independently extracted study characteristics, participant information, static postural stability assessment methods, and results. We calculated the pooled standardized mean difference (SMD) and 95% confidence interval (CI) using a random-effects meta-analysis and assessed the certainty of the evidence.

Evidence Synthesis: Fourteen cross-sectional studies (293 participants with CAI and 284 healthy controls) were included. The meta-analysis showed no significant differences between the CAI and healthy groups in the double-leg stance condition of the Balance Error Scoring System (BESS) (SMD, –0.03; low-certainty evidence). Significant group differences were found in the BESS single-leg stance (SLS) on firm and foam surfaces (SLS firm: SMD, 0.47, very low-certainty evidence; SLS foam: SMD, 0.80, very low-certainty evidence), the tandem stance (TS) on firm and foam surfaces (TS firm: SMD, 0.39, low-certainty evidence; TS foam: SMD, 0.76, low-certainty evidence), and the total BESS in the foam conditions (SMD, 1.12, very low certainty evidence). Significant differences were also found between the CAI and healthy groups in the foot-lift (SMD, 1.24; very low certainty evidence) and time-in-balance tests (SMD, –0.94; very low certainty evidence).

Conclusions: Due to the large magnitude of the differences, the SLS foam, TS foam, and the total BESS in the foam conditions, as well as the foot-lift test or time-in-balance test, may be the most appropriate to clinically identify static postural stability impairment in individuals with CAI.

Key words: ankle sprain; ankle ligament injury; balance; postural control; rehabilitation

CONTEXT

Lateral ankle sprains (LASs) are one of the most common sports injuries and have a high recurrence rate.[1] The proportion of recurrent ankle sprains in the sports population ranges from 12% to 47%.[2] Of those who sustain an LAS, 40% develop chronic ankle instability (CAI),[3] which can lead to perceived instability, giving way, and reinjury of the ankle.[4] Up to 10% to 20% of athletes who continue to participate in sports activities have CAI.[5] Furthermore, CAI is associated with ankle osteoarthritis and reduced health-related quality of life.[1,6,7]

Several types of impairments are associated with CAI, including pathomechanical, motor-behavioral, and sensory-perceptual impairments.[8] Among these, many studies have been conducted on postural stability deficits, which are classified as motor-behavioral impairments.[9,10] A recent meta-analysis found that individuals with CAI have static postural stability deficits during single-leg standing.[10] Static postural stability is the most commonly assessed outcome in intervention studies,[9] and its assessment in clinical practice guidelines for post-LAS and CAI is recommended.[11] Thus, the importance of assessing static postural stability in individuals with CAI is widely recognized.

A number of methods have been used to assess static postural stability in CAI.[9,10,12] A systematic review of instrument-based assessment of static postural stability in CAI reported that assessment by the center of pressure trajectory using a force plate can identify static postural stability impairment in CAI.[10] However, force plates may not always be available in clinical practice because of their high cost. There are several non- instrumented methods to assess static postural stability, such as the Balance Error Scoring System (BESS), foot-lift test, time-in-balance test, etc.[13] The BESS consists of a total of six conditions, including three standing conditions and two floor conditions. The BESS conditions used varied among the studies; additionally, only single-leg standing on a firm or foam surface was used.[14-16]

However, it remains unclear which specific BESS conditions or other tests (i.e., foot-lift test, time-in-balance test, etc.) should be used to detect differences in static postural stability between individuals with and without CAI. Evidence-based clarification of this issue will provide helpful insights for clinical decision making when selecting an assessment method. Therefore, this study aimed to identify appropriate non-instrumented assessment methods to detect differences in static postural stability between individuals with and without CAI through a systematic review and meta-analysis.

METHODS

Design

This study was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 statement.[17] The protocol for this review was pre-registered with PROSPERO (registration number: blind).

Evidence acquisition

Search strategy

Four databases (PubMed, CINAHL, Cochrane Library, and Web of Sciences) were searched from their inceptions until February 1, 2023. A search strategy was created for each database (Supplemental Content 1), and the search results were exported to Endnote X9 (Thomson Reuters, New York, NY, USA). Duplicate papers were removed using software. Two independent reviewers screened the titles and abstracts of the studies according to eligibility criteria described below. Two reviewers screened the full text of the remaining studies. At each screening stage, disagreements between the reviewers were resolved by consensus.

81

82 ***Study selection***

83 The inclusion criteria for the studies included in this systematic review were as
84 follows: (1) they compared static postural stability between individuals with CAI and healthy
85 controls, (2) static postural stability was assessed using non-instrumented methods, and (3)
86 the criteria for CAI included a history of at least one ankle sprain, and a history of giving way,
87 recurrent sprains, and/or a feeling of instability.[4] The exclusion criteria were as follows (1)
88 studies that did not include healthy participants as a control group, and (2) non-English
89 articles. Static postural stability was defined as the measure of postural stability when the
90 body is stationary.[18]

91

92 ***Data extraction***

93 Two reviewers independently extracted the following data: study characteristics
94 (author name and year of publication), participant information (sample size and definitions of
95 the injured group), static postural stability assessment methods (type of test and conditions),
96 and the mean and standard deviation (SD) of the static postural stability assessment results
97 (injured and healthy control groups). When the median and quartiles were reported, the mean
98 and SD were estimated using the Box-Cox method.[19] If static postural stability data were not
99 reported in the paper, we requested the data from the corresponding author by e-mail. Studies
100 with unavailable data were excluded from the meta-analysis.

101

102 ***Risk of bias***

103 Two reviewers independently assessed the risk of bias of the included studies. There
104 is no gold standard for assessing the risk of bias in cross-sectional studies,[20] but the
105 Newcastle-Ottawa Scale (NOS)[21] is often used with modifications. Therefore, we adapted the

NOS for use in a cross-sectional study based on previous studies.[20,22-24] For sample selection, we assessed four items: (1) The representativeness of the sample (was the sample truly or somewhat representative of the target population?), (2) Sample size (justified or satisfactory?), (3) definition of injured population (was a validated measurement tool used?), and (4) non-included participants (was comparability established and was the inclusion rate satisfactory?). In addition, we assessed the comparability of cases and controls based on design or analysis. For outcomes, the following two items were assessed: (1) Outcome assessment (was an independent blind assessment conducted using an objectively validated method?) (2) Statistical tests (is an appropriate test used and clearly described?). Scores were assigned for each item assessment and interpreted as follows: 8–10 points, low risk of bias; 6–7 points, medium risk of bias; 5 points or less, high risk of bias.[23]

Data analysis

Meta-analysis (random effects) was conducted using Review Manager 5.4 (RevMan).[25] Data for similar postural stability assessment methods from each study were pooled, and standardized mean differences (SMDs) and 95% confidence intervals (CIs) were calculated using RevMan. The SMD, rather than the mean difference, was calculated to examine the magnitude of the differences between the CAI and healthy groups based on the method of assessment. A positive SMD, except for the time-in-balance test, indicated that the CAI group had decreased postural stability compared to the healthy control group. Differences were considered statistically significant if the 95% CI of the SMD did not include a 0. The SMD was interpreted as small if less than 0.40, moderate if 0.40 to 0.70, and large if greater than 0.70.[26] Statistical heterogeneity of 0% was interpreted as possibly unimportant, 30% as moderate, 50% as substantial, and 75% as considerable, based on the I^2 statistic.[26] Publication bias was assessed using a funnel plot when more than 10 studies were available in

131 the meta-analysis.[27]

132 Finally, two reviewers independently assessed the certainty of the evidence for each
133 meta-analysis using the Grading of Recommendations Assessment, Development, and
134 Evaluation methodology.[28] The certainty of the evidence, based on previous studies,[29,30] was
135 downgraded if the following five criteria were met: more than 75% of the studies were not
136 considered to be at low risk of bias; I^2 statistic greater than 50%; wide 95% CI (upper or
137 lower 95% CI >0.5 of the effect size in either direction) and a sample size of fewer than 400
138 in the meta-analysis for each variable; studies with indirect comparisons; asymmetric funnel
139 plots. The certainty of evidence was downgraded from high to moderate, low, or very low
140 when each of the above criteria was met.

141

142 **Evidence synthesis**

143 ***Study selection and characteristics***

144 Fourteen studies (293 patients with CAI and 284 healthy controls) were included
145 after screening (Figure 1). Table 1 summarizes the characteristics of the included studies. All
146 the included studies examined static postural stability in individuals with CAI using a cross-
147 sectional design. Non-instrumented assessments included the BESS (n=8),[31-38] foot-lift test
148 (n=4),[36,39-41] time-in-balance test (n=4),[36,40-42] Clinical Test of Sensory Integration in Balance
149 (n=1),[43] and Romberg test (n=1).[44] Table 2 describes each assessment method.

150

151 ***Risk of bias***

152 Table 3 presents the results of the risk of bias assessment. Eight studies (57.1%) were
153 considered high risk, six (42.9%) were considered medium risk, and none were considered
154 low risk. Most studies had a high risk of sample representativeness (92.9%), non-included
155 participants (100.0%), and sample size justification (64.3%). In addition, only 1.4% and 0.7%

of the studies scored 2 points for comparability and outcome assessment, respectively.

Meta-analysis of non-instrumented static postural stability

Balance Error Scoring System

Eight studies used the BESS, of which four studies examined double-leg stance (DLS) conditions on firm (DLS firm) and foam (DLS foam) surfaces.[31,32,34,36] The SMD of the DLS firm could not be estimated because the mean and SD were zero in most studies. The pooled data for DLS foam showed no significant difference between the CAI and control groups (SMD, –0.03; 95% CI, –0.34-0.28; I^2=7%; low-certainty evidence) (Figure 2 and Table 4). In the single-leg stance condition of the BESS, six and four studies examined firm (SLS firm) and foam (SLS foam) surfaces, respectively.[31,32,34-36,38] In the SLS firm condition, the CAI group showed significantly more errors than the control group (SMD: 0.47; 95% CI, 0.07-0.86; I^2=57%; very low certainty evidence) (Figure 2 and Table 4). The CAI group also had significantly more errors than the control group in the SLS foam condition (SMD, 0.80; 95% CI, 0.24-1.35; I^2=67%; very low certainty evidence) (Figure 2 and Table 4). The tandem stance (TS) on firm (TS firm) and foam (TS foam) surfaces was examined in four studies.[31,32,34,36] The CAI group had significantly more errors than the control group in both the TS firm (SMD, 0.39; 95% CI, 0.09-0.69; I^2=0%; low certainty evidence) and TS foam (SMD, 0.76; 95% CI, 0.45-1.07; I^2=0%; low certainty evidence) conditions.

Four studies were used for the total scores for all BESS conditions.[32-34,36] The pooled SMD showed no significant difference between the CAI and control groups (SMD, 0.40; 95% CI, –0.25-1.05; I^2=76%; very low certainty evidence) (Figure 3 and Table 4). This statistical heterogeneity was considerable, and the result was significant when the study with the highest heterogeneity was excluded (SMD, 0.70; 95% CI, 0.24-1.15, I^2=34%). In addition, 2 studies reported the total scores for the firm (firm total) and foam (foam total) surface conditions,

181 respectively.[34,37] The meta-analysis showed no significant group differences in the firm total
182 (SMD, 0.61; 95% CI, −1.52-2.73; I^2=91%; very low certainty evidence) (Figure 3 and Table
183 4), whereas the foam total had a significantly higher number of errors in the CAI group
184 (SMD, 1.12; 95% CI, 0.23-2.02; I^2=51%; very low certainty evidence) (Figure 3 and Table 4).
185 Statistical heterogeneity was considerable for total firm scores; however, there were only two
186 studies.
187
188 *Foot-lift test*
189 Four studies used the foot-lift test,[36,39-41] but only three were included in the meta-
190 analysis.[36,39,41] This was because the mean and SD could not be extracted from one study.[40]
191 The pooled data showed that the CAI group had a significantly greater number of foot lifts
192 than the control group (SMD, 1.24; 95% CI, 0.82-1.66; I^2=90%; very low certainty evidence)
193 (Figure 4 and Table 4). Statistical heterogeneity was significant. When the study that
194 contributed most to the heterogeneity was excluded,[39] the I^2 statistic was 0%, and the
195 statistical result was similar (SMD, 0.72; 95% CI, 0.24-1.20; I^2, 0%).
196
197 *Time-in-balance test*
198 Four studies used the time-in-balance test,[36,40-42] but only three were included in the
199 meta-analysis.[36,41,42] This was because the mean and SD could not be extracted from one
200 study.[40] A meta-analysis showed that the CAI group maintained postural stability for a
201 significantly shorter time than the control group (SMD, −0.94; 95% CI, −1.29 to −0.59;
202 I^2=50%; very low certainty evidence) (Figure 5 and Table 4).
203
204 *Other tests*
205 One study used the Clinical Test of Sensory Integration in Balance (CTSIB) to

compare a CAI group with a control group.[43] A meta-analysis could not be performed because only one study was available for the CTSIB. The CAI group had significantly worse scores for conditions 2 and 6 of the CTSIB, CTSIB composite score, and CTSIB somatosensory ratio than the control group.[43] The Romberg test was used in one study, and no significant group differences were found.[44]

DISCUSSION

This systematic review and meta-analysis aimed to determine which non-instrumented static postural stability assessment methods are useful for individuals with CAI. The present meta-analysis suggests that the SLS firm, SLS foam, TS firm, TS foam, and foam total in the BESS as well as the foot-lift and time-in-balance tests can detect static postural stability deficits in the CAI group. In particular, the pooled SMD showed that the SLS foam, TS foam, and foam total in the BESS, foot-lift test, and time-in-balance test had large effects (SMD >0.7). Therefore, the present study suggests that these assessments can be used in clinical practice to detect static postural stability deficits associated with CAI. However, it should be noted that most of the certainty of the evidence was very low to low.

The BESS is the most commonly used non-instrumented static postural stability assessment tool for individuals with CAI. This is also the clinical assessment recommended in the consensus statement of the International Ankle Consortium.[45] The present study suggests that the SLS and TS conditions can detect static postural stability impairment in the CAI group but not in the DLS condition. No significant group differences were observed in DLS. The DLS conditions of the BESS are easy and may not be necessary for static postural stability assessment in the CAI group. The SLS and TS conditions detected significant group differences for each surface condition; however, these SMDs were larger when using the foam surface than when using the firm condition. The total number of errors for the foam condition

also exhibited a high SMD. These findings suggest that assessing errors in SLS or TS on a foam surface is useful for detecting static postural stability impairments caused by CAI.

A meta-analysis of the foot-lift test showed a large pooled SMD, suggesting that it is useful for assessing static postural stability in the CAI group. As this statistical heterogeneity was substantial, we confirmed that excluding the study with high heterogeneity resulted in a large SMD (0.72).[39] A previous meta-analysis also showed that the foot-lift test can discriminate between the CAI and healthy groups.[13] This test is performed on a firm surface, and unlike the BESS, the stance is maintained for 30 seconds and the number of times any part of the foot lifts from the floor is counted.[36,39,41] A cross-sectional study by Linens et al also reported a larger effect size for the difference between the CAI and healthy groups in the foot-lift test than in the SLS firm of the BESS.[36] In the single-leg standing on a firm surface, the assessment of foot-focused errors for 30 seconds may be useful to detect static postural stability impairments in CAI.

For the time-in-balance test, the CAI and healthy groups were significantly different, with a large pooled SMD. The synthesized effect size for the time-in-balance test has not been reported in a previous meta-analysis,[13] whereas a cross-sectional study reported that the time-in-balance test had an effect size as large as that of the foot-lift test for the difference between the CAI and healthy groups.[36] The present meta-analysis suggests that the time-in-balance test would be useful for detecting static postural stability impairment in CAI. It is noteworthy that the mean value of the time-in-balance test for the CAI group was less than 10 seconds in two studies,[41,42] while it was 28.99 seconds in one study.[36] As a possible explanation for this, Wikstrom et al. discussed the potential variation in the definition of balance loss in this test among different studies.[41] In the time-in-balance test, loss of balance is defined as the movement of the tested foot or the touching of the ground by the contralateral foot during single-leg standing. However, the definition of a moving test foot was not specific. In the

future, it will be necessary to define this as any part of the foot moving off the floor (e.g., medial foot elevation, heel lateral shift, etc.), similar to the study by Wikstrom et al.[41]

In clinical practice, the SLS foam, TS foam, or the total score of the BESS foam surface condition, as well as either the foot-lift test or the time-in-balance test, can be used to detect static postural stability impairments in individuals with CAI because these conditions or tests have large SMDs. For example, in patients who have difficulty standing on a single leg, static postural stability may be adequately assessed by assessing them in the TS condition, preferably using a foam surface. Additionally, if clinicians do not have access to a foam pad, a foot-lift test or time-in-balance test may be a better option. Furthermore, unlike the BESS, which necessitates the observation of errors throughout the body (from checking eye openings to hip and foot movements), these two tests only require observation of the test foot. This may enable the examiner to observe small foot movements without overlooking them, and may facilitate the detection of reduced postural stability due to CAI. Additionally, clinical postural stability assessment in the DLS condition may be unnecessary because of the difficulty in detecting postural stability impairments associated with CAI. Assessment of the total score of the BESS foam surface condition requires three conditions for a 20-second postural stability test: DLS, SLS, and TS foam conditions. This is not more time-efficient than other tests, such as SLS foam only, TS foam only, foot-lift test, and time-in-balance test. This information may also be useful in time-constrained settings.

Study limitations

This study had several limitations. First, there were no low risk of bias studies in the present review. Therefore, the results should be interpreted with caution. Second, the statistical heterogeneity of most of the meta-analyses in this review ranged from substantial to considerable. This indicates that the effect sizes varied between the studies included in the

meta-analysis and that the results should be interpreted with caution. Third, the definition of CAI varied among included studies. Finally, although we were unable to perform a meta-analysis, a cross-sectional study found significant differences between CAI and healthy participants on several scores of the CTSIB scores.[43] This test differs from the other tests in that it has a visual conflict dome condition to produce a visual sway referencing effect.[43] The significant group difference was found in the visual-conflict dome condition.[43] Static postural stability was also assessed using a stroboscopic glass that disrupts visual information,[46] which is cost-effective and easy to use in clinical practice. Future studies that use such tools to manipulate visual information are likely to contribute to the development of clinical assessment methods for postural stability in individuals with CAI.

CONCLUSIONS

Our systematic review and meta-analysis showed that SLS firm, SLS foam, TS firm, TS foam, and foam total in the BESS as well as the foot-lift and time-in-balance tests could detect static postural stability impairment in the CAI group. In particular, the pooled SMD showed that the SLS foam, TS foam, and foam total in the BESS, foot-lift test, and time-in-balance test had a large effect (SMD >0.7), and therefore can be used to detect static postural stability impairment in individuals with CAI. However, the certainty of the evidence was very low to low, and therefore the present results should be interpreted with caution.

REFERENCES

1. Gribble PA, Bleakley CM, Caulfield BM, et al. Evidence review for the 2016 International Ankle Consortium consensus statement on the prevalence, impact and long-term consequences of lateral ankle sprains. *Br J Sports Med*. Dec 2016;50(24):1496-1505. doi:10.1136/bjsports-2016-096189
2. Herzog MM, Kerr ZY, Marshall SW, Wikstrom EA. Epidemiology of Ankle Sprains and Chronic Ankle Instability. *J Athl Train*. Jun 2019;54(6):603-610. doi:10.4085/1062-6050-447-17
3. Doherty C, Bleakley C, Hertel J, Caulfield B, Ryan J, Delahunt E. Recovery From a First-Time Lateral Ankle Sprain and the Predictors of Chronic Ankle Instability: A Prospective Cohort Analysis. *Am J Sports Med*. Apr 2016;44(4):995-1003. doi:10.1177/0363546516628870
4. Gribble PA, Delahunt E, Bleakley C, et al. Selection criteria for patients with chronic ankle instability in controlled research: a position statement of the International Ankle Consortium. *Br J Sports Med*. Jul 2014;48(13):1014-8. doi:10.1136/bjsports-2013-093175
5. Koshino Y, Samukawa M, Murata H, et al. Prevalence and characteristics of chronic ankle instability and copers identified by the criteria for research and clinical practice in collegiate athletes. *Phys Ther Sport*. Sep 2020;45:23-29. doi:10.1016/j.ptsp.2020.05.014
6. Golditz T, Steib S, Pfeifer K, et al. Functional ankle instability as a risk factor for osteoarthritis: using T2-mapping to analyze early cartilage degeneration in the ankle joint of young athletes. *Osteoarthritis Cartilage*. Oct 2014;22(10):1377-85. doi:10.1016/j.joca.2014.04.029
7. Houston MN, Hoch JM, Hoch MC. Patient-Reported Outcome Measures in Individuals With

Chronic Ankle Instability: A Systematic Review. *J Athl Train*. Oct 2015;50(10):1019-33. doi:10.4085/1062-6050-50.9.01

8. Hertel J, Corbett RO. An Updated Model of Chronic Ankle Instability. *J Athl Train*. Jun 2019;54(6):572-588. doi:10.4085/1062-6050-344-18
9. Nozu S, Takemura M, Sole G. Assessments of Sensorimotor Deficits Used in Randomized Clinical Trials With Individuals With Ankle Sprains and Chronic Ankle Instability: A Scoping Review. *PM R*. Aug 2021;13(8):901-914. doi:10.1002/pmrj.12487
10. Xue X, Wang Y, Xu X, et al. Postural Control Deficits During Static Single-leg Stance in Chronic Ankle Instability: A Systematic Review and Meta-Analysis. *Sports Health*. Mar 5 2023:19417381231152490. doi:10.1177/19417381231152490
11. Martin RL, Davenport TE, Fraser JJ, et al. Ankle Stability and Movement Coordination Impairments: Lateral Ankle Ligament Sprains Revision 2021. *J Orthop Sports Phys Ther*. Apr 2021;51(4):CPG1-CPG80. doi:10.2519/jospt.2021.0302
12. Koshino Y, Kobayashi T. Effects of Conservative Interventions on Static and Dynamic Balance in Individuals With Chronic Ankle Instability: A Systematic Review and Meta-analysis. *Arch Phys Med Rehabil*. Apr 2023;104(4):673-685. doi:10.1016/j.apmr.2022.10.014
13. Rosen AB, Needle AR, Ko J. Ability of Functional Performance Tests to Identify Individuals With Chronic Ankle Instability: A Systematic Review With Meta-Analysis. *Clin J Sport Med*. Nov 2019;29(6):509-522. doi:10.1097/JSM.0000000000000535
14. Hall EA, Chomistek AK, Kingma JJ, Docherty CL. Balance- and Strength-Training Protocols to Improve Chronic Ankle Instability Deficits, Part I: Assessing Clinical Outcome Measures. *J Athl Train*. Jun 2018;53(6):568-577. doi:10.4085/1062-6050-385-16
15. McKeon PO, Wikstrom EA. Sensory-Targeted Ankle Rehabilitation Strategies for Chronic

Ankle Instability. *Med Sci Sports Exerc*. May 2016;48(5):776-84. doi:10.1249/mss.0000000000000859

16. Beazell JR, Grindstaff TL, Sauer LD, Magrum EM, Ingersoll CD, Hertel J. Effects of a proximal or distal tibiofibular joint manipulation on ankle range of motion and functional outcomes in individuals with chronic ankle instability. *J Orthop Sports Phys Ther*. Feb 2012;42(2):125-34. doi:10.2519/jospt.2012.3729
17. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. Mar 29 2021;372:n71. doi:10.1136/bmj.n71
18. Gonçalves C, Bezerra P, Clemente F, et al. The relationship between static and dynamic balance in active young adults. *Human Movement*. 2022;23(2):65-75. doi:10.5114/hm.2021.106165
19. McGrath S, Zhao X, Steele R, Thombs BD, Benedetti A, Collaboration DESD. Estimating the sample mean and standard deviation from commonly reported quantiles in meta-analysis. *Stat Methods Med Res*. Jan 30 2020:962280219889080. doi:10.1177/0962280219889080
20. Moskalewicz A, Oremus M. No clear choice between Newcastle-Ottawa Scale and Appraisal Tool for Cross-Sectional Studies to assess methodological quality in cross-sectional studies of health-related quality of life and breast cancer. *J Clin Epidemiol*. Apr 2020;120:94-103. doi:10.1016/j.jclinepi.2019.12.013
21. Wells GA, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. 1/12, 2020. Accessed 9 Feb 2022, http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
22. Nyawo TA, Dludla PV, Mazibuko-Mbeje SE, et al. A systematic review exploring the significance of measuring epicardial fat thickness in correlation to B-type natriuretic peptide

levels as prognostic and diagnostic markers in patients with or at risk of heart failure. *Heart Fail Rev*. Mar 2022;27(2):665-675. doi:10.1007/s10741-021-10160-3

23. Ribeiro CM, Beserra BTS, Silva NG, et al. Exposure to endocrine-disrupting chemicals and anthropometric measures of obesity: a systematic review and meta-analysis. *BMJ Open*. Jun 21 2020;10(6):e033509. doi:10.1136/bmjopen-2019-033509
24. West TJ, Bruder AM, Crossley KM, Culvenor AG. Unilateral tests of lower-limb function as prognostic indicators of future knee-related outcomes following anterior cruciate ligament injury: a systematic review and meta-analysis of 13 150 adolescents and adults. *Br J Sports Med*. Jan 20 2023;doi:10.1136/bjsports-2022-105736
25. The Cochrane Collaboration. Review Manager (RevMan) [Computer program]. Version 5.4. .
26. Higgins JPT, Thomas J, Chandler J, et al. Cochrane Handbook for Systematic Reviews of Interventions version 6.2 (updated February 2021). *Cochrane*. 2021;Available from www.training.cochrane.org/handbook.
27. Sterne JA, Sutton AJ, Ioannidis JP, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ*. Jul 22 2011;343:d4002. doi:10.1136/bmj.d4002
28. Balshem H, Helfand M, Schunemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol*. Apr 2011;64(4):401-6. doi:10.1016/j.jclinepi.2010.07.015
29. Karanasios S, Korakakis V, Whiteley R, Vasilogeorgis I, Woodbridge S, Gioftsos G. Exercise interventions in lateral elbow tendinopathy have better outcomes than passive interventions, but the effects are small: a systematic review and meta-analysis of 2123 subjects in 30 trials. *Br J Sports Med*. May 2021;55(9):477-485. doi:10.1136/bjsports-2020-102525

30. Ardern CL, Buttner F, Andrade R, et al. Implementing the 27 PRISMA 2020 Statement items for systematic reviews in the sport and exercise medicine, musculoskeletal rehabilitation and sports science fields: the PERSiST (implementing Prisma in Exercise, Rehabilitation, Sport medicine and SporTs science) guidance. *Br J Sports Med*. Feb 2022;56(4):175-195. doi:10.1136/bjsports-2021-103987
31. Dobo B, White A, Linens S. Balance Error Scoring System Stances That Identify Division I Athletes With Chronic Ankle Instability Most in Need of Rehabilitation. *Athl Train Sports Health Care*. 2015;7(5):190-196. doi:10.3928/19425864-20150831-04
32. Docherty CL, Valovich McLeod TC, Shultz SJ. Postural control deficits in participants with functional ankle instability as measured by the balance error scoring system. *Clin J Sport Med*. May 2006;16(3):203-8. doi:10.1097/00042752-200605000-00003
33. John C, Stotz A, Gmachowski J, et al. Is an Elastic Ankle Support Effective in Improving Jump Landing Performance, and Static and Dynamic Balance in Young Adults With and Without Chronic Ankle Instability? *J Sport Rehabil*. Aug 1 2020;29(6):789-794. doi:10.1123/jsr.2019-0147
34. Koshino Y, Akimoto M, Kawahara D, et al. Inertial Sensor-Based Assessment of Static Balance in Athletes with Chronic Ankle Instability. *J Sports Sci Med*. Mar 2023;22(1):36-43. doi:10.52082/jssm.2023.36
35. Kwon YU. Static Postural Stability in Chronic Ankle Instability, an Ankle Sprain and Healthy Ankles. *Int J Sports Med*. Jul 2018;39(8):625-629. doi:10.1055/a-0608-4552
36. Linens SW, Ross SE, Arnold BL, Gayle R, Pidcoe P. Postural-stability tests that identify individuals with chronic ankle instability. *J Athl Train*. Jan-Feb 2014;49(1):15-23. doi:10.4085/1062-6050-48.6.09

37. Powell MR, Powden CJ, Houston MN, Hoch MC. Plantar cutaneous sensitivity and balance in individuals with and without chronic ankle instability. *Clin J Sport Med*. Nov 2014;24(6):490-6. doi:10.1097/jsm.0000000000000074
38. Ross SE, Linens SW, Wright CJ, Arnold BL. Balance assessments for predicting functional ankle instability and stable ankles. *Gait Posture*. Oct 2011;34(4):539-42. doi:10.1016/j.gaitpost.2011.07.011
39. Hiller CE, Refshauge KM, Herbert RD, Kilbreath SL. Balance and recovery from a perturbation are impaired in people with functional ankle instability. *Clin J Sport Med*. Jul 2007;17(4):269-75. doi:10.1097/JSM.0b013e3180f60b12
40. Ko J, Rosen AB, Brown CN. Comparison Between Single and Combined Clinical Postural Stability Tests in Individuals With and Without Chronic Ankle Instability. *Clin J Sport Med*. Jul 2017;27(4):394-399. doi:10.1097/jsm.0000000000000354
41. Wikstrom EA, Kyeongtak S, Cug M, Bentley RL. Clinical Balance Measures in Patients With Chronic Ankle Instability, Copers, and Uninjured Controls. *Athl Train Sports Health Care*. 2017;9(1):11-16. doi:10.3928/19425864-20160822-02
42. Chrintz H, Falster O, Roed J. Single-leg postural equilibrium test. *Scand J Med Sci Sports*. 1991;1(4):244-246. doi:10.1111/j.1600-0838.1991.tb00305.x
43. Song K, Wikstrom EA. Laboratory- and Clinician-Oriented Measures of Sensory Organization Strategies in Those With and Without Chronic Ankle Instability. *Athl Train Sports Health Care*. 2020;12(6):257-264. doi:10.3928/19425864-20200821-01
44. Yalfani A, Gandomi F, Kohboomi M. The Effect of G-Max and G-Med Muscles Fatigue on Functional Performance and Balance in Athletes with and without Chronic Ankle Instability. *Asian J Sports Med*. 2017;8(3):1-6. doi:10.5812/asjsm.55444

45. Delahunt E, Bleakley CM, Bossard DS, et al. Clinical assessment of acute lateral ankle sprain injuries (ROAST): 2019 consensus statement and recommendations of the International Ankle Consortium. *Br J Sports Med*. Oct 2018;52(20):1304-1310. doi:10.1136/bjsports-2017-098885

46. Han S, Lee H, Son SJ, Hopkins JT. The effects of visual feedback disruption on postural control with chronic ankle instability. *J Sci Med Sport*. Jan 2022;25(1):53-57. doi:10.1016/j.jsams.2021.07.014

Figure legends

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram.

Figure 2. Meta-analyses on each condition of the Balance Error Scoring System. This forest plot was generated by RevMan 5.4.[25] CAI, chronic ankle instability; DLS foam, double leg stance on the foam surface; SLS firm, single leg stance on the firm surface; SLS foam, single leg stance on the foam surface; TS firm, tandem stance on the firm surface; TS firm, tandem stance on the firm surface.

Figure 3. Meta-analyses on the total scores of the Balance Error Scoring System. This forest plot was generated by RevMan 5.4.[25] CAI, chronic ankle instability.

Figure 4. Meta-analyses on the foot lift test. This forest plot was generated by RevMan 5.4.[25] CAI, chronic ankle instability.

Figure 5. Meta-analyses on the time-in-balance test. This forest plot was generated by RevMan 5.4.[25] CAI, chronic ankle instability.

Table 1. Study characteristics

Author	N	Key definitions of CAI	Clinical balance assessment	Findings
Chrintz[42]	CAI: 29 CON: 41	1) a history of recurrent ASs and GW incidents resistant to training; 2) prior to operative exploration of the lateral ligaments of the ankle.	· **Time in Balance test in EC** and EO	· Significantly less time in the CAI group than in the CON group. **Mean and SD were estimated based on reported medians and quartiles for meta-analysis.*
Dobo[31]	CAI: 32 CON: 19	1) at least one significant AS; 2) self-reported sensations of ankle GW during activity.	· BESS 6 conditions: **(1) DLS firm, (2) DLS foam, (3) SLS firm, (4) SLS foam, (5) TS firm, (6) TS foam**	· No results of group comparisons were reported. · Significant cutoffs were found in the SLS firm, SLS foam, and TS foam conditions that could identify individuals with CAI.
Docherty[32]	CAI: 30 CON: 30	1) at least one AS; 2) repeated feeling of GW or instability in the ankle	· BESS 6 conditions: **(1) DLS firm, (2) DLS foam, (3) SLS firm, (4) SLS foam, (5) TS firm, (6) TS foam** · **BESS total errors**	· Significantly more errors in the SLS firm, SLS foam, and TS foam conditions in the CAI group than in the CON group. · Significantly more total BESS errors in the CAI group than in the CON group.
Hiller[39]	CAI: 31 CON: 15	1) at least one AS; 2) CAIT score ≤ 24.	· **Foot lift test**	· Significantly more foot lifts in the CAI group than in the CON group. **Data from the unilateral and bilateral instability groups were combined for meta-analysis.*
John[33]	CAI: 20 CON: 20	1) a history of at least one AS, GW and/or RAS and/or feelings of instability; 2) CAIT score ≤ 24; 3) FAAM score < 90%; 4) FAAM Sports < 80%	· **BESS total errors**	· No significant group effect on the total BESS errors.
Ko[40]	CAI: 25 CON: 33	1) a history of moderate–severe AS that required at least 3 days of partial or non-weight bearing; 2) a history of GW with activity; 3) CAIT score ≤ 25	· Foot lift test · Time in Balance test	· No results of group comparisons were reported.
Koshino[34]	CAI: 16 CON: 16	1) at least one AS; 2) the initial sprain over 1 year ago; 3) at least two GW within 6 months and/or recurrent AS; 4) CAIT score ≤ 25.	· BESS 6 conditions: **(1) DLS firm, (2) DLS foam, (3) SLS firm, (4) SLS foam, (5) TS firm, (6) TS foam** · BESS scores: **(1) total errors, (2) firm total errors, (3) foam total errors**	· Significantly more total foam errors in the CAI group than in the CON group.

Kwon[35]	CAI: 20 CON: 20	1) at least one AS; 2) multiple episodes of GW in the 6 months and/or 'feeling of instability'; 3) CAIT score < 24; 4) FAAM score < 90%.	· **BESS SLS firm condition in EC** and EO	· No significant group differences in errors for the SLS firm condition in EC and EO.
Linens[36]	CAI: 17 CON: 17	1) at least one significant AS; 2) self-reported sensations of GW at least twice; 3) CAIT score < 27; 4) no signs or symptoms of an acute injury.	· BESS 6 conditions: **(1) DLS firm, (2) DLS foam, (3) SLS firm, (4) SLS foam, (5) TS firm, (6) TS foam** · **BESS total errors** · **Foot lift test** · **Time in Balance test**	· No results of group comparisons were reported. · Significant cutoffs were found in the BESS SLS firm, total BESS, foot lift test, time-in balance test and that could identify individuals with CAI.
Powell[37]	CAI: 10 CON: 10	1) at least one AS; 2) at least 2 episodes of GW in the past 3 months; 3) AII score ≥ 4.	· BESS scores: **(1) firm total errors, (2) foam total errors**	· Significantly more total firm and total foam errors in the CAI group than in the CON group. **Mean and SD were estimated based on reported medians and quartiles for meta-analysis.*
Ross[38]	CAI: 17 CON: 17	1) self-reported history of AS; 2) least two episodes of GW sensations within the year prior to enrolling.	· **BESS SLS firm condition**	· Although not compared between groups, regression analysis indicated more errors in the SLS firm in the CAI group than in the CON group.
Song[43]	CAI: 15 CON: 15	1) at least one AS; 2) at least two episodes of GW within the past 6 months; 3) IdFAI score > 11; 4) FAAM score ≤ 90%; 4) FAAM Sports ≤ 80%.	· Ten parameters of the CTSIB in single leg standing.	· Significantly more scores of the condition 2 and 6 in CTSIB, CTSIB composite score, and CTSIB somatosensory ratio in the CAI group than in the CON group.
Wikstrom[41]	CAI: 19 CON: 19	1) at least one AS; 2) a minimum of two GW episodes within the past 12 months.	· **Foot lift test** · **Time in Balance test**	· No significant group differences in foot lift test and time-in balance test.
Yalfani[44]	CAI: 12 CON: 12	1) history of ankle injury which caused limping or pain for at least 24 hours; 2) at least two episodes of GW within the past 6 months; 3) FADI score ≤ 89%; 4) FADI Sports ≤ 84%.	· Romberg test in EO and EC	· No significant group differences in Romberg test in EO and EC.

Bold and underlined text indicate data used in the meta-analysis. AII, Ankle Instability Instrument; AS, lateral ankle sprain; BESS, Balance Error Scoring System; CAIT, Cumberland Ankle Instability Tool; CTSIB, Clinical Test of Sensory Integration in Balance; DLS firm, double leg stance on the firm surface; DLS foam, double leg stance on the foam surface; EC, eyes closed; EO, eyes open; FAAM, Foot and Ankle Ability Measure; FADI, Foot and Ankle Disability Index; GW, giving way; IdFAI, Identification of Functional Ankle Instability; SD, standard deviation; SLS firm, single leg stance on the firm surface; SLS foam, single leg stance on the foam surface; TS firm, tandem stance on the firm surface; TS firm, tandem stance on the firm surface.

Table 2. Non-instrumented static postural stability assessment methods

Test	Posture	Conditions	Test time	Outcome measure
Balance Error Scoring System	Maintaining posture in each condition with EC and hands on hips	Six conditions: 1) DLS firm, 2) DLS foam, 3) SLS firm, 4) SLS foam, 5) TS firm, 6) TS foam.	Each condition for 20 sec	Count the number of errors. The maximum number of errors for each condition is 10. The following outcomes were reported: number of errors in each condition, total number of errors in all conditions, firm surface conditions, and foam surface conditions, respectively.
Foot-lift test	Single leg standing on a firm surface with EC	None.	For 30 sec	Count as an error if part of the foot is lifted, the contralateral foot touches the floor, and the contralateral foot remains on the floor (every second).
Time-in-Balance test	Single leg standing on a firm surface with EC	None.	Maximum 60 sec	Measure the time (seconds) until a part of the test foot moves off on the floor or the contralateral foot touches the floor.
Clinical Test of Sensory Integration in Balance	Single leg standing	Six sensory conditions: 1) firm and EO, 2) firm and EC, 3) firm and visual-conflict dome, 4) foam and EO, 5) foam and EC, 6) foam and visual-conflict dome.	Maximum 30 sec	Measure the time (in seconds) that the participant maintains postural stability.
Romberg test	Standing with both feet together	EO and EC.	N/A	Measure the time that the participant maintains postural stability.

DLS firm, double leg stance on the firm surface; DLS foam, double leg stance on the foam surface; EC, eyes closed; EO, eyes open; N/A, not applicable; SLS firm, single leg stance on the firm surface; SLS foam, single leg stance on the foam surface; TS firm, tandem stance on the firm surface; TS firm, tandem stance on the firm surface.

Table 3. Risk of bias assessment of each study using the modified Newcastle-Ottawa Scale

	Selection				Comparability	Outcome		
Author	Representativeness of the sample	Definition of injured population	Non-included participants	Sample size justification	Comparability of cases and controls on the basis of the design or analysis	Assessment of outcome	Statistical test	Total scores
Chrintz[42]	1	1	0	0	0	1	1	4
Dobo[31]	0	1	0	0	2	1	1	5
Docherty[32]	0	1	0	0	2	1	1	5
Hiller[39]	0	2	0	0	1	1	1	5
John[33]	0	2	0	0	1	1	1	5
Ko[40]	0	2	0	1	1	2	1	7
Koshino[34]	0	2	0	1	1	1	1	6
Kwon[35]	0	2	0	1	1	1	1	6
Linens[36]	0	2	0	0	1	1	1	5
Powell[37]	0	2	0	0	1	1	1	5
Ross[38]	0	1	0	1	1	1	1	5
Song[43]	0	2	0	1	1	1	1	6
Wikstrom[41]	0	1	0	0	1	1	1	4
Yalfani[44]	0	2	0	0	1	1	1	5


Identification
Records identified from:
PubMed (n = 1798)
CINAHL (n = 1682)
Web of Science (n = 2754)
Cochrane Library (n = 997)
Records removed before screening:
Duplicate records removed (n = 2976)
Records identified from:
Websites (n = 1)
Citation searching (n = 1)
Screening
Records screened (n = 4225)
Records excluded** (n = 3892)
Reports sought for retrieval (n = 333)
Reports not retrieved (n = 0)
Reports sought for retrieval (n = 2)
Reports not retrieved (n = 0)
Reports assessed for eligibility (n = 12)
Reports excluded:
Participants (n = 46)
Outcome (n = 244)
Design (n = 9)
Not in English (n = 10)
Not full-text (n = 3)
Multiple reports from the same cohort (n = 9)
Reports assessed for eligibility (n = 2)
Reports excluded (n = 0)
Included
Studies included in review (n = 14)

Figure 1

Figure 2


CAI
Control
Std. Mean Difference
Std. Mean Difference
Study or Subgroup Mean SD Total Mean SD Total Weight IV, Random, 95% CI
IV, Random, 95% CI
1.1.1 DLS foam
Dobo 2015 0.06 0.25 32 0.16 0.5 19 27.4% -0.27 [-0.84, 0.30]
Docherty 2006 0.17 0.53 30 0.2 0.48 30 34.1% -0.06 [-0.56, 0.45]
Koshino 2023 0.75 1.18 16 0.25 0.58 16 18.4% 0.52 [-0.18, 1.23]
Linens 2014 0.06 0.24 17 0.11 0.33 17 20.1% -0.17 [-0.84, 0.50]
Subtotal (95% CI) 95 82 100.0% -0.03 [-0.34, 0.28]
Heterogeneity: Tau² = 0.01; Chi² = 3.23, df = 3 (P = 0.36); I² = 7%
Test for overall effect: Z = 0.20 (P = 0.84)
1.1.2 SLS firm
Dobo 2015 2.28 1.8 32 1.84 1.3 19 18.3% 0.27 [-0.31, 0.84]
Docherty 2006 2.9 2.19 30 1.63 1.25 30 19.5% 0.70 [0.18, 1.23]
Koshino 2023 1.38 1.54 16 2.31 2.44 16 15.3% -0.44 [-1.15, 0.26]
Kwon 2018 2.7 1.87 20 1.81 1.98 20 16.9% 0.45 [-0.18, 1.08]
Linens 2014 2.53 2.37 17 1.29 1.05 17 15.5% 0.66 [-0.03, 1.35]
Ross 2011 2.1 1.4 17 0.8 0.59 17 14.6% 1.18 [0.45, 1.92]
Subtotal (95% CI) 132 119 100.0% 0.47 [0.07, 0.86]
Heterogeneity: Tau² = 0.14; Chi² = 11.65, df = 5 (P = 0.04); I² = 57%
Test for overall effect: Z = 2.32 (P = 0.02)
1.1.3 SLS foam
Dobo 2015 5.84 1.37 32 3.95 0.71 19 24.7% 1.59 [0.94, 2.24]
Docherty 2006 7.07 1.6 30 5.6 1.77 30 27.9% 0.86 [0.33, 1.39]
Koshino 2023 7.13 1.89 16 6.38 2.25 16 23.4% 0.35 [-0.35, 1.05]
Linens 2014 6 1 17 5.59 1.33 17 24.0% 0.34 [-0.34, 1.02]
Subtotal (95% CI) 95 82 100.0% 0.80 [0.24, 1.35]
Heterogeneity: Tau² = 0.21; Chi² = 9.00, df = 3 (P = 0.03); I² = 67%
Test for overall effect: Z = 2.83 (P = 0.005)
1.1.4 TS firm
Dobo 2015 1.22 1.24 32 0.58 0.77 19 27.1% 0.58 [-0.00, 1.16]
Docherty 2006 1.23 1.25 30 0.53 1.04 30 33.9% 0.60 [0.08, 1.12]
Koshino 2023 1.44 2.06 16 1.56 1.9 16 19.0% -0.06 [-0.75, 0.63]
Linens 2014 1.29 1.53 17 1 1.17 17 20.0% 0.21 [-0.47, 0.88]
Subtotal (95% CI) 95 82 100.0% 0.39 [0.09, 0.69]
Heterogeneity: Tau² = 0.00; Chi² = 2.93, df = 3 (P = 0.40); I² = 0%
Test for overall effect: Z = 2.54 (P = 0.01)
1.1.5 TS foam
Dobo 2015 3.81 1.28 32 2.47 0.84 19 25.5% 1.16 [0.55, 1.77]
Docherty 2006 4.33 2.35 30 2.73 1.6 30 34.6% 0.79 [0.26, 1.31]
Koshino 2023 6.5 2.73 16 5.06 2.29 16 19.1% 0.56 [-0.15, 1.27]
Linens 2014 3.71 1.65 17 3.06 1.48 17 20.8% 0.40 [-0.28, 1.08]
Subtotal (95% CI) 95 82 100.0% 0.76 [0.45, 1.07]
Heterogeneity: Tau² = 0.00; Chi² = 2.99, df = 3 (P = 0.39); I² = 0%
Test for overall effect: Z = 4.79 (P < 0.00001)
Test for subgroup differences: Chi² = 14.55, df = 4 (P = 0.006), I² = 72.5%
-4
-2
0
2
4
Favours [CAI]
Favours [control]

Figure 3


CAI
Control
Std. Mean Difference
Std. Mean Difference
Study or Subgroup Mean SD Total Mean SD Total Weight IV, Random, 95% CI
IV, Random, 95% CI
1.2.1 Total score
Docherty 2006 15.7 6 30 10.7 3.2 30 26.9% 1.03 [0.49, 1.57]
John 2020 9.8 3.8 20 11.4 3.7 20 25.2% -0.42 [-1.05, 0.21]
Koshino 2023 16.9 4.5 16 15.6 5.8 16 23.9% 0.24 [-0.45, 0.94]
Linens 2014 13.59 4 17 11.06 3.01 17 24.0% 0.70 [0.00, 1.39]
Subtotal (95% CI) 83 83 100.0% 0.40 [-0.25, 1.05]
Heterogeneity: Tau² = 0.33; Chi² = 12.54, df = 3 (P = 0.006); I² = 76%
Test for overall effect: Z = 1.19 (P = 0.23)
1.2.2 Firm total score
Koshino 2023 2.5 2 16 3.9 3.9 16 51.7% -0.44 [-1.14, 0.26]
Powell 2014 4.55 2.45 10 1.18 0.98 10 48.3% 1.73 [0.67, 2.79]
Subtotal (95% CI) 26 26 100.0% 0.61 [-1.52, 2.73]
Heterogeneity: Tau² = 2.14; Chi² = 11.17, df = 1 (P = 0.0008); I² = 91%
Test for overall effect: Z = 0.56 (P = 0.58)
1.2.3 Foam total score
Koshino 2023 14.4 3.7 16 11.7 3.4 16 58.8% 0.74 [0.02, 1.46]
Powell 2014 7.98 1.78 10 4.46 2.23 10 41.2% 1.67 [0.62, 2.72]
Subtotal (95% CI) 26 26 100.0% 1.12 [0.23, 2.02]
Heterogeneity: Tau² = 0.22; Chi² = 2.05, df = 1 (P = 0.15); I² = 51%
Test for overall effect: Z = 2.46 (P = 0.01)
Test for subgroup differences: Chi² = 1.66, df = 2 (P = 0.44), I² = 0%
-4
-2
0
2
4
Favours [CAI]
Favours [control]

Figure 4


CAI
Control
Std. Mean Difference
Std. Mean Difference
Study or Subgroup Mean SD Total Mean SD Total Weight IV, Fixed, 95% CI
IV, Fixed, 95% CI
Hiller 2007 22.6 3.29 31 13.3 2.3 15 22.3% 3.04 [2.14, 3.93]
Linens 2014 5.57 2.38 17 3.2 2.68 17 35.4% 0.91 [0.20, 1.62]
Wikstrom 2017 12.3 6.08 19 9.3 4.09 19 42.3% 0.57 [-0.08, 1.22]
Total (95% CI) 67 51 100.0% 1.24 [0.82, 1.66]
Heterogeneity: Chi² = 20.41, df = 2 (P < 0.0001); I² = 90%
Test for overall effect: Z = 5.75 (P < 0.00001)
-4
-2
0
2
4
Favours [CAI]
Favours [control]

Figure 5


CAI
Control
Std. Mean Difference
Std. Mean Difference
Study or Subgroup Mean SD Total Mean SD Total Weight IV, Fixed, 95% CI
IV, Fixed, 95% CI
Chrints 1991 6.69 9.04 29 24.66 16.36 41 45.4% -1.29 [-1.81, -0.76]
Linens 2014 28.99 17.3 17 46.01 19.64 17 24.7% -0.90 [-1.61, -0.19]
Wikstrom 2017 9.83 7.02 19 14.55 12.94 19 29.9% -0.44 [-1.09, 0.20]
Total (95% CI) 65 77 100.0% -0.94 [-1.29, -0.59]
Heterogeneity: Chi² = 3.96, df = 2 (P = 0.14); I² = 50%
Test for overall effect: Z = 5.21 (P < 0.00001)
-4
-2
0
2
4
Favours [control]
Favours [CAI]