HOKKAIDO UNIVERSITY

Title	Genotypes and virulence-related activities of Candida albicans derived from oral cavity of patients in Hokkaido
Author(s)	Ouchi, Chisato; Hasebe, Akira; Sakata, Ken-ichiro et al.
Citation	Archives of oral biology, 157 https://doi.org/10.1016/j.archoralbio.2023.105827
Issue Date	2024-01
Doc URL	https://hdl.handle.net/2115/93315
Rights	©2024. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	Ouchi AOB Suppl Fig 1.R2.pdf, Supplementary material

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Supplementary Material




Figure S1. Measurement of ESE activities. (A) A typical picture of the agar plate for the measurement of ESE activity. Large white colonies of C. albicans and the destained or precipitated zones around them caused by ESE activity were observed. (B) The diameter of the colonies (denoted as a) and the diameter of the destained or precipitated zones (denoted as b) were measured. The Pz values were calculated as a ÷ b.