



HOKKAIDO UNIVERSITY

Title	New Species of Insects from Japan described during the Year 1935
Citation	Insecta matsumurana, 10(4), 163-173
Issue Date	1936-06
Doc URL	https://hdl.handle.net/2115/9332
Type	departmental bulletin paper
File Information	10(4)_p163-173.pdf



noted that "from the short ULMER's description of a female (*A. amurensis*) from Japan it may be conclude, that this female belongs not to *A. amurensis*, but probably, to *A. sachalinensis*." On examination of a large series of the examples, I come to express the opinion that the separation between both species is quite unstable. So, in this paper, I reserve the name *A. amurensis* on the Japanese examples.

Trivial Name: *Amur-tobikera*.

NEW SPECIES OF INSECTS FROM JAPAN DESCRIBED DURING THE YEAR 1935*

HYMENOPTERA

MICKEL, C. E.: The Mutillid Wasps of the Islands of the Pacific Ocean (Hymenoptera; Mutillidae) (Trans. Ent. Soc. London, LXXXIII, pp. 177-312).

Cystomutilla (?) *teranishii*, p. 196; *Smicromyrme lewisi lewisi* (nom. n.), p. 288; *S. lewisi nigricula* (n. subsp.), p. 289; *S. lewisi yanoi* (n. subsp.), p. 289.

MIWA, Y. et SONAN, J.: Taiwansan Hariganemushi ni kisei suru Arigatabachi no 1 Shinshu (Description of a new species of the Genus *Pristocera* parasitic on the larvae of Elateridae from Formosa) (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 90-92).

Pristocera formosana, p. 91.

—————: Description of a new Egg-parasite of *Melanauster chinensis* FORST. from Formosa (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 406-407).

Aprostoctus fukutai, p. 406.

SONAN, J.: Descriptions of two new Species of the Family Vespidae (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 370-372).

Vespa matsumurai, p. 370; *V. esakii*, p. 371.

YASUMATSU, K.: The Genus *Pison* SPINOLA of the Japanese Empire (Hymenoptera, Trypoxylonidae) (Ann. Zool. Japon, XV, pp. 227-238).

Pison strandi, p. 231; *P. iwatai*, p. 233; *P. tosawai*, p. 234.

—————: Notes on some Hymenoptera collected by Mr. C. TAKEYA on Sado Island with Descriptions of two unrecorded *Megachile*-species from Japan and Amami-Oshima Island (Fukuoka Hakubutsu. Zasshi, I, No. 6, pp. 384-389).

Megachile (*Chelostomoides*) *esakii*, p. 387.

—————: Two new Eumenidae from Tsushima and Formosa (*Mushi*, VIII, pp. 86-89).

Rhynchium flavopunctatum SMITH f. *tsushimarum* (n. f.), p. 86; *Ancistrocerus esakii*, p. 86.

—————: The Oxybelidae of Japan and Korea (Hymenoptera) (Trans. Sapporo Nat. Hist. Soc., XIV, pp. 38-41).

Oxybelus strandi, p. 39.

* Insects described in this magazine are omitted.

DIPTERA

- ALEXANDER, C. P.: New or little-known Tipulidae from Eastern Asia (Diptera), XXIII (Philip. Journ. Sci., LVI, pp. 339-369).
- Trichocera arisanensis, p. 339; Longurio (Longurio) variceps, p. 340; Tipula (Acutipula) albo-plagiata, p. 341; T. gressitti, p. 348; Nephrotoma atrolatera, p. 350; Dolichozepeza (Nesopeza) lugubri-vestis, p. 351; Limonia (Limonia) subhostilis, p. 352; Dicranota (Rhaphidolabis) pallidithorax, p. 356; D. (Amalopina) fumicostata, p. 357; Styrgomyia separata, p. 368.
- _____ : l. c., XXIV (Philip. Journ. Sci., LVI, pp. 525-562).
- Trichocera sapporensis, p. 525; Tipula (Oreomyza) coxitalis, p. 532; T. (Oreomyza) stenotuber-culata, p. 534; T. (Schummelia) strictiva, p. 541; Limonia (Libnotes) immetata, p. 546; L. (Limonia) tuta, p. 547.
- _____ : l. c., XXVI (Philip. Journ. Sci., LVII, pp. 195-225).
- Nephrotoma definita, p. 202; N. caudifera, p. 203; Hexatoma (Eriocera) ishigakiensis, p. 212; H. (Eriocera) iriomotensis, p. 213; Teucholabis (Teucholabis) iriomotensis, p. 218; Gonomyia (Ptilostena) longipennis, p. 219; G. (Gonomyia) bibarensis, p. 220; G. (Gonomyia) foliacea, p. 221.
- _____ : l. c., XXVII (Philip. Journ. Sci., LVIII, pp. 213-252).
- Tipula (Arctotipula) miyadaii, p. 220; T. (Schummelia) stipata, p. 221; Nephrotoma integra, p. p. 224; N. sublamellata, p. 231; N. chosensis, p. 233; N. daisensis, p. 236; Limonia (Libnotes) perparvuloides, p. 241; L. (Limonia) edura, p. 242; L. (Limonia) juvenca, p. 243; L. (Pseudoglochina) monocycla, p. 244; L. (Pseudoglochina) dimelania, p. 246; Antocha (Antocha) platyphallus, p. 247; Gonomyia (Gonomyia) latifolia, p. 248; G. (Lipophleps) praedita, p. 249.
- _____ : l. c., XXVIII (Philip. Journ. Sci., LVIII, pp. 385-423).
- Ormosia seclusa, p. 422.
- ENDERLEIN, G.: Eisentrautius ibizanus nov. gen., nov. spec. und über weitere Astiiden (Dipt.) (Mitt. Deutsch. Ent. Ges., VI, pp. 46-47).
- Chaetastia (n. g.) for C. sexsetosa (DUDA 1927), p. 47.
- KARIYA, S.: On the Family Tanyderidae of Japan (Mushi, VIII, pp. 39-41).
- Protanyderus alexanderi, p. 40.
- KARL, O.: Aussereuropäische Musciden (Anthomyiden) aus dem Deutschen Entomologischen Institut (Arb. Morph. Taxon. Ent. Berlin-Dahlem, II, pp. 29-49).
- Stomoxys latipennis, p. 30; Hydrotaea affinis, p. 38; Pegomyia quadrivittata, p. 44; Pygophora trimaulata, p. 48.
- SASAKI, C.: On a new Phorid-fly infesting our Edible Mushroom (Proc. Imp. Acad. Japan, XI, pp. 112-114).
- Ophiochaeta matsutakei, p. 112.
- SHIRAKI, T.: Simuliidae of the Japanese Empire (Mem. Fac. Sc. Agr. Taihoku Imp. Univ., Formosa, XVI, pp. 1-90).
- Prosimulium yezoense, p. 3; P. kiotoense, p. 6; Helodon sapporensis, p. 10; Eusimulium falcoe, p. 13; E. taipei, p. 15; E. geniculare, p. 19; E. bonninense, p. 21; Stilboplax 5-striatum, p. 27; Odagmia bidentata, p. 34; O. oitana, p. 37; O. iwatensis, p. 40; O. karenkoensis, p. 42; O. japonica, p. 45; Simulium annulipes, p. 49; S. katoi, p. 53; S. oshimanum, p. 56; S. nipponese, p. 59; S. minutum, p. 62; S. ambignum, p. 71; S. yokotense, p. 74; S. nikkoense, p. 77; S. arisanum, p. 80.
- TOKUNAGA, M.: Chironomidae from Japan, V. Supplementary Report on the Clunioninae (Mushi, VIII, pp. 1-20).
- Telmatogeton pacificus, p. 15.
- _____ : Three new Crane-flies from Japan (Ann. Zool. Japon, XV, pp. 194-198).
- Limonia (Idioglochina) pacifica, p. 194; Limmophila nigrilunae, p. 196; Alfredia imanishii, p. 198.

TOKUNAGA, M. et KURODA, M.: Unrecorded Chironomid Flies from Japan (Diptera), with a Description of a new Species (Trans. Kansai Ent. Soc., No. 6, pp. 1-8).
Chironomus (Stenochironomus) nelumbus, p. 4.

COLEOPTERA

ADACHI, T.: New Staphylinid Beetles from Japan (Studies on Staphylinidae of Japan I) (Kontyû IX, pp. 125-182).

Coprophilus fulvomaculatus, p. 125; *Deleaster yokoyamai*, p. 126.

BERNHAEUER, M.: Neuheiten der paläarktischen Staphylinidenfauna I (Koleopt. Rund., XXI, pp. 39-48).

Pinophilus sachtlebeni, *P. sautteri*, *P. formosae*, p. 42; *Paederus jessoensis*, p. 44; *Leucoparyphus silphoides* L. var. *koreanus* (n. var.), p. 48.

CHUJÔ, M.: Chrysomelidae of Loo-Choo Archipelago (I) (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 65-89).

Pedrilla flavonotata, p. 69; *Cryptocephalus loochooensis*, p. 72; *C. takahashii*, p. 73; *Coenobius obscuripennis*, p. 74; *Lamprosoma okinawense*, p. 75; *Chrysolampa fulva*, p. 76; *Nodostoma hirayamai*, p. 77; *N. okinawense*, p. 77; *Aulacophora (Ceratia) nitidipennis*, p. 82; *Theopea aureoviridis*, p. 85.

—————: l. c. (II) (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 203-211).

Lypesthes ungeri, p. 204; *Luperodes flavomaculatus*, p. 206; *Spaheroderma quadrimaculata*, p. 207.

—————: Studies on the Chrysomelidae in the Japanese Empire (VIII), Subfamily Halticinae (1) (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 354-369).

Nonarthra coreanum, p. 359; *N. formosense*, p. 360; *Psylliodes taiwanica*, p. 364; *P. gigantea*, p. 367.

—————: l. c., Subfamily Halticinae (2) (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 392-400).

Sangaria costata, p. 394; *Liprus montanus*, p. 396; *Lipromima fulvipes* (n. sp.) & ab. *bicolor* (n. ab.), p. 399.

—————: l. c., Subfamily Halticinae (3) (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 459-476).

Chaetocnema formosensis, p. 471.

—————: H. SAUTER'S Formosa-Ausbeute: Subfamily Galerucinae (Coleoptera, Chrysomelidae) (Arb. Morph. Taxon. Ent. Berlin-Dahlem, II, pp. 160-174).

Luperodes nitidissimus, p. 161; *Luperus (Luperus) sauteri*, p. 162; *Paraulaca costata*, p. 164; *P. flavipennis*, p. 165; *P. sauteri*, p. 166; *P. taiwana*, p. 167; *Galerucida nigrita*, p. 168; *Theopea sauteri*, p. 169, *T. sauteri*, var. *minuta* (n. var.), p. 171; *Aenidea taiwana*, p. 171; *Monolepta formosana*, p. 172; *M. sauteri*, p. 173.

—————: Description of a new Genus and two new Species of Chrysomelidae (Col.) (Arb. Morph. Taxon. Ent. Berlin-Dahlem, II, pp. 219-220).

Eriosardella (n. g.) *costata*, p. 219; *Cryptocephalus kraatzi*, p. 220.

—————: Descriptions of three new Mordellid-species from Formosa (Sylvia, VI, pp. 72-74).

Mordella auromaculata, p. 72; *Tomoxia formosana*, p. 73; *Mordellistena nigrofasciata*, p. 74.

—————: A systematic Catalogue of Formosan Phipiphoridae (Coleoptera) (Sylvia, VI, pp. 129-135).

Macrosiagon uninotaticolle, p. 133.

GRESSITT, J. L.: New Species and Records of Longicorns from Formosa (I) (Coleoptera: Cerambycidae) (Philip. Journ. Sci., LVII, pp. 181-194).

Noemia incompta, p. 181; *Aglaophis decemmaculatus*, p. 185; *Cleomenida pulchella*, p. 187;

Monochamus flocculatus, p. 188; *Glenea luteicollis*, p. 190; ? *Glenea lata*, p. 191; *Anastathes parva*, p. 193.

GRESSITT, J. L.: New Longicorn Beetles from Formosa (II) (Philip. Journ. Sci., LVIII, pp. 253-266).

Pseudopidonia aenipennis, p. 253; *P. subaenus*, p. 254; *P. occipitalis*, p. 255; *Leptura* (?) *ochraceoventra*, p. 256; *Anoplodera* (?) *tricolor*, p. 258; *Anoplodermorpha binotata*, p. 259; *A. breva*, p. 260; *Strangalina angustissima*, p. 261; *S. sturalis*, p. 262; *S. subapicalis*, p. 263; *Leptepania minuta*, p. 265.

—————: New Longicorn Beetles of the Subfamily Lamiinae from Formosa (Coleoptera: Cerambycidae) (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 286-292).

Exocentrus rufithorax, p. 286; *Neoserixia longicollis*, p. 287; *N. longicollis infasciatus*, p. 289; *N. schwarzeri*, p. 289; *Serixia albopleura*, p. 291.

—————: New Japanese Longicorn Beetles (Coleoptera, Cerambycidae) (Kontyû, IX, pp. 166-179).

Pachypidonia (n. g.), p. 166; *P. crassicornis*, p. 168; *Leptura stygica*, p. 168; *Stenomachus lighti*, p. 170; *Phymatodes quadrimaculatus*, p. 171; *P. vandykei*, p. 172; *Epicyltus* (n. g.) for *Clytus yokoyamai* KANO, p. 173; *Exocentrus fisheri*, p. 175; *Praolia yuasa*, p. 176; *Phytoecia gleneoides*, p. 177.

KOBAYASHI, K. et SEKI, K.: A new Cerambycid Beetle from Honshû, Japan (Mushi, VIII, pp. 42-43).

Menesia tokioensis, p. 42.

KÔNO, H.: Die Mordelliden Japans. Fünfter Nachtrag (Trans. Sapporo Nat. Hist. Soc., XIV, pp. 123-130).

Higehananomia (n. g.) *palpalis*, p. 123; *Yakuhananomia* (n. g.) for *Tomoxia yakui* KÔNO, p. 124; *Hoshihananomia* (n. g.), p. 124; *H. pirika*, p. 125; *Mordellistena okamotoi*, p. 127; *M. shizuokana*, p. 128.

KORSCHESKY, R.: Neue Coccinelliden aus Afrika, Brasilien und Formosa (Arb. Morph. Taxon. Ent. Berlin-Dahlem, II, pp. 252-256).

Rodolia formosana, p. 255.

MASUDA, T.: *Doromushi-ka no 1 Shinshu* (A new Dryopid-Beetle, *Psephenoides japonicus* sp. nov.) (Trans. Kansai Ent. Soc., No. 6, pp. 9-10).

Psephenoides japonicus, p. 9.

MATSUSHITA, M.: Bemerkungen zu den japanischen Cerambyciden nebst Beschreibung einiger neuer Arten (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 308-313).

Leptura (*Vadonia*) *kochiana*, p. 308; *Strangalia* (s. str.) *shikokensis*, p. 399; *Japonopsimus* (n. g.), p. 310; *Uraecha angusta* PASCOE subsp. *horishana* (n. subsp.), p. 312; *Mesosella kuraruana*, p. 312; *Exocentrus saitoi*, p. 313.

—————: Beitrag zur Cerambyciden-Fauna von Mikronesien (Trans. Sapporo Nat. Hist. Soc., XIV, pp. 115-122).

Rhaphipodus (s. str.) *carolinensis*, p. 115; *Ceresium yoshinoi*, p. 116; *C. nanyoanum*, p. 117; *Nanyohammus* (n. g.), p. 117; *N. luteoparsus*, p. 118; *Niphohammus auripilis*, p. 119; *Pterolophia palauana*, p. 119; *P. uchiyamai*, p. 120; *Sybra carolina*, p. 121.

MITONO, T.: Descriptions of new Species of Longicorn Beetles from Formosa II (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 404-405).

Oplatocera mandibulata MIWA et MITONO (n. sp.), p. 404.

MIWA, Y.: Description of a new Species of Hydroscaphidae from Formosa (II) (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 352-353).

Hydroscapha takahashii, p. 352.

—————: Descriptions of two new Species of Lymexylonidae from Formosa (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 456-458).

- Hylecoetus formosanus, p. 456; Atractocerus mirabilis, p. 457.
- MIWA, Y.: Coleoptera from Amami-Islands in Loo-Choo Archipelago (Trans. Kansai Ent. Soc., No. 6, pp. 11-30).
- Pachylanguria oshimana, p. 17; Anadatus amamiensis, p. 17, Setenis oshimanus, p. 21.
- MIWA, Y. et CHUJŌ, M.: Some new Buprestids from the Japanese Empire (Entomological World, Tokio, III, pp. 270-282).
- Chalcophora satsumae var. abnormale (n. var.), p. 270; Poecilonota cupreomaculata, Eurythyrea kawakamii, p. 271; Buprestis japonensis sachalinensis (n. subsp.), B. unicus, p. 272; B. aurulenta niponica (n. subsp.), p. 273; Tamamushia (n. g.), p. 274; T. virida, Anthaxia proteus var. minuta (n. var.), A. proteus var. matsumurai (n. var.), p. 275; A. proteus var. rubromarginata (n. var.), A. proteus var. viridomarginata (n. var.), Chrysobothris shirakii, p. 276; Coraebus mushaensis, p. 277; C. pruni, C. sonani, p. 278; C. formosanus, p. 279; C. insulicora var. rubrifrons (n. var.), Sambus formosanus, p. 280; S. quiquefasciatus, Cylindromorphus formosanus, p. 281.
- MURAYAMA, J.: Notes au point de vue scientifique, sur les espèces nonreconnues, rares et nouvelles de Scarabéidés de la Corée (Journ. Chosen Nat. Hist. Soc., XX, pp. 1-9).
- Serica septentrionalis, p. 7; S. koryoensis, p. 8.
- PIC. M.: Nouveautés diverses (Mélanges exot.-ent., LXVI, pp. 1-36).
- Byrrhinus formosanus, p. 2; Podabrinus obscuricolor, p. 12.
- SCHEDL, K. E.: Scolytidae and Platypodidae: New Species from the Philippine Islands and Formosa (Philip. Journ. Sci., LVII, pp. 479-489).
- Xylechinus formosanus, p. 479.
- SEKI, K.: Two new Longicorn-Beetles from Japan (Kontyû, IX, pp. 91-93).
- Stenothalmus koshunensis, p. 91; Xylotrechus nipponicus, p. 92.
- : Description on a new Longicorn-Beetle from Japan (Entomological World, Tokio, III, pp. 224-225).
- Clytus fukukii, p. 224.
- : Notes on the Cerambycid-Coleoptera of the Genus Psacotha, with Description of one new Variety (Entomological World, Tokio, III, pp. 289-293).
- Psacotha hilaris var. machidai (n. var.), p. 292.
- SEKI, K. et KOBAYASHI, K.: Description on a new Longicorn-Beetle from Japan (Entomological World, Tokio, III, pp. 225-227).
- Glenea kasugensis, p. 225.
- SEKI K. et SUEMATSU, T.: Descriptions of two new Longicorn-Beetles of the Tribe Clytini from Japan and Formosa (Ins. World, Gifu, XXXIX, pp. 5-7).
- Perissus arisanus, p. 5; Chlorophorus minomensis, p. 6.

LEPIDOPTERA

- DOI, H.: New or unrecorded Butterflies from Corea (Zephyrus, VI, pp. 15-16).
- Argynnis amathusia nansetsuzana (n. subsp.), p. 15.
- MEYRICK, E.: Exotic Microlepidoptera IV, pt. 18, pp. 545-576; pt. 19, pp. 577-608.
- Drosophantis (n. g.), p. 546; D. corusca, p. 546; Sympetalistis (n. g.), p. 556; S. petrograpta, p. 557; Thamnoscelis (n. g.), p. 558; T. prisciformis, p. 558; Mnesipatris (n. g.) phaedrospora, p. 559; Thiotricha dissobola, p. 561; Cratinitis (n. g.) tubigera, p. 561; Phrixocrita (n. g.), p. 561; P. aegidopis, p. 562; Musurga oxycarpa, p. 562; Frisilia homalistic, p. 563; Thubana deltaspis, p. 563; Lecithocera fascinatrix, p. 563; Athymoris (n. g.) martialis, p. 564; Pachnistis silens, p. 565; Cynicocrates (n. g.), p. 565; C. tachytoma, p. 566; Satrapia pyrotechnica, p. 566; Borkhausenia tyropis, p. 566; Cryptolechia argometra, p. 567; Blastobasis indirecta, p. 567; Stagmatophora semioceros, p. 572; Balionebrs (n. g.) bacteriota, p. 573; Imma tricocota, p. 574; Tinea metathyris, p. 575; T.

limentis, p. 575; T. protograpta, p. 578; T. endochrysa, p. 579; T. trapezoides, p. 579; Ceratosticha (n. g.), p. 579; C. leptodeta, p. 580; Aristotelia cleodora, p. 583; Stenolechia bathrodyas, p. 583; S. notomochla, p. 583; Telphusa comprobata, p. 584; Phthorimaea aganocarpa, p. 585; Thiotricha microrrhoda, p. 586; Chelaria triorthias, p. 589; Promalactis autoclina, p. 592; P. symbolopa, p. 593; Depressaria spectrocentra, p. 593; Commatarcha (n. g.) palaeosema, p. 594; Lithocolletis tritorrhecta, p. 596; Epicephala venenata, p. 596; Hyponomeuta mochlocrossa, p. 602; H. hexabola, p. 602; H. spodocrossa, p. 602; Ethmia mesozyga, p. 603; Rhabdocosma (n. g.), p. 603; R. aglaophanes, p. 604; Labdia bitabulata, p. 605; Chrysoclista hexachrysa, p. 607.

MORI, T. et CHO, F.: Description of a new Butterfly and two interesting Butterflies from Korea (Zephyrus, VI, pp. 11-14).

Erebia shajitsuzanensis, p. 11.

NAGAMI, K. et ISHIGA, T.: Tottorisan Chō-ūi no 1 Shinshu (A new Species of Butterfly from Tottori) (Fukuoka Hakubutsu. Zasshi, I, pp. 303-304).

Zephyrus hisamatsusanus, p. 303.

NAKAHARA, W.: A remarkable aberrant Form of Hebomoia glaucippe from Formosa (Zephyrus, VI, pp. 20-21).

Hebomoia glaucippe formosana FRUHSTORFER ab. dortheae (n. ab.), p. 20.

NOMURA, K.: Notes on some Butterflies of the Genus Neptis from Formosa and Corea (Zephyrus, VI, pp. 25-41).

Neptis esakii, p. 29; Neptis themis nirei (n. subsp.), p. 31.

ŌKUBO, J.: On a new Subspecies of Lycaena pryeri MURRAY from Shikoku (Zephyrus, VI, pp. 45-50).

Lycaena pryeri shikokuana (n. subsp.), p. 45.

SONAN, J.: Psychidae of Formosa (Trans. Nat. Hist. Soc. Formosa, XXV, pp. 448-455).

Kotochallia (n. g.) shirakii, p. 449; Mahasena oolona, p. 450; M. kotoensis, p. 451; Acanthopsyche (Pteronema) postica, p. 452; A. (Metisa) saccharivora, p. 453; Psyche (Eurycyttarus) takahashii, p. 454.

WATARI, M.: On some Species of the Genus Zephyrus (Zephyrus, VI, pp. 42-44).

Zephyrus saphirina STAUDINGER ab. immaculata (n. ab.), p. 42.

RHYNCHOTA

ESAKI, T.: A new Species of Cicadidae from Formosa (Proc. Imp. Acad. Japan, XI, pp. 201-202).

Tibicen chujoi, p. 201.

—————: New Henicocephalidae from Japan and Formosa (Mushi, VIII, pp. 21-24).

Henicocephalus yami, p. 21; H. chipon, p. 22; H. japonicus, p. 23.

—————: New name for Dikraneura dorsalis ESAKI, 1931. (Mushi, VIII, p. 24).

Dikraneura fragi (nom. nov.) for Dikraneura dorsalis.

—————: A new Species of Scutellerinae from Japan (Mushi, VIII, pp. 105-107).

Poecilocoris splendidulus, p. 105.

KANDA, S.: Descriptions of new Pseudococcus (Coccidae) from Japan (Kontyū, IX, pp. 83-90).

Pseudococcus flavidus, p. 83; P. matsudoensis, p. 85; P. sasae, p. 87.

—————: On the Genus Heliococcus (Coccidae) (Ann. Zool. Japon., XV, pp. 70-76).

Heliococcus takahashii, p. 73.

KATO, M.: Description on a new Membracidae from Chosen (Entomological World, Tokio, III, pp. 476-478).

Neomachaerotypus (n. g.) eguchii, p. 476.

SHUNJI, O.: Hōsan Higemadara-abura Agriopaphis-Zoku no Shu no Kensakuhyō to 3 Shinshu

(A Key to the Japanese Species of the Genus *Agrioaphis* with Descriptions of three new Species) (*Oyo-Dobutsugaku-Zasshi*, VII, pp. 281-287).

Agrioaphis morioka, p. 282; *A. hashibamii*, p. 284; *A. sasacola*, p. 285.

SHINJI O.: Tōhokusan Hadaka-kaigaramushi no 2 Shinshu (Descriptions of two new Species of Coccidae from Tōhoku, Japan) (*Oyo-Dobutsugaku-Zasshi*, VIII, pp. 288-290).

Monophlebus akebiae, p. 288; *Lecanium gigantea*, p. 289.

—————: 1 Shinka o mooku beki Hadaka-kaigaramushi no 1 Shu ni tsuite (A new Genus and Species of Coccidae) (*Oyo-Dobutsugaku-Zasshi*, VII, pp. 106-108).

Serrolecanium (n. g.) *bambusae* p. 106.

—————: Naichisan Kobuabura-zoku no Shu no Kensaku-hyō to 1 Shinshu (A Key to the Japanese Species of the Genus *Myzus* with Description of one new Species) (*Kontyū*, IX, pp. 233-238).

Myzus geranicola, p. 234.

—————: Naichisan Higenagaabura-zoku no Shu no Kensakuhyō to 1 Shinshu (A Key to the Japanese Species of *Macrosiphum* with Description of one new Species) (*Kontyū*, IX, pp. 239-243).

Macrosiphum maazamii, p. 241.

—————: Naichisan Aminashihigenagaabura-zoku no Shu no Kensakuhyō to 1 Shinshu (A Key to the Japanese Species of the Genus *Acyrtosiphum* with Description of one new Species) (*Kontyū*, IX, pp. 249-253).

Acyrtosiphum circifoliae, p. 251.

—————: Hosan Aūra-zoku no Kensakuhyō to 1 Shinshu no Kisai (A Key to the Japanese Species of the Genus *Aphis* with Description of one new Species) (*Dobutsu. Zasshi*, XLVII, pp. 737-741).

Aphis sanguisorbae, p. 740.

—————: A List of the Unarmed Scale Insects collected in the Prefecture of Iwate, with the Descriptions of 7 new Species (*Dobutsu. Zasshi*, XLVII, pp. 767-776).

Pluvinaria araliae, p. 771; *P. euonymii*, p. 771; *P. katurae*, p. 772; *P. acericola*, p. 772; *Pseudococcus hibae*, p. 773; *P. katurae*, p. 773; *Lecanium sansho*, p. 773.

SHIRAIWA, H.: Studies on Mealy Bugs infesting Pear in Japan (*Kontyū*, IX, 64-75).

Pseudococcus matsumotoi, p. 66 & 68; *P. piricola*, p. 67 & 69.

SONAN, J.: Phymatidae of Formosa (*Trans. Nat. Hist. Soc. Formosa*, XXV, pp. 401-403).

Amblythyreus taiwana, p. 401; *Cnizocoris shirakii*, p. 402.

TAKAHASHI, R.: Scale Insects of the Genus *Coccoxymytilus* from Formosa (Hemiptera) (*Trans. Nat. Hist. Soc. Formosa*, XXV, pp. 444-447).

Coccoxymytilus kanoi, p. 444.

—————: Some Oriental Aphididae (Hemiptera). I, (*Konowia*, XIV, pp. 318-322).

Macrosiphum holsti, p. 318; *Prociphilus formosanus*, p. 320.

—————: Three new Aphids of the Genus *Thoracaphis* Van der Goot (Hem.) (*Stylops*, IV, pp. 86-89).

Thoracaphis piyananensis, p. 86; *T. saramaensis*, p. 88; *T. fici* TAKAHASHI var. *foveolatus* (n. var.), p. 89.

—————: Observations on the Coccidae of Formosa, Part V (Dept. Agr. Govern. Res. Inst. Formosa, No. 66, pp. 1-37).

Paracardiococcus (n. g.) *actinodaphnis*, p. 6; *Saissetia bobuae*, p. 8; *Aulacaspis kuzunoi* KUWANA var. *divergens* (n. var.) p. 10; *A. mangiferae* NEWST. var. *aceris* (n. var.), p. 12; *Diaspis machilicola*, p. 13 & var. *cinnamomicola* (n. var.), p. 15; *Phenacaspis rotunda*, p. 16; *P. taiwana*, p. 17; *Leucodiaspis vitis*, p. 19; *Lepidosaphes takaoensis*, p. 23; *L. cycadicola* KUWANA var. *yamahoi* (n. var.), p. 26; *Fiorinia rhododendri*, p. 27; *Parlatoria lithocarpi*, p. 29; *Aspidiotus shakunagi*, p. 32; *A. chipponsanensis*, p. 33; *Aonidia tentaculata*, GREEN var. *formosana* (n. var.), p. 35.

—————: Aleyrodidae of Formosa, Part IV (Dept. Agr. Govern. Res. Inst. Formosa, No.

66, pp. 39-64).

Dialeurodes subrotunda, p. 41; *D. citri* ASHM. var. *kinyana* (n. var.), p. 43; *D. agalmae*, p. 44; *D. elaeagni*, p. 45; *D. vanieriae*, p. 46; *D. ficicola*, p. 47; *D. tristylis*, p. 48; *D. ardisiae*, p. 50; *Aleurotuberculatus rhododendri*, p. 51; *A. piperis*, p. 52; *A. multipori*, p. 53; *Tuberaleyrodes machili* TAKAHASHI var. *actinodaphnis* (n. var.), p. 56; *Aleurotrachelus maesae*, p. 57; *A. pyracanthae*, p. 58; *Pealius psychotriae*, p. 59; *P. damnacanthi*, p. 61; *P. machii*, p. 62; *Aleurolobus hederæ*, p. 63.

—————: Notes on the Aleyrodidae of Japan (Homoptera), II (Kontyû, IX, pp. 25-27).

Acanthobemisia (n. g.) *distylis*, p. 25.

—————: I. c., III (Kontyû, IX, pp. 279-283).

Pealius rhododendri, p. 279; *Aleurocanthus woglumi* ASHBY var. *formosana* (n. var.), p. 281.

ORTHOPTERA

FURUKAWA, H.: On a small Collection of Orthopteroidea from Miyazaki Prefecture (Kyûshû) (Mushi, VIII, pp. 108-116).

Cycloptilum ohzii, p. 111.

SHIRAKI, T.: Orthoptera of the Japanese Empire, Part IV. Phasmidae (Mem. Fac. Sci. Agr. Taihoku Imp. Univ., XIV, No. 3, Entom., No. 6, pp. 23-88).

Baculum formosanum, p. 26; *B. granulatus*, p. 27; *Entoria ishigakiensis*, p. 32; *E. koshunensis*, p. 35; *E. okinawaensis*, p. 37; *E. miyakoensis*, p. 38; *E. banshoryoensis*, p. 40; *E. taihokuensis*, p. 41; *E. shinchikuensis*, p. 43; *E. nagoensis*, p. 47; *E. takaensis*, p. 51; *E. domonensis*, p. 53; *E. taitoensis*, p. 54; *E. longiopercula*, p. 57; *Phraortes formosanus*, p. 62; *P. koyasanensis*, p. 64; *P. kumamotoensis*, p. 66; *P. miyakoensis*, p. 68; *Micadina rotundata*, p. 72; *M. yasumatsui*, p. 74; *M. sonani*, p. 75; *Sipyloidea okunii*, p. 80; *S. truncata*, p. 82.

EPHEMERIDA

IMANISHI, K.: Mayflies from Japanese Torrents, V. Notes on the Genera *Cinygma* and *Heptagenia* (Ann. Zool. Japon., XV, pp. 213-221).

Cinygma hirasana, p. 214; *C. kurobensis*, p. 215; *C. adusta*, p. 217; *C. vernalis*, p. 217; *C. dorsalis*, p. 218.

THYSANOPTERA

PRIESNER, H.: New or little-known Oriental Thysanoptera (Philip. Journ. Sci., LVII, pp. 351-375).

Sericothrips tabulifer, p. 351; *Dendrothrips minowai*, p. 353; *Taeniothrips oreophilus*, p. 355; *Bolacothrips orientalis*, p. 359; *Liothrips heptapleurinus*, p. 360; *L. piperinus*, p. 361; *Dolichothrips pumilus*, p. 362; *Smerinthrips vitivorus*, p. 364; *Haplothrips* (*Odontoplothrips*) *dentifer*, p. 365; *H. chinensis* PRIESNER var. *montivage* (n. var.), p. 366; *H. allii*, p. 367; *Neosmerinthothrips formosensis*, p. 368; *Plectrothrips corticinus*, p. 371; *Elaphrothrips takahashii*, p. 372; *Leeuwenia pugnatix*, p. 373.

—————: Neue Exotische Thysanopteren (Stylops, IV, Part 6, pp. 125-131).

Paraliothrips (n. g.) *takahashii*, p. 125; *P. grandiceps*, p. 126; *Docdothrips imitans*, p. 127; *Taeniothrips congnaticeps*, p. 127; *T. alliorum*, p. 128.

TAKAHASHI, R.: An interesting Thrips from Amami-Oshima, Locchoos (Mushi, VIII, pp. 61-63). *Bactrothrips brevitubus*, p. 61.

SUPPLEMENT TO THE YEAR 1934*

HYMENOPTERA

SONAN, J.: A few Xylocopidae from Japan (Trans. Nat. Hist. Soc. Formosa, XXIV, pp. 226-232).

Xylocopa amamensis, p. 230.

YASUMATSU, K.: Les Hyménoptères de l'île Yakushima (Mushi, VII, pp. 61-67).

Henicospilus fuscomaculatus yakushimensis (n. subsp.), p. 67.

DIPTERA

BARANOV, N.: Uebersicht der orientalischen Gattungen und Arten des Carcelia-Komplexes (Tachinidae) (Trans. Ent. Soc. London, LXXXII, pp. 387-408).

Catacarcelia rondaniella, p. 392; *Eucarcelia* (n. g.) for *Tachina excisa* and *E. grossa*, p. 393; *Myxocarcelia* (n. g.) for *Carcelia hirsuta*, *Carceliella* (n. g.) for *Carcelia octava*, p. 398; *Parexorista latistylata*, p. 405; *Isocarceliopsis* (n. g.) *hemimacquartioides*, p. 406; *Asiocarcelia* (n. g.) for *Carcelia caudata*, p. 407.

—————: Neue Gattungen und Arten der orientalischen Raupen-fliegen (Encycl. Ent. Sér. B. II. Diptera, VII, pp. 160-165).

Trichiformosomyia (n. g.) *sauteri*, p. 164; *Kosempomyiella* (n. g.) *rufiventris*, p. 165.

CRESSON, JR. E. T.: Descriptions of new Genera and Species of the Dipterous Family Ephydriidae, XL (Trans. Amer. Ent. Soc., LX, pp. 199-222).

Hyadaina sauteri, p. 206; *Scatella bullacosta*, p. 219.

DUDA, O.: Weitere neue und wenig bekannte orientalische und australische Chloropiden (Diptera) des Deutschen Entomologischen Instituts in Berlin Dahlem (Arbeit. Morph. Taxon. Ent. Berlin-Dahlem, I, pp. 39-60).

Eutropha oldenbergi, p. 58.

EDWARDS, F. W.: Four new Diptera in the Collection of the Berlin Zoological Museum (Stylops, III, pp. 185-187).

Olbiogaster zonatus, p. 185.

HENDEL, F.: Revision der Tethiniden (Dipt. Muscid. acal.) (Tijdschr. Ent., pp. 37-54).

Rhinoessa orientalis, p. 47.

—————: Uebersicht ueber die Gattungen der Pyrgotiden, nebst Beschreibung neuer Gattung und Arten (Encycl. Ent. Sér. B, II, Diptera, VII, pp. 111-156).

Adapsilia facialis, p. 148.

SZIADY, Z.: Zwei neue oriental-asiatisch Rhagio-Arten (Dipt.) (Konowia, XIII, pp. 8-9).

Rhagio shirakii, p. 9.

COLEOPTERA

BALTHASAR, V.: Neue Coprinen-Arten und -Abarten (Ent. Blät., Berlin, XXX, pp. 146-149).

Copris frankenbergeri, p. 149.

BREUNING, S.: Ueber Carabini (Folia Zool. Hydrobiol., Riga, VIII, pp. 29-40).

Acoptopterus arrowianus, p. 31.

CHEN, S. II.: Coléoptères Halticinae recueillis par M. H. SAUTER à Formosa (Ann. Soc. Ent. France, CIII, pp. 175-185).

* Ins. Mats., IX, pp. 175-179 (1935).

- Eucycla sauteri*, p. 181; *Taizonia* (n. g.), p. 182; *T. bella*, p. 183; *Longitarsus horni*, p. 183; *L. taiwanicus*, p. 184; *Aphthona formosana*, p. 185.
- CHEN, S. H.: On some Species of Chrysomelidae in the Britisch Museum (Stylops, III, pp. 66-78). *Ambrostoma sublaevis*, p. 66; *Eucycla duodecimmaculata*, p. 75.
- CHUJÓ, M.: H. SAUTER'S Formosa-Ausbeute: Subfamilies Criocerinae, Clytrinae and Cryptocephalinae (Arbeit. Morph. Taxon. Ent., Berlin-Dahlem, I, pp. 281-291).
- Lema koshunensis*, p. 282; *Cyaniris* (*Cyaniris*) *japonicus* var. *formosana* (n. var.), p. 284; *C. (Cyaniris) sauteri*, p. 286; *Cryptocephalus formosanus* p. 287; *C. sauteri*, p. 289; *C. nitidissimus*, p. 290; *Melixanthus formosensis*, p. 291.
- : Description of a New *Temnaspis*-Species from Corea (Journ. Chosen Nat. Hist. Soc., XIX, p. 34).
- Temnaspis coreana*, p. 34.
- MURAYAMA, J.: Une nouvelle espèce de Scarabéididé de la Corée (Journ. Chosen Nat. Hist. Soc., XIX, p. 35).
- Aserica fusania*, p. 35.
- : Supplementary Notes on the Platypodidae of Formosa, IV (Journ. Facul. Agr. Hokkaido Imp. Univ., Sapporo, XXXV, pp. 133-149).
- Platypus arisannensis*, p. 135; *Crossotarsus emancipatus*, p. 138; *Diapus truncatus*, p. 143.
- PIC, M.: Notes diverses, nouveautés (Echange, Moulins, 50, pp. 17-19).
- Leptura bodemeyeri*, p. 17; *Strangalia circaocularis*, p. 18.
- : Neue exotische Coleopteren (Malacodermata) (Ent. Anz., XIV, pp. 45-47, and 54).
- Malthinus sauteri*, p. 46; *Themus kickeri*, p. 54.
- SCHAUBERGER, E.: Zur Kenntnis der indo-orientalischen Harpalinen (Dritter Beitrag) (Ent. Anz., XIV, pp. 9-13).
- Dioryche torta formosensis* (n. subsp.), p. 12.

LEPIDOPTERA

- DRAUDT, M.: The Macrolepidoptera of the World (in Seitz) Supplement III.
- Rhynchaglaea scitula albibasis*, p. 148.
- HIROSE, T.: On two new aberrant Forms in *Hypolimnas bolina jaluita* FRUHSTORFER from Marshall-Island (Trans. Kansai Ent. Soc., No. 5, pp. 25-28).
- Hypolimnas bolina jaluita* FRUHSTORFER ab. *marshallensis* (n. ab.), p. 25, ab. *ebonensis* (n. ab.), p. 26.
- KATO, M.: Description on a new Lycaenidae from Formosa (Entomological World, Tokio, II, pp. 386-388).
- Aphanaeus sozanensis*, p. 386.
- WAKABAYASHI, H.: Description on a new Form of *Papilio demetrius* from Tokio (Entomological World, Tokio, II, pp. 389-390).
- Papilio demetrius* CRAMER ab. *tokionis* (n. ab.), p. 389.
- WEHRLI, E.: Ueber neue paläarktische Geometrinae und ein neues Subgenus (Ins. Ent. Zeit., XXVII, pp. 509-513).
- Cystidia truncagulata*, p. 512.

RHYNCHOTA

- HEINZE, K.: Ueber Plataspidae (Hem.) aus Asien (Stett. Ent. Zeit., XCV, pp. 283-290).
- Ponsilasia* (n. g.), p. 283; *P. formosana*, p. 285.
- HUNGERFORD, H. B. et EVANS, N. E.: The Hydrometridae of the Hungarian National Museum

and other Studies in the Family (Hem.) (Ann. Mus. Nat. Hung., Budapest, XXVIII, pp. 31-112).

Hydrometra annamana, p. 68.

KANDA, S.: Two New Genera of Japanese Coccidae (Ins. World, Gifu, XXXVIII, pp. 308-312).

Saliococcus (n. g.) for *Phenacoccus takae* KUWANA, p. 308; *Rosanococcus* (n. g.) for *Phenacoccus suwakoensis* KUWANA et TOYODA, p. 311.

MONZEN, K.: Some Contributions to the Knowledges of the new and recorded Species of the Gall producing Aphides (Bull. Sci. Res. Alumni Ass. Morioka Coll., Agr. Forest., IX, pp. 13-34).

Nipponaphis distyliicola, p. 16; *N. globuli*, p. 17; *N. yanonis* MATS. var. *autumnus*, p. 21; *Anuraphis floris*, p. 33.

TAKAHASHI, R.: Another Aphid producing Galls on *Styrax* in Formosa (Hemiptera) (Trans. Nat. Hist. Soc. Formosa, XXIV, pp. 312-314).

Astegopteryx taiwana, p. 312.

—————: Two new Genera of Aphididae (Hemiptera) (Stylops, III, pp. 54-58).

Anomalosiphum (n. g.) *pithecolobii*, p. 54; *Taiwanaphis* (n. g.) *decasperimi*, p. 56.

THYSANOPTERA

PRIESNER, H.: Indomalayische Thysanopteren, V. Revision der indomalayischen Arten der Gattung *Haplothrips* SERV. (Record. Ind. Mus., XXXV, pp. 347-369).

Haplothrips vernoninal, p. 360, & var. *grandior* (n. var.), p. 361.