



HOKKAIDO UNIVERSITY

Title	Non-very generic intersections and their combinatorics [an abstract of dissertation and a summary of dissertation review]
Author(s)	Das, Pragnya
Degree Grantor	北海道大学
Degree Name	博士(理学)
Dissertation Number	甲第16083号
Issue Date	2024-09-25
Doc URL	https://hdl.handle.net/2115/93606
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Type	doctoral thesis
File Information	Pragnya_Das_abstract.pdf, 論文内容の要旨



Abstract of Doctoral Dissertation

Degree requested: Doctor of Science Applicant's name: Pragnya Das

Non-very generic intersections and their combinatorics

(ノンベリジェネリック交叉とその組合せ論)

The discriminantal arrangement $\mathcal{B}(n, k, \mathcal{A})$ defined by Manin and Schechtman is an arrangement of hyperplanes constructed from a generic arrangement $\mathcal{A}$ of n hyperplanes in a k -dimensional space generalizing the classical braid arrangement. In particular, $\mathcal{B}(n, 1, \mathcal{A})$ is the Braid arrangement.

The combinatorics of $\mathcal{B}(n, k, \mathcal{A})$ is known to be constant when $\mathcal{A}$ varies in an open Zariski set $\mathcal{Z}$ in the space of all generic arrangements. In particular $\mathcal{A}$ is called very generic if $\mathcal{A} \in \mathcal{Z}$ and non-very generic otherwise. In the later case, when $\mathcal{A}$ is a non-very generic arrangement, the combinatorics of the discriminantal arrangement $\mathcal{B}(n, k, \mathcal{A})$ can be studied by exploiting its relation with the special configurations of points in the space.

In this dissertation, we begin by reviewing the definition of a discriminantal arrangement and some basic properties of its combinatorics. We rewrite the orchard and the Sylvester's problems in terms of the combinatorics of $\mathcal{B}(n, 2, \mathcal{A})$ and then we build two arrangements of 12 lines in the projective real plane having, respectively, the minimum number of ordinary points and the maximum number of multiplicity 3 intersections. These arrangements of 12 lines in the projective real plane can help finding solutions for the orchard and the Sylvester's problems in a purely combinatorial way using the combinatorics of the intersection lattice of the discriminantal arrangements.

An intersection in $\mathcal{B}(n, k, \mathcal{A})$ when $\mathcal{A} \in \mathcal{Z}$ is called a very generic intersection and an intersection which doesn't exist in $\mathcal{B}(n, k, \mathcal{A})$ when $\mathcal{A} \in \mathcal{Z}$ is called a non-very generic intersection. We distinguish between the two types of non-very generic intersections as *arithmetic non-very generic* or *geometric non-very generic* considering whether the non-very generic intersection satisfies a condition (which we term as the BBA-condition) provided by Athanasiadis, Bayer and Brandth and then study the *arithmetic non-very generic intersections* which can be classified using the combinatorics of the discriminantal arrangement $\mathcal{B}(n, k, \mathcal{A})$.