



Title	Study on the therapeutic strategies of psoriasis targeting novel regulatory pathways in $\gamma\delta$ T cells [an abstract of dissertation and a summary of dissertation review]
Author(s)	王, 上一
Degree Grantor	北海道大学
Degree Name	博士(獣医学)
Dissertation Number	甲第16158号
Issue Date	2024-09-25
Doc URL	https://hdl.handle.net/2115/93825
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Type	doctoral thesis
File Information	Shangyi_Wang_abstract.pdf, 論文内容の要旨



学位論文内容の要旨
Abstract of the dissertation

博士の専攻分野の名称：博士（獣医学）

氏名：王 上一

Name

学位論文題名
The title of the doctoral dissertation

Study on the therapeutic strategies of psoriasis targeting novel
regulatory pathways in $\gamma\delta$ T cells

($\gamma\delta$ T 細胞の新規制御経路とこれを標的とした乾癬の治療戦略に関する研究)

< abstract >

Psoriasis is a chronic inflammatory skin disease characterized by epidermal hyperproliferation and cellular infiltration. Previous studies have shown that psoriasis development depends on proinflammatory cytokines, such as interleukin (IL)-23 and IL-17. IL-23 produced by innate immune cells, such as dendritic cells and macrophages, in psoriatic lesions activates a subset of helper T cells to release IL-17, promoting neutrophil recruitment and epidermal keratinocyte proliferation. However, recent studies have revealed that innate $\gamma\delta$ T cells are involved in psoriasis pathogenesis as the primary source of IL-17. The purpose of the present study was to unravel the regulatory mechanisms of IL-17-producing $\gamma\delta$ T ($\gamma\delta$ T17) cells and establish the basis of psoriasis therapies targeting $\gamma\delta$ T17 cells.

In Chapter 1, $\gamma\delta$ T17 cell regulation through nuclear receptors, REV-ERBs, and the therapeutic strategy of psoriasis targeting this regulatory pathway were shown. REV-ERBs are ligand-dependent transcription factors and generally known to regulate circadian rhythm. Although a recent study revealed that REV-ERBs suppress the function of IL-17-producing helper T cells, the role of REV-ERBs in $\gamma\delta$ T17 cells regulation had been unclear. $\gamma\delta$ T17 cells were significantly increased in the secondary lymphoid organs of REV-ERB-deficient mice. IL-17 production by $\gamma\delta$ T cells was suppressed by short-term treatment with a synthetic REV-ERB agonist, SR9009, both *in vitro* and *in vivo*. Furthermore, topical application of SR9009 to the skin reduced the inflammatory symptoms of imiquimod (IMQ)-induced psoriasiform dermatitis in mice. These results demonstrate

that $\gamma\delta$ T17 cells associated with psoriasis development are regulated by REV-ERBs and that modulation of REV-ERB activity in $\gamma\delta$ T17 cells using synthetic ligands may become a novel therapeutic approach for psoriasis.

Next in Chapter 2, the role of an inflammatory cytokine called tumor necrosis factor-like ligand 1A (TL1A) and its receptor, death receptor 3 (DR3), in $\gamma\delta$ T17 cell activation and its relevance with psoriasis pathogenesis were examined. Neutralization of TL1A by anti-TL1A antibody injection inhibited the cytokines production by $\gamma\delta$ T17 cells in IMQ-treated mouse skin. DR3 was constitutively expressed in splenic and dermal $\gamma\delta$ T17 cells. TL1A induced cytokine production by splenic $\gamma\delta$ T17 cells synergistically with IL-23. TL1A exacerbated the dermal symptoms of the murine psoriasis model generated by intradermal IL-23 injection in wild-type but not in $\gamma\delta$ T cell-deficient mice. These findings suggest a novel regulatory mechanism of $\gamma\delta$ T17 cells through TL1A and DR3 and its involvement in psoriasis pathogenesis as a possible therapeutic target. In conclusion, the present study on the novel regulatory mechanism of $\gamma\delta$ T17 cells may provide insights into psoriasis pathogenesis and therapeutics.