



HOKKAIDO UNIVERSITY

Title	A Braconid Parasite of <i>Theretra japonica</i> De L'orzo (Host Record of Braconidae, 3)
Author(s)	Watanabe, Chihisa
Citation	<i>Insecta matsumurana</i> , 14(2-3), 94-94
Issue Date	1940-03
Doc URL	https://hdl.handle.net/2115/9442
Type	departmental bulletin paper
File Information	14(2-3)_p94-94.pdf



KUROSAWA as a solitary parasite of *Margaronia pyloalis* at Kyoto.

Cocoon: Dark brown to light brown, spindle-shaped, covered with a small amount of loose silk. Length, 7-10 mm., width, 3-4 mm.

Habitat: Honshu (Kyoto, 7 ♀♀, 9 ♂♂, VI, 1939, M. KUROSAWA leg.).

Gen. Distr.: Japan; Formosa; Manchoukuo; China; The Philippines.

摘 要

本文に記述せるクハノメイガの寄生蜂の學名及び和名は次の如し。

<i>Goniozus japonicus</i> ASHMEAD	ハマキアリガタバチ (ハマキタマゴバチ*)
<i>Epiurus persimilis</i> ASHMEAD	クロヒゲヒラタヒメバチ
<i>Spilocryptus japonicus</i> UCHIDA	シロフトガリヒメバチ
<i>Chelonus tabonus</i> SONAN	ワタノメイガカウラコマユ
<i>Phanerotoma planifrons</i> (NEES)	ヒメキイロカウラコマユ
<i>Apanteles minor</i> (ASHMEAD)	ヒメクロツヤサムラヒコマユ (ヒメクロツヤヤドリ**)
<i>Apanteles kurosawai</i> WATANABE (sp. nov.)	クハノメイガサムラヒコマユ
<i>Macrocentrus philippinensis</i> ASHMEAD	キマダラヒゲナガコマユ

*松村松年: 日本益蟲目錄, p. 96 (1908).

**———: 同上, p. 212 (1908).

A BRACONID PARASITE OF *THERETRA* *JAPONICA* DE L'ORZO (HOST RECORD OF BRACONIDAE, III)

By

CHIHIISA WATANABE

Microplitis theretrae WATANABE

Microplitis theretrae WATANABE, Jour. Facul. Agr., Hokkaido Imp. Univ., XLII, p. 108, ♀, Pl. III, fig. 3 (1937).

Host: *Theretra japonica* DE L'ORZO

This species was originally described as a parasite of *Theretra nessus* DRURY. The writer has received a female bred from the larva of *Theretra japonica* DE L'ORZO by Mr. T. ISHIZAWA at Ogikubo, Tokyo.

Habitat: Honshu (Ogikubo, Tokyo, 1 ♀, 12. V, 1939, T. ISHIZAWA leg.).

General Distribution: Japan (Honshu).