



Title	A new Species of Hesperidae from Formosa
Author(s)	Matsumura, Shonen
Citation	Insecta matsumurana, 15(1-2), 1-2
Issue Date	1940-12
Doc URL	<a href="https://hdl.handle.net/2115/9453">https://hdl.handle.net/2115/9453</a>
Type	departmental bulletin paper
File Information	15(1-2)_p1-2.pdf



A NEW SPECIES OF *HESPERIDAE*  
FROM FORMOSA

By

SHONEN MATSUMURA

(松村松年)

(With two Text-figures)

*Bibacis argenteola* n. sp.

♂. Wings chest-nut brown, with a yellowish shade towards the base.

Upperside: Primaries in the middle with 4 golden yellowish patches in a longitudinal row, 2 of which laying on both ends being much smaller; another smaller patch at the base of the interspace 5, near the apex in the interspaces 5, 8, 9 and 10 with each a white speck in an oblique row; veins somewhat darker in colour; fringes grayish, at the base of each vein darker. Secondaries at the outer margin checked with yellowish gray fringes; the anal lobe produced moderately.




Fig. 1

*Bibacis argenteola* n. sp. (♂) ×15  
Upperside

Underside: Purplish brown, the 5th part of the termen grayish. Primaries with the markings as on the upperside. Secondaries in the middle with a broad silvery band which becomes much narrower at the vein 8; at the interspace 5 produced outwardly and at the interspace 2 roundly ended; outwardly with an obscure grayish wavy band which is not distinct in the interspaces 4 and 5; the 5th part of the terminal portion as in the primaries.

Exp.: 52 mm.




Fig. 2

*Bibacis argenteola* n. sp. (♂) × 1.5

Underside

Hab.: Formosa; one male specimen was collected at Nôkô in the Prov. Taichu near Hori in June, 1940, by my collector. This species comes near *B. sena* MOOR. from the Philippines and Indian Region.

This extraordinary species may belong to a new genus, but the author puts it in the above genus.