



HOKKAIDO UNIVERSITY

Title	Comprehensive overview of disease models for Wolfram syndrome : toward effective treatments
Author(s)	Morikawa, Shuntaro; Tanabe, Katsuya; Kaneko, Naoya et al.
Citation	Mammalian genome, 35, 1-12 https://doi.org/10.1007/s00335-023-10028-x
Issue Date	2024-03-01
Doc URL	https://hdl.handle.net/2115/94707
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature' s AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: https://doi.org/10.1007/s00335-023-10028-x
Type	journal article
File Information	HUSCAP_★Wolfram_Models_Review_ver13.pdf



Comprehensive Overview of Disease Models for Wolfram Syndrome: Towards Effective Treatments

Shuntaro Morikawa^{1*}, Katsuya Tanabe², Naoya Kaneko¹, Nozomi Hishimura¹, Akie Nakamura¹

¹ Department of Pediatrics, Hokkaido University Hospital, Sapporo, Japan

² Division of Endocrinology, Metabolism, Haematological Science and Therapeutics, Yamaguchi University Graduate School of Medicine, Ube, Japan.

*Correspondence:

Shuntaro Morikawa, MD, PhD

Department of Pediatrics, Hokkaido University Hospital

North 14, West 5, Kita-ku, Sapporo, Japan, Zip 060-8638

Email: shuntaro.morikawa@med.hokudai.ac.jp

Acknowledgments

S.M. thanks Dr. Fumihiko Urano (Washington University School of Medicine in St. Louis), Dr. Toshihiro Tajima (Department of Pediatrics, Jichi Medical University), and Dr. Atsushi Manabe (Department of Pediatrics, Hokkaido University Graduate School of Medicine) for their mentorship and Sachiko Nakayama, Yui Sakurai, Ryoko Miyoshi, Kaoru Yokoki, and Miki Kaiho (Department of Pediatrics, Hokkaido University) for their invaluable secretarial assistance.

Abstract

Wolfram syndrome (OMIM 222300) is a rare autosomal recessive disease with a devastating array of symptoms, including diabetes mellitus, optic nerve atrophy, diabetes insipidus, hearing loss, and neurological dysfunction. The discovery of the causative gene, *WFS1*, has propelled research on this disease. However, a comprehensive understanding of the function of *WFS1* remains unknown, making the development of effective treatment a pressing challenge. To bridge these knowledge gaps, disease models for Wolfram syndrome are indispensable, and understanding the characteristics of each model is critical. This review will provide a summary of the current knowledge regarding *WFS1* function and offer a comprehensive overview of established disease models for Wolfram syndrome, covering animal models such as mice, rats, flies, and zebrafish, along with induced pluripotent stem cell (iPSC)-derived human cellular models. These models replicate key aspects of Wolfram syndrome, contributing to a deeper understanding of its pathogenesis and providing a platform for discovering potential therapeutic approaches.

Key words

Wolfram syndrome, *WFS1*, disease model, ER stress, iPSC

1. Introduction

Wolfram Syndrome (OMIM 222300) is a rare autosomal recessive disease characterized by several symptoms, including diabetes mellitus, optic nerve atrophy, diabetes insipidus, hearing loss, and neurological dysfunction. Two types of Wolfram syndrome have been identified: type 1 and type 2, caused by pathogenic variants of *WFS1* and *CISD2*, respectively. Most Wolfram syndrome cases are classified as type 1, so we will refer to them as “Wolfram syndrome” throughout this review. Since the discovery of *WFS1* as the causative gene of Wolfram syndrome, several animal and cellular models have been established. These disease models are indispensable tools in the research on Wolfram syndrome. However, a comprehensive summary of the characteristics and phenotypes of these experimental models is lacking. Therefore, this review aims to fill this gap by providing an overview of the animal and cellular models of Wolfram syndrome established to date. In this review, we first discuss (1) the clinical symptoms of Wolfram syndrome, (2) the function of *WFS1* and the pathophysiology caused by its abnormalities, and (3) the characteristics of disease models of Wolfram syndrome.

2. Clinical features of Wolfram syndrome and the discovery of *WFS1*

This section describes the clinical features of Wolfram syndrome and the disease concept associated with the causative gene *WFS1*.

Major symptoms of Wolfram syndrome

The first cases of Wolfram syndrome were reported by Wolfram and Wagener in 1938 (Wolfram and Wagener 1938). The report described four siblings who developed juvenile-onset diabetes mellitus and optic nerve atrophy. Subsequently, these patients also developed hearing loss and neurogenic bladder (Paley and Tunbridge 1956). In 1956, Paley and Tunbridge reported two additional cases of diabetes and optic nerve atrophy, suggesting a hereditary component (Paley and Tunbridge 1956). In 1977, a comprehensive review of 91 cases by Cremers et al. (Cremers et al. 1977) led to the official designation of this condition as “Wolfram syndrome.” Since then, numerous reports on the clinical symptoms of patients have emerged. The first nationwide survey was conducted and reported by Barrett et al. in 1995 (Barrett et al. 1995), revealing a prevalence of Wolfram syndrome at 1 in 770,000. Typically, patients with Wolfram syndrome develop diabetes mellitus at a median age of 6 years and optic atrophy around 11 years. Diabetes insipidus and sensorineural hearing loss typically manifest in the second decade of life, while renal tract abnormalities and neurological complications such as cerebellar ataxia appear by the fourth decade. Neurological and urinary tract manifestations are the most common causes of morbidity and mortality (Kinsley et al. 1995). Wolfram syndrome is characterized by a reduced size of intracranial volume, particularly in the brainstem and cerebellum, leading to balance impairment (Hershey et al. 2012; Pickett et al. 2012). However, the severity of these symptoms can vary among individuals with Wolfram syndrome (De Franco et al. 2017).

Discovery of *WFS1*

Wolfram syndrome was previously believed to be a mitochondrial disease due to the similarities in symptoms and the presence of mitochondrial DNA deletions in some patients (Bu and Rotter 1993; Bunday et al. 1993; Rötig et al. 1993; Vora and Lilleyman 1993). However, subsequent cases demonstrated the absence of mitochondrial DNA abnormalities, and the inheritance pattern suggested an abnormality in the nuclear genome of patients with Wolfram syndrome. Linkage analysis suggested the presence of a causative gene for Wolfram syndrome on chromosome 4p16 (Polymeropoulos et al. 1994). In 1998, Inoue et al., followed soon after by Strom et al., identified the *WFS1* gene (Inoue et al. 1998; Strom et al. 1998) (Fig. 1). Since the discovery of *WFS1*, the understanding of the pathogenesis of Wolfram syndrome has remarkably advanced. Approximately 200 *WFS1* variants associated with Wolfram syndrome have been reported, mostly found in exon 8 of *WFS1* (Smith et al. 2004). *WFS1* encodes the protein WFS1, also known as wolframin, which is an endoplasmic reticulum (ER)-membrane protein with multiple transmembrane domains comprising 890 amino acids (Strom et al. 1998; Takeda et al. 2001). WFS1 is expressed ubiquitously in various tissues, with the highest levels observed in the pancreas, heart, and brain, particularly in the hippocampus, amygdaloid area, and olfactory tubercle (Hofmann et al. 2003; Takeda et al. 2001). Recent studies have reported predominant localization of WFS1 in the ER membrane, particularly in the mitochondria-associated ER membranes, which serve as contact sites between the ER and mitochondria (Angebault et al. 2018; La Morgia et al. 2020).

Wolfram syndrome and WFS1-related disorders

Wolfram syndrome, a rare disorder caused by recessive *WFS1* variants, is now recognized to exhibit a phenotypic spectrum. Recent findings suggest that its frequency may be higher in certain races and populations than previously assumed (Bansal et al. 2018; De Franco et al. 2017; Li et al. 2020; Marchand et al. 2021). Moreover, there are related disorders caused by a dominant inheritance form of *WFS1*, known as “*WFS1*-related disorders” or “Wolfram-like syndrome.” These disorders manifest with one or more symptoms observed in Wolfram syndrome. The severity and symptoms of *WFS1*-related disorders can vary, ranging from mild diabetes mellitus, hearing loss, congenital cataracts, and optic atrophy developing independently to severe neonatal cases that exhibit all these symptoms (Kobayashi et al. 2018; Mets et al. 2010; Morikawa et al. 2017). It has also been reported that specific heterozygous *WFS1* variants can lead to non-syndromic low-frequency sensorineural hearing loss (Bespalova et al. 2001; Cryns et al. 2002; Young et al. 2001).

3. *WFS1* functions and their anomalies

Some *WFS1* variants have also been implicated in other types of diabetes. Several large-scale genome-wide association studies have reported that several single nucleotide polymorphisms (SNPs) in *WFS1* are associated with an increased genetic risk for type 2 diabetes mellitus (T2DM), irrespective of racial differences (Cheurfa et al. 2011; Elek et al. 2015; Heni et al. 2010; Lee et al. 2008; Long et al. 2012; Lyssenko et al. 2008; Sandhu et al. 2007; Sparsø et al. 2008; van Hoek et al. 2008; Westermarck et al. 2010). However, a meta-analysis has shown that two specific *WFS1* SNPs, rs734312 and rs10010131, have a significant protective effect against the risk of developing T2DM (Cheng et al. 2013). Accordingly, understanding the pathogenesis of Wolfram syndrome and *WFS1*-related disorders may provide valuable insights into the pathogenesis of other types of diabetes, including T2DM. In this section, we describe the known physiological functions of *WFS1* and the pathological conditions that arise due to its abnormalities.

Interaction between WFS1 and Ca²⁺ channels or pumps

The ER serves as an intracellular Ca²⁺ store, and WFS1 is expressed on the ER membrane. Therefore, numerous studies have explored the relationship between WFS1 and intracellular Ca²⁺ homeostasis (La Morgia et al. 2020; Nguyen et al. 2020; Osman et al. 2003). The first study to demonstrate a connection between WFS1 and intracellular Ca²⁺ was reported by Osman et al. (Osman et al. 2003), who showed elevated levels of intracellular Ca²⁺ in *Xenopus* oocytes overexpressing *WFS1*.

Takei et al. first revealed the relationship between Ca²⁺ concentration in the ER and the expression level of WFS1 (Takei et al. 2006). They found that WFS1 regulates ER Ca²⁺ storage and cytosolic Ca²⁺ homeostasis by increasing the Ca²⁺ uptake and store-operated Ca²⁺ entry (SOCE). This regulation involves the modulation of Ca²⁺ pumps and channels, such as sarcoendoplasmic reticulum ATPase (SERCA) and inositol 1,4,5-trisphosphate receptor (IP₃R), localized on the

ER membrane (Kunnappallil and Hasan 2022). WFS1 deficiency results in upregulated SERCA expression, leading to increased Ca^{2+} pumping into the ER (Zatyka et al. 2015) (Fig. 2A). Additionally, WFS1 forms a complex with neuronal calcium sensor 1 (NCS1) and IP_3R to facilitate Ca^{2+} transfer between the ER and mitochondria. WFS1 deficiency reduces the expression level of NCS1, followed by impaired ER-mitochondrial contact and subsequent Ca^{2+} uptake into the mitochondria (Angebault et al. 2018) (Fig. 2B). Therefore, therapeutic approaches targeting intracellular Ca^{2+} signaling have been explored as potential treatments for Wolfram syndrome (Abreu et al. 2021; Akiyama et al. 2009; Clark et al. 2017; Crouzier et al. 2022a; Lu et al. 2014; Nguyen et al. 2020). Apart from its interactions with Ca^{2+} pumps and channels, WFS1 also interacts with the Na^+/K^+ ATPase beta 1 subunit, the V1A subunit of the H1 ATPase, and the voltage-dependent anion channel isoform 1 (VDAC1) (Gharanei et al. 2013; Zatyka et al. 2008; Zatyka et al. 2023). In *WFS1*-depleted cells, the H1 ATPase V1A subunit is degraded more rapidly (Gharanei et al. 2013).

WFS1 as a component of unfolded protein response

The link between WFS1 and ER stress was first described by Ueda et al. (Ueda et al. 2005). They demonstrated that the *WFS1* expression level increased in response to ER stress. The gene expression levels of *Wfs1* are considered ER stress markers (Lipson et al. 2006) due to the presence of a conserved sequence in the promoter region similar to the ER stress response element (Kakiuchi et al. 2006). ER stress-induced *WFS1* upregulation requires the activation of inositol requiring 1 and PKR-like endoplasmic reticulum kinase, which are key regulators of the unfolded protein response (UPR) (Fonseca et al. 2005). Another UPR regulator, activating transcription factor (ATF) 6 β , binds to the *Wfs1* gene promoter and induces both gene and protein expression (Odisho et al. 2015).

WFS1 has additional roles in the regulation of ER stress and protein degradation. It stabilizes the E3 ubiquitin ligase HRD1 and regulates the degradation of ATF6 α by facilitating ATF6 transport to the proteasome (Fonseca et al. 2010) (Fig. 2C). This means WFS1 acts as a UPR regulator; its deficiency in pancreatic β -cells induces pathogenic ER stress, leading to impaired cell cycle and accelerated cell apoptosis (Yamada et al. 2006). However, considering their close interaction, both ER stress and intracellular Ca^{2+} homeostasis may be involved in the pathogenesis of Wolfram syndrome. In fact, *WFS1* deficiency induces ER stress, resulting in IP_3R dysfunction and disturbed cytosolic Ca^{2+} homeostasis, which subsequently affects mitochondrial dynamics (Blackstone et al. 2016). This leads to inhibited mitochondrial fusion and trafficking, as well as augmented mitophagy, contributing to delayed neuronal development. ER stress is recognized for disrupting intracellular Ca^{2+} homeostasis and inducing cellular inflammation. The concept of “sterilized inflammation,” which refers to non-infectious inflammation triggered by pathogenic ER-stress (Lerner et al. 2012; Osowski et al. 2012), has recently emerged. In line with these observations, ER-stress-induced sterilized inflammation caused by WFS1 dysfunction accelerates disease progression in Wolfram syndrome (Morikawa et al. 2022; Panfili et al. 2021).

WFS1 function in intracellular trafficking

WFS1 localization extends beyond the ER to include secretory granules (Hatanaka et al. 2011). In *WFS1*-deficient

pancreatic β -cells, insulin secretory granule acidification becomes impaired, leading to impaired insulin exocytosis (Hatanaka et al. 2011) (Fig. 2D). The acidification of insulin granule is necessary for efficient proinsulin processing because endopeptidase PC1/3 and PC2 activate with a low pH optimum. While the precise interaction between WFS1 expressed on the insulin granule membrane and impaired insulin granule acidification has not been elucidated, it is speculated that WFS1 regulate the activities of V-type H^+ -ATPase and CLC-3 Cl^- channels expressed on the insulin granule membrane (Hatanaka et al. 2011). More recently, it was demonstrated that WFS1 serves as a vesicular cargo protein involved in intracellular trafficking (Wang et al. 2021). Specifically, the C-terminus of WFS1 directly binds to transported proteins within the ER lumen, while the N-terminus of WFS1 interacts with SEC24, a subunit of the coat protein complex II (COPII) located in the cytoplasm. This interaction allows for vesicular transport from the ER to the Golgi (Fig. 2E). This function of WFS1 has been shown to be necessary for the transportation of proteins such as proinsulin from the ER to the Golgi in pancreatic β -cells (Wang et al. 2021).

4. Experimental models of Wolfram syndrome

Experimental models are indispensable tools for conducting research to understand the pathogenesis of Wolfram syndrome and to develop therapeutic strategies. Over the years, several disease models have been established to study this condition. In the 2000s, *Wfs1*-knockout mice were generated as an early animal model. Subsequently, in the 2010s, disease models were extended to include rats, flies, and zebrafish. Recently, a *Wfs1* pathogenic variant knock-in mouse model has also been reported. In this section, we summarize the phenotypic characteristics observed in these Wolfram syndrome animal models and also include information on induced pluripotent stem cell (iPSC)-derived cellular models (Table).

Animal models of Wolfram syndrome

The first animal model, Wolfram syndrome, was established by Ishihara et al. in 2004 (Ishihara et al. 2004). The whole-body *Wfs1*-deficient B6 background mice exhibited decreased pancreatic insulin content at 2 weeks old, impaired glucose homeostasis, and decreased insulin secretion around 17 weeks old. The diabetic phenotype in these mice resulted from ER stress-induced pancreatic β -cell death and was accelerated by crossing with agouti lethal yellow mice, a model that develops obesity, insulin resistance, and pancreatic β -cell hyperplasia (Akiyama et al. 2009). Similar to Wolfram syndrome patients, these mice also exhibited symptoms of central diabetes insipidus due to arginine vasopressin (AVP) deficiency (Kurimoto et al. 2021), impaired retinal functions (Bonnet Wersinger et al. 2014), and psychiatric symptoms observed in behavioral studies (Kato et al. 2008). In addition to the B6 strain, another whole-body *Wfs1* knockout mouse model on the 129S6 background was developed (Köks et al. 2009; Luuk et al. 2008). These 129S6 mice, in which LacZ replaces exon 8 of *Wfs1*, showed an earlier onset of diabetes compared to the B6 strain (Abreu et al. 2020). Interestingly, males were found to be at higher risk of developing diabetes than females (Noormets et al. 2011). The phenotype of this 129S6-background mouse model is well studied, and besides the diabetic phenotype, this mouse model recapitulates many

of the symptoms observed in patients with Wolfram syndrome (Blackstone et al. 2016; Ivask et al. 2021; Luuk et al. 2009; Noormets et al. 2009; Noormets et al. 2014; Richard et al. 2023; Visnapuu et al. 2013a; Visnapuu et al. 2013b; Waszczykowska et al. 2020). The only *Wfs1*-conditional knockout mouse model was reported by Riggs et al. These B6 background mice, with specifically knocked out *Wfs1* in pancreatic β -cells, develop glucose intolerance and insulin deficiency by 12 weeks of age (Riggs et al. 2005). Most recently, a *Wfs1* E864K knock-in mouse model was reported (Richard et al. 2023). These *Wfs1* E864K knock-in mice develop an early onset of severe vestibular dysfunction besides hearing loss. Human patients with Wolfram syndrome carrying the *WFS1* p.E864K variant develop low-frequency sensorineural hearing loss, optic atrophy, and impaired glucose tolerance. However, no vestibular neuropathy has been reported. These facts suggest that the role of *WFS1* in the human vestibular system may not be as crucial as in rodents.

Besides the mouse model, the *Wfs1*-deficient rat is a well-established animal model for Wolfram syndrome (Plaas et al. 2017). This model involves whole-body *Wfs1* knockout in Sprague-Dawley background rats, which show impaired glucose homeostasis starting at 3 months of age. At 15 months of age, these rats exhibit retinal gliosis, optic nerve atrophy, and reduced medullary volume. Although the onset age of each symptom is older than that in the mouse models, the larger body size of rats makes it advantageous for conducting therapeutic interventions and sample collection. Recently, the drosophila and zebrafish models with *WFS1* homolog knockout have been reported as additional models for Wolfram syndrome. Optical transparency of the zebrafish embryo enables the observation of neural development *in vivo*. Drosophila has a short generation time and a small body size, making it possible to test many potential therapeutic molecules. Using these Wolfram syndrome zebrafish and Drosophila models, phenotypes caused by *WFS1* deficiency, including neurodegeneration, visual dysfunction, and mitochondrial dysfunction, have been well-studied (Cairns et al. 2021; Crouzier et al. 2022b; Kunnappallil and Hasan 2022; Sakakibara et al. 2018).

The pursuit of novel therapeutic approaches to combat Wolfram syndrome is actively underway. One promising strategy involves repurposing existing diabetic medications for the treatment of Wolfram syndrome, such as GLP-1 receptor agonists (Panfili et al. 2023). Notably, exenatide and dulaglutide have been reported to improve glucose tolerance in Wolfram syndrome mouse models (Gorgogietas et al. 2023; Kondo et al. 2018; Sedman et al. 2016). Similarly, another GLP-1 receptor agonist, liraglutide, was found to prevent the development of glucose intolerance, reduce neuroinflammation, protect against retinal ganglion cell death, and delay the progression of hearing and vision loss in *Wfs1* mutant rat models (Jagom e et al. 2021; Seppa et al. 2019; Seppa et al. 2021; Toots et al. 2018).

iPSC-derived models of Wolfram syndrome

The first successfully developed insulin-producing cells from patients with Wolfram syndrome were reported by Shang et al. in 2014 (Shang et al. 2014). In their study, the authors generated insulin-producing cells from skin fibroblast-derived iPSCs of Wolfram syndrome patients. These patient-derived insulin-producing cells showed lower insulin content and reduced tolerance to ER stress compared to controls. This cellular model provided crucial insights into ER stress-related diseases derived from human patients. Maxwell et al. improved the differential protocol for iPSC-derived β -cells (SC- β -cells) and used CRISPR/Cas9 techniques to explore the potential of gene therapy and personalized cell therapy for

Wolfram syndrome (Maxwell et al. 2020). In their study, they successfully restored insulin secretion in patient-derived SC- β -cells through the correction of *WFS1*. Transplanting the gene-edited SC- β -cells into diabetic mice resulted in improved blood glucose profile. In addition, they found that SC- β -cells differentiated from Wolfram syndrome patients with mild *WFS1* variants retained their insulin secretory capacity (Kitamura et al. 2022). This indicates that these SC- β -cell models exhibit phenotypes consistent with the symptoms observed in patients. Using these SC- β -cell models, a combination treatment of 4-phenylbutyric acid and tauroursodeoxycholic acid has been demonstrated to effectively address the diabetic phenotype in Wolfram syndrome patients with the *WFS1* p.R558C variant (Kitamura et al. 2022). Besides pancreatic β -cells, neural cells have also been developed as a disease model from Wolfram syndrome patient iPSCs. Impaired neurite outgrowth was observed in neurons derived from Wolfram syndrome patients, and this morphological change was prevented by treatment with valproic acid (Pourtoy-Brasselet et al. 2021). Valproic acid has been reported to reduce ER stress and act as a histone deacetylase inhibitor for neurons. It is not clear how valproic acid promotes neurite outgrowth in the neuronal cell model of Wolfram syndrome. However, valproic acid may exert neuroprotective effects by reducing ER stress and through pleiotropic mechanisms. Zatyka et al. demonstrated that the interaction between *WFS1* and *VDAC1* is essential for mitochondrial function and dynamics in iPSC-derived neural cells (Zatyka et al. 2023). These models could prove useful for studying the pathogenesis of brainstem atrophy and neurodegeneration observed in Wolfram syndrome.

5. Future expectations

Numerous clinical and basic studies have significantly contributed to understanding the function of *WFS1* and advancing towards a cure for Wolfram syndrome. In the future, further research is expected to uncover the unknown pathogenesis, particularly regarding the development of optic nerve atrophy and neurological symptoms associated with *WFS1* dysfunction. Furthermore, the development of therapeutic agents against Wolfram syndrome necessitates the use of patient-derived disease models. In addition to pancreatic β -cells and neural cells, the development of retinal ganglion, inner ear, and AVP-producing neural cells is important. These patient-derived cellular models can serve as invaluable tools for developing pre-clinical and personalized therapeutic reagents against visual impairment, hearing loss, and central diabetes insipidus in Wolfram syndrome. Research on Wolfram syndrome, including the development of various disease models, continues to serve as a prototype for studying ER-stress-related diseases.

Statements and Declarations

Competing interests

The authors have no conflict of interest to declare.

Author contributions

S.M. performed the literature search and wrote the draft. S.M., K.T., N.K., N.H., and A.K. critically reviewed the manuscript. All authors approved the final version of the manuscript to be published.

Data availability statement

Data sharing is not applicable to this review article.

Funding

S.M. received funding from The Japanese Society for Pediatric Endocrinology Future Development Grant, which is supported by Novo Nordisk Pharma Ltd. K.T. received funding from Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology, Japan (23K08011).

References

- Abreu D, Asada R, Revilla JMP, Lavagnino Z, Kries K, Piston DW, Urano F (2020) Wolfram syndrome 1 gene regulates pathways maintaining beta-cell health and survival. *Lab Invest* 100:849–862. <https://doi.org/10.1038/s41374-020-0408-5>
- Abreu D, Stone SI, Pearson TS, Bucelli RC, Simpson AN, Hurst S, Brown CM, Kries K, Onwumere C, Gu H, Hoekel J, Tychsen L, Van Stavern GP, White NH, Marshall BA, Hershey T, Urano F (2021) A phase Ib/Ila clinical trial of dantrolene sodium in patients with Wolfram syndrome. *JCI Insight* 6. <https://doi.org/10.1172/jci.insight.145188>
- Akiyama M, Hatanaka M, Ohta Y, Ueda K, Yanai A, Uehara Y, Tanabe K, Tsuru M, Miyazaki M, Saeki S, Saito T, Shinoda K, Oka Y, Tanizawa Y (2009) Increased insulin demand promotes while pioglitazone prevents pancreatic beta cell apoptosis in Wfs1 knockout mice. *Diabetologia* 52:653–663. <https://doi.org/10.1007/s00125-009-1270-6>
- Angebault C, Fauconnier J, Patergnani S, Rieusset J, Danese A, Affortit CA, Jagodzinska J, Mégy C, Quiles M, Cazevielle C, Korchagina J, Bonnet-Wersinger D, Milea D, Hamel C, Pinton P, Thiry M, Lacampagne A, Delprat B, Delettre C (2018) ER-mitochondria cross-talk is regulated by the Ca(2+) sensor NCS1 and is impaired in Wolfram syndrome. *Sci Signal* 11. <https://doi.org/10.1126/scisignal.aag1380>
- Bansal V, Boehm BO, Darvasi A (2018) Identification of a missense variant in the WFS1 gene that causes a mild form of Wolfram syndrome and is associated with risk for type 2 diabetes in Ashkenazi Jewish individuals. *Diabetologia* 61:2180–2188. <https://doi.org/10.1007/s00125-018-4690-3>
- Barrett TG, Bunday SE, Macleod AF (1995) Neurodegeneration and Diabetes: UK nationwide study of Wolfram (DIDMOAD) syndrome. *Lancet* 346:1458–1463. [https://doi.org/10.1016/s0140-6736\(95\)92473-6](https://doi.org/10.1016/s0140-6736(95)92473-6)
- Bespalova IN, Van Camp G, Bom SJ, Brown DJ, Cryns K, DeWan AT, Erson AE, Flothmann K, Kunst HP, Kurnool P, Sivakumaran TA, Cremers CW, Leal SM, Burmeister M, Lesperance MM (2001) Mutations in the Wolfram syndrome 1 gene (WFS1) are a common cause of low frequency sensorineural hearing loss. *Hum Mol Genet* 10:2501–2508. <https://doi.org/10.1093/hmg/10.22.2501>

314 Blackstone C, Cagalinec M, Liiv M, Hodurova Z, Hickey MA, Vaarmann A, Mandel M, Zeb A, Choubey V, Kuum M, Safiulina D,
 315 Vasar E, Veksler V, Kaasik A (2016) Role of mitochondrial dynamics in neuronal development: mechanism for Wolfram
 316 syndrome. *PLOS Biol* 14

317 Bonnet Wersinger D, Benkafadar N, Jagodzinska J, Hamel C, Tanizawa Y, Lenaers G, Delettre C (2014) Impairment of visual
 318 function and retinal ER stress activation in Wfs1-deficient mice. *PLOS ONE* 9:e97222.
 319 <https://doi.org/10.1371/journal.pone.0097222>

320 Bu X, Rotter JI (1993) Wolfram syndrome: a mitochondrial-mediated disorder? *Lancet* 342:598–600.
 321 [https://doi.org/10.1016/0140-6736\(93\)91416-j](https://doi.org/10.1016/0140-6736(93)91416-j)

322 Bunday S, Fielder A, Poulton K (1993) Wolfram syndrome: mitochondrial disorder. *Lancet* 342:1059–1060.
 323 [https://doi.org/10.1016/0140-6736\(93\)92918-j](https://doi.org/10.1016/0140-6736(93)92918-j)

324 Cairns G, Burté F, Price R, O'Connor E, Toms M, Mishra R, Moosajee M, Pyle A, Sayer JA, Yu-Wai-Man P (2021) A mutant wfs1
 325 zebrafish model of Wolfram syndrome manifesting visual dysfunction and developmental delay. *Sci Rep* 11:20491.
 326 <https://doi.org/10.1038/s41598-021-99781-0>

327 Cheng S, Wu Y, Wu W, Zhang D (2013) Association of rs734312 and rs10010131 polymorphisms in WFS1 gene with type 2
 328 diabetes mellitus: a meta-analysis. *Endocr J* 60:441–447. <https://doi.org/10.1507/endocrj.EJ12-0325>

329 Cheurfa N, Brenner GM, Reis AF, Dubois-Laforgue D, Roussel R, Tichet J, Lantieri O, Balkau B, Fumeron F, Timsit J, Marre M,
 330 Velho G (2011) Decreased insulin secretion and increased risk of type 2 diabetes associated with allelic variations of the WFS1
 331 gene: the Data from Epidemiological Study on the insulin Resistance Syndrome (DESIR) prospective study. *Diabetologia*
 332 54:554–562. <https://doi.org/10.1007/s00125-010-1989-0>

333 Clark AL, Kanekura K, Lavagnino Z, Spears LD, Abreu D, Mahadevan J, Yagi T, Semenkovich CF, Piston DW, Urano F (2017)
 334 Targeting cellular calcium homeostasis to prevent cytokine-mediated beta cell death. *Sci Rep* 7:5611.
 335 <https://doi.org/10.1038/s41598-017-05935-4>

336 Cremers CW, Wijdeveld PG, Pinckers AJ (1977) Juvenile diabetes mellitus, optic atrophy, hearing loss, diabetes insipidus, atonia
 337 of the urinary tract and bladder, and other abnormalities (Wolfram syndrome). A review of 88 cases from the literature with
 338 personal observations on 3 new patients. *Acta Paediatr Scand Suppl* (264):1–16. <https://doi.org/10.1111/j.1651-2227.1977.tb15069.x>

339

340 Crouzier L, Danese A, Yasui Y, Richard EM, Liévens JC, Patergnani S, Couly S, Diez C, Denus M, Cubedo N, Rossel M, Thiry M, Su
 341 TP, Pinton P, Maurice T, Delprat B (2022a) Activation of the sigma-1 receptor chaperone alleviates symptoms of Wolfram
 342 syndrome in preclinical models. *Sci Transl Med* 14:eabh3763. <https://doi.org/10.1126/scitranslmed.abh3763>

343 Crouzier L, Richard EM, Diez C, Alzaeem H, Denus M, Cubedo N, Delaunay T, Glendenning E, Baxendale S, Liévens JC, Whitfield
 344 TT, Maurice T, Delprat B (2022b) Morphological, behavioral and cellular analyses revealed different phenotypes in Wolfram
 345 syndrome wfs1a and wfs1b zebrafish mutant lines. *Hum Mol Genet* 31:2711–2727. <https://doi.org/10.1093/hmg/ddac065>

346 Cryns K, Pfister M, Pennings RJ, Bom SJ, Flothmann K, Caethoven G, Kremer H, Schatteman I, Köln KA, Tóth T, Kupka S, Blin N,
 347 Nürnberg P, Thiele H, van de Heyning PH, Reardon W, Stephens D, Cremers CW, Smith RJ, Van Camp G (2002) Mutations in the
 348 WFS1 gene that cause low-frequency sensorineural hearing loss are small non-inactivating mutations. *Hum Genet* 110:389–
 349 394. <https://doi.org/10.1007/s00439-002-0719-1>

350 De Franco E, Flanagan SE, Yagi T, Abreu D, Mahadevan J, Johnson MB, Jones G, Acosta F, Mulaudzi M, Lek N, Oh V, Petz O,
 351 Caswell R, Ellard S, Urano F, Hattersley AT (2017) Dominant ER stress-inducing WFS1 mutations underlie a genetic syndrome
 352 of neonatal/infancy-onset diabetes, congenital sensorineural deafness, and congenital cataracts. *Diabetes* 66:2044–2053.
 353 <https://doi.org/10.2337/db16-1296>
 354 Elek Z, Németh N, Nagy G, Németh H, Somogyi A, Hosszufalusi N, Sasvári-Székely M, Rónai Z (2015) Micro-RNA Binding Site
 355 polymorphisms in the WFS1 gene are risk factors of diabetes mellitus. *PLOS ONE* 10:e0139519.
 356 <https://doi.org/10.1371/journal.pone.0139519>
 357 Fonseca SG, Fukuma M, Lipson KL, Nguyen LX, Allen JR, Oka Y, Urano F (2005) WFS1 is a novel component of the unfolded
 358 protein response and maintains homeostasis of the endoplasmic reticulum in pancreatic beta-cells. *J Biol Chem* 280:39609–
 359 39615. <https://doi.org/10.1074/jbc.M507426200>
 360 Fonseca SG, Ishigaki S, Osowski CM, Lu S, Lipson KL, Ghosh R, Hayashi E, Ishihara H, Oka Y, Permutt MA, Urano F (2010)
 361 Wolfram syndrome 1 gene negatively regulates ER stress signaling in rodent and human cells. *J Clin Invest* 120:744–755.
 362 <https://doi.org/10.1172/JCI39678>
 363 Gharanei S, Zatyka M, Astuti D, Fenton J, Sik A, Nagy Z, Barrett TG (2013) Vacuolar-type H⁺-ATPase V1A subunit is a molecular
 364 partner of Wolfram syndrome 1 (WFS1) protein, which regulates its expression and stability. *Hum Mol Genet* 22:203–217.
 365 <https://doi.org/10.1093/hmg/ddt400>
 366 Gorgogietas V, Rajaei B, Heeyoung C, Santacreu BJ, Marín-Cañas S, Salpea P, Sawatani T, Musuaya A, Arroyo MN, Moreno-
 367 Castro C, Benabdallah K, Demarez C, Toivonen S, Cosentino C, Pachera N, Lytrivi M, Cai Y, Carnel L, Brown C, Urano F, Marchetti
 368 P, Gilon P, Eizirik DL, Cnop M, Igoillo-Esteve M (2023) GLP-1R agonists demonstrate potential to treat Wolfram syndrome in
 369 human preclinical models. *Diabetologia* 66:1306–1321. <https://doi.org/10.1007/s00125-023-05905-8>
 370 Hatanaka M, Tanabe K, Yanai A, Ohta Y, Kondo M, Akiyama M, Shinoda K, Oka Y, Tanizawa Y (2011) Wolfram syndrome 1 gene
 371 (WFS1) product localizes to secretory granules and determines granule acidification in pancreatic beta-cells. *Hum Mol Genet*
 372 20:1274–1284. <https://doi.org/10.1093/hmg/ddq568>
 373 Heni M, Ketterer C, Thamer C, Herzberg-Schäfer SA, Guthoff M, Stefan N, Machicao F, Staiger H, Fritsche A, Häring HU (2010)
 374 Glycemia determines the effect of type 2 diabetes risk genes on insulin secretion. *Diabetes* 59:3247–3252.
 375 <https://doi.org/10.2337/db10-0674>
 376 Hershey T, Lugar HM, Shimony JS, Rutlin J, Koller JM, Perantie DC, Paciorkowski AR, Eisenstein SA, Permutt MA, Washington
 377 University Wolfram Study Group (2012) Early brain vulnerability in Wolfram syndrome. *PLOS ONE* 7:e40604.
 378 <https://doi.org/10.1371/journal.pone.0040604>
 379 Hofmann S, Philbrook C, Gerbitz KD, Bauer MF (2003) Wolfram syndrome: structural and functional analyses of mutant and
 380 wild-type wolframin, the WFS1 gene product. *Hum Mol Genet* 12:2003–2012. <https://doi.org/10.1093/hmg/ddg214>
 381 Inoue H, Tanizawa Y, Wasson J, Behn P, Kalidas K, Bernal-Mizrachi E, Mueckler M, Marshall H, Donis-Keller H, Crock P, Rogers
 382 D, Mikuni M, Kumashiro H, Higashi K, Sobue G, Oka Y, Permutt MA (1998) A gene encoding a transmembrane protein is
 383 mutated in patients with diabetes mellitus and optic atrophy (Wolfram syndrome). *Nat Genet* 20:143–148.
 384 <https://doi.org/10.1038/2441>
 385 Ishihara H, Takeda S, Tamura A, Takahashi R, Yamaguchi S, Takei D, Yamada T, Inoue H, Soga H, Katagiri H, Tanizawa Y, Oka Y

(2004) Disruption of the WFS1 gene in mice causes progressive beta-cell loss and impaired stimulus-secretion coupling in insulin secretion. *Hum Mol Genet* 13:1159–1170. <https://doi.org/10.1093/hmg/ddh125>

Ivask M, Volke V, Raasmaja A, Kõks S (2021) High-fat diet associated sensitization to metabolic stress in Wfs1 heterozygous mice. *Mol Genet Metab* 134:203–211. <https://doi.org/10.1016/j.ymgme.2021.07.002>

Jagomäe T, Seppa K, Reimets R, Pastak M, Plaas M, Hickey MA, Kukker KG, Moons L, De Groef L, Vasar E, Kaasik A, Terasmaa A, Plaas M (2021) Early intervention and lifelong treatment with GLP1 receptor agonist liraglutide in a Wolfram syndrome rat model with an emphasis on visual neurodegeneration, sensorineural hearing loss and diabetic phenotype. *Cells* 10. <https://doi.org/10.3390/cells10113193>

Kakiuchi C, Ishiwata M, Hayashi A, Kato T (2006) XBP1 induces WFS1 through an endoplasmic reticulum stress response element-like motif in SH-SY5Y cells. *J Neurochem* 97:545–555. <https://doi.org/10.1111/j.1471-4159.2006.03772.x>

Kato T, Ishiwata M, Yamada K, Kasahara T, Kakiuchi C, Iwamoto K, Kawamura K, Ishihara H, Oka Y (2008) Behavioral and gene expression analyses of Wfs1 knockout mice as a possible animal model of mood disorder. *Neurosci Res* 61:143–158. <https://doi.org/10.1016/j.neures.2008.02.002>

Kinsley BT, Swift M, Dumont RH, Swift RG (1995) Morbidity and mortality in the Wolfram syndrome. *Diabetes Care* 18:1566–1570. <https://doi.org/10.2337/diacare.18.12.1566>

Kitamura RA, Maxwell KG, Ye W, Kries K, Brown CM, Augsornworawat P, Hirsch Y, Johansson MM, Weiden T, Ekstein J, Cohen J, Klee J, Leslie K, Simeonov A, Henderson MJ, Millman JR, Urano F (2022) Multidimensional analysis and therapeutic development using patient iPSC-derived disease models of Wolfram syndrome. *JCI Insight* 7. <https://doi.org/10.1172/jci.insight.156549>

Kobayashi M, Miyagawa M, Nishio SY, Moteki H, Fujikawa T, Ohyama K, Sakaguchi H, Miyanochara I, Sugaya A, Naito Y, Morita SY, Kanda Y, Takahashi M, Ishikawa K, Nagano Y, Tono T, Oshikawa C, Kihara C, Takahashi H, Noguchi Y, Usami SI (2018) WFS1 mutation screening in a large series of Japanese hearing loss patients: massively parallel DNA sequencing-based analysis. *PLOS ONE* 13:e0193359. <https://doi.org/10.1371/journal.pone.0193359>

Kõks S, Soomets U, Paya-Cano JL, Fernandes C, Luuk H, Plaas M, Terasmaa A, Tillmann V, Noormets K, Vasar E, Schalkwyk LC (2009) Wfs1 gene deletion causes growth retardation in mice and interferes with the growth hormone pathway. *Physiol Genomics* 37:249–259. <https://doi.org/10.1152/physiolgenomics.90407.2008>

Kondo M, Tanabe K, Amo-Shiinoki K, Hatanaka M, Morii T, Takahashi H, Seino S, Yamada Y, Tanizawa Y (2018) Activation of GLP-1 receptor signalling alleviates cellular stresses and improves beta cell function in a mouse model of Wolfram syndrome. *Diabetologia* 61:2189–2201. <https://doi.org/10.1007/s00125-018-4679-y>

Kunnappallil RS, Hasan G (2022). <https://doi.org/10.1101/2022.11.10.515972>

Kurimoto J, Takagi H, Miyata T, Hodai Y, Kawaguchi Y, Hagiwara D, Suga H, Kobayashi T, Sugiyama M, Onoue T, Ito Y, Iwama S, Banno R, Tanabe K, Tanizawa Y, Arima H (2021) Deficiency of WFS1 leads to the impairment of AVP secretion under dehydration in male mice. *Pituitary* 24:582–588. <https://doi.org/10.1007/s11102-021-01135-6>

La Morgia C, Maresca A, Amore G, Gramegna LL, Carbonelli M, Scimonelli E, Danese A, Patergnani S, Caporali L, Tagliavini F, Del Dotto V, Capristo M, Sadun F, Barboni P, Savini G, Evangelisti S, Bianchini C, Valentino ML, Liguori R, Tonon C, Giorgi C, Pinton P, Lodi R, Carelli V (2020) Calcium mishandling in absence of primary mitochondrial dysfunction drives cellular

pathology in Wolfram syndrome. *Sci Rep* 10:4785. <https://doi.org/10.1038/s41598-020-61735-3>

Lee YH, Kang ES, Kim SH, Han SJ, Kim CH, Kim HJ, Ahn CW, Cha BS, Nam M, Nam CM, Lee HC (2008) Association between polymorphisms in SLC30A8, HHEX, CDKN2A/B, IGF2BP2, FTO, WFS1, CDKAL1, KCNQ1 and type 2 diabetes in the Korean population. *J Hum Genet* 53:991–998. <https://doi.org/10.1007/s10038-008-0341-8>

Lerner AG, Upton JP, Praveen PV, Ghosh R, Nakagawa Y, Igbaria A, Shen S, Nguyen V, Backes BJ, Heiman M, Heintz N, Greengard P, Hui S, Tang Q, Trusina A, Oakes SA, Papa FR (2012) IRE1alpha induces thioredoxin-interacting protein to activate the NLRP3 inflammasome and promote programmed cell death under irremediable ER stress. *Cell Metab* 16:250–264. <https://doi.org/10.1016/j.cmet.2012.07.007>

Li M, Wang S, Xu K, Chen Y, Fu Q, Gu Y, Shi Y, Zhang M, Sun M, Chen H, Han X, Li Y, Tang Z, Cai L, Li Z, Shi Y, Yang T, Polychronakos C (2020) High prevalence of a monogenic cause in Han Chinese diagnosed with Type 1 diabetes, partly driven by nonsyndromic recessive WFS1 mutations. *Diabetes* 69:121–126. <https://doi.org/10.2337/db19-0510>

Lipson KL, Fonseca SG, Ishigaki S, Nguyen LX, Foss E, Bortell R, Rossini AA, Urano F (2006) Regulation of insulin biosynthesis in pancreatic beta cells by an endoplasmic reticulum-resident protein kinase IRE1. *Cell Metab* 4:245–254. <https://doi.org/10.1016/j.cmet.2006.07.007>

Long J, Edwards T, Signorello LB, Cai Q, Zheng W, Shu XO, Blot WJ (2012) Evaluation of genome-wide association study-identified type 2 diabetes loci in African Americans. *Am J Epidemiol* 176:995–1001. <https://doi.org/10.1093/aje/kws176>

Lu S, Kanekura K, Hara T, Mahadevan J, Spears LD, Oslowski CM, Martinez R, Yamazaki-Inoue M, Toyoda M, Neilson A, Blanner P, Brown CM, Semenkovich CF, Marshall BA, Hershey T, Umezawa A, Greer PA, Urano F (2014) A calcium-dependent protease as a potential therapeutic target for Wolfram syndrome. *Proc Natl Acad Sci U S A* 111:E5292–E5301. <https://doi.org/10.1073/pnas.1421055111>

Luuk H, Koks S, Plaas M, Hannibal J, Rehfeld JF, Vasar E (2008) Distribution of Wfs1 protein in the central nervous system of the mouse and its relation to clinical symptoms of the Wolfram syndrome. *J Comp Neurol* 509:642–660. <https://doi.org/10.1002/cne.21777>

Luuk H, Plaas M, Raud S, Innos J, Sütt S, Lasner H, Abramov U, Kurrikoff K, Kõks S, Vasar E (2009) Wfs1-deficient mice display impaired behavioural adaptation in stressful environment. *Behav Brain Res* 198:334–345. <https://doi.org/10.1016/j.bbr.2008.11.007>

Lyssenko V, Jonsson A, Almgren P, Pulizzi N, Isomaa B, Tuomi T, Berglund G, Altshuler D, Nilsson P, Groop L (2008) Clinical risk factors, DNA variants, and the development of type 2 diabetes. *N Engl J Med* 359:2220–2232. <https://doi.org/10.1056/NEJMoa0801869>

Marchand L, Li M, Leblicq C, Rafique I, Alarcon-Martinez T, Lange C, Rendon L, Tam E, Courville-Le Bouyonnet A, Polychronakos C (2021) Monogenic causes in the Type 1 diabetes genetics consortium cohort: low genetic risk for autoimmunity in case selection. *J Clin Endocrinol Metab* 106:1804–1810. <https://doi.org/10.1210/clinem/dgab056>

Maxwell KG, Augsornworawat P, Velazco-Cruz L, Kim MH, Asada R, Hogrebe NJ, Morikawa S, Urano F, Millman JR (2020) Gene-edited human stem cell-derived beta cells from a patient with monogenic diabetes reverse preexisting diabetes in mice. *Sci Transl Med* 12. <https://doi.org/10.1126/scitranslmed.aax9106>

Mets RB, Emery SB, Lesperance MM, Mets MB (2010) Congenital cataracts in two siblings with Wolfram syndrome. *Ophthalmic*

Genet 31:227–229. <https://doi.org/10.3109/13816810.2010.516056>

Morikawa S, Blacher L, Onwumere C, Urano F (2022) Loss of function of WFS1 causes ER stress-mediated inflammation in pancreatic beta-cells. *Front Endocrinol (Lausanne)* 13:849204. <https://doi.org/10.3389/fendo.2022.849204>

Morikawa S, Tajima T, Nakamura A, Ishizu K, Ariga T (2017) A novel heterozygous mutation of the WFS1 gene leading to constitutive endoplasmic reticulum stress is the cause of Wolfram syndrome. *Pediatr Diabetes* 18:934–941. <https://doi.org/10.1111/pedi.12513>

Nguyen LD, Fischer TT, Abreu D, Arroyo A, Urano F, Ehrlich BE (2020) Calpain inhibitor and ibudilast rescue beta cell functions in a cellular model of Wolfram syndrome. *Proc Natl Acad Sci U S A* 117:17389–17398. <https://doi.org/10.1073/pnas.2007136117>

Noormets K, Kõks S, Ivask M, Aunapuu M, Arend A, Vasar E, Tillmann V (2014) Energy metabolism and thyroid function of mice with deleted wolframin (Wfs1) gene. *Exp Clin Endocrinol Diabetes* 122:281–286. <https://doi.org/10.1055/s-0034-1372582>

Noormets K, Kõks S, Kavak A, Arend A, Aunapuu M, Keldrimaa A, Vasar E, Tillmann V (2009) Male mice with deleted Wolframin (Wfs1) gene have reduced fertility. *Reprod Biol Endocrinol* 7:82. <https://doi.org/10.1186/1477-7827-7-82>

Noormets K, Kõks S, Muldmaa M, Mäuring L, Vasar E, Tillmann V (2011) Sex differences in the development of diabetes in mice with deleted wolframin (Wfs1) gene. *Exp Clin Endocrinol Diabetes* 119:271–275. <https://doi.org/10.1055/s-0030-1265163>

Odisho T, Zhang L, Volchuk A (2015) ATF6 β regulates the Wfs1 gene and has a cell survival role in the ER stress response in pancreatic beta-cells. *Exp Cell Res* 330:111–122. <https://doi.org/10.1016/j.yexcr.2014.10.007>

Osowski CM, Hara T, O’Sullivan-Murphy B, Kanekura K, Lu S, Hara M, Ishigaki S, Zhu LJ, Hayashi E, Hui ST, Greiner D, Kaufman RJ, Bortell R, Urano F (2012) Thioredoxin-interacting protein mediates ER stress-induced beta cell death through initiation of the inflammasome. *Cell Metab* 16:265–273. <https://doi.org/10.1016/j.cmet.2012.07.005>

Osman AA, Saito M, Makepeace C, Permutt MA, Schlesinger P, Mueckler M (2003) Wolframin expression induces novel ion channel activity in endoplasmic reticulum membranes and increases intracellular calcium. *J Biol Chem* 278:52755–52762. <https://doi.org/10.1074/jbc.M310331200>

Paley RG, Tunbridge RE (1956) Primary optic atrophy in diabetes mellitus. *Diabetes* 5:295–296. <https://doi.org/10.2337/diab.5.4.295>

Panfili E, Frontino G, Pallotta MT (2023) GLP-1 receptor agonists as promising disease-modifying agents in WFS1 spectrum disorder. *Front Clin Diabetes Healthc* 4:1171091. <https://doi.org/10.3389/fcdhc.2023.1171091>

Panfili E, Mondanelli G, Orabona C, Belladonna ML, Gargaro M, Fallarino F, Orecchini E, Prontera P, Proietti E, Frontino G, Tirelli E, Iacono A, Vacca C, Puccetti P, Grohmann U, Esposito S, Pallotta MT (2021) Novel mutations in the WFS1 gene are associated with Wolfram syndrome and systemic inflammation. *Hum Mol Genet* 30:265–276. <https://doi.org/10.1093/hmg/ddab040>

Pickett KA, Duncan RP, Paciorkowski AR, Permutt MA, Marshall B, Hershey T, Earhart GM, Washington University Wolfram Study Group (2012) Balance impairment in individuals with Wolfram syndrome. *Gait Posture* 36:619–624. <https://doi.org/10.1016/j.gaitpost.2012.06.008>

Plaas M, Seppa K, Reimets R, Jagomäe T, Toots M, Koppel T, Vallisoo T, Nigul M, Heinla I, Meier R, Kaasik A, Piirsoo A, Hickey

494 MA, Terasmaa A, Vasar E (2017) Wfs1- deficient rats develop primary symptoms of Wolfram syndrome: insulin-dependent
 495 diabetes, optic nerve atrophy and medullary degeneration. *Sci Rep* 7:10220. <https://doi.org/10.1038/s41598-017-09392-x>
 496 Polymeropoulos MH, Swift RG, Swift M (1994) Linkage of the gene for Wolfram syndrome to markers on the short arm of
 497 chromosome 4. *Nat Genet* 8:95–97. <https://doi.org/10.1038/ng0994-95>
 498 Pourtoy-Brasselet S, Sciauvaud A, Boza-Moran MG, Cailleret M, Jarrige M, Polvèche H, Polentes J, Chevet E, Martinat C,
 499 Peschanski M, Aubry L (2021) Human iPSC-derived neurons reveal early developmental alteration of neurite outgrowth in the
 500 late-occurring neurodegenerative Wolfram syndrome. *Am J Hum Genet* 108:2171–2185.
 501 <https://doi.org/10.1016/j.ajhg.2021.10.001>
 502 Richard EM, Brun E, Korchagina J, Crouzier L, Affortit C, Alves S, Cazevielle C, Mausset-Bonnefont AL, Lenoir M, Puel JL,
 503 Maurice T, Thiry M, Wang J, Delprat B (2023) Wfs1(E864K) knock-in mice illuminate the fundamental role of Wfs1 in
 504 endocochlear potential production. *Cell Death Dis* 14:387. <https://doi.org/10.1038/s41419-023-05912-y>
 505 Riggs AC, Bernal-Mizrachi E, Ohsugi M, Wasson J, Fatrai S, Welling C, Murray J, Schmidt RE, Herrera PL, Permutt MA (2005)
 506 Mice conditionally lacking the Wolfram gene in pancreatic islet beta cells exhibit diabetes as a result of enhanced endoplasmic
 507 reticulum stress and apoptosis. *Diabetologia* 48:2313–2321. <https://doi.org/10.1007/s00125-005-1947-4>
 508 Rötig A, Cormier V, Chatelain P, Francois R, Saudubray JM, Rustin P, Munnich A (1993) Deletion of mitochondrial DNA in a case
 509 of early-onset diabetes mellitus, optic atrophy, and deafness (Wolfram syndrome, MIM 222300). *J Clin Invest* 91:1095–1098.
 510 <https://doi.org/10.1172/JCI116267>
 511 Sakakibara Y, Sekiya M, Fujisaki N, Quan X, Iijima KM (2018) Knockdown of wfs1, a fly homolog of Wolfram syndrome 1, in the
 512 nervous system increases susceptibility to age- and stress-induced neuronal dysfunction and degeneration in *Drosophila*. *PLOS*
 513 *Genet* 14:e1007196. <https://doi.org/10.1371/journal.pgen.1007196>
 514 Sandhu MS, Weedon MN, Fawcett KA, Wasson J, Debenham SL, Daly A, Lango H, Frayling TM, Neumann RJ, Sherva R, Blech I,
 515 Pharoah PD, Palmer CN, Kimber C, Tavendale R, Morris AD, McCarthy MI, Walker M, Hitman G, Glaser B, Permutt MA,
 516 Hattersley AT, Wareham NJ, Barroso I (2007) Common variants in WFS1 confer risk of type 2 diabetes. *Nat Genet* 39:951–953.
 517 <https://doi.org/10.1038/ng2067>
 518 Sedman T, Rünkorg K, Krass M, Luuk H, Plaas M, Vasar E, Volke V (2016) Exenatide is an effective antihyperglycaemic agent in
 519 a mouse model of Wolfram syndrome 1. *J Diabetes Res* 2016:9239530. <https://doi.org/10.1155/2016/9239530>
 520 Seppa K, Jagomäe T, Kukker KG, Reimets R, Pastak M, Vasar E, Terasmaa A, Plaas M (2021) Liraglutide, 7,8-DHF and their co-
 521 treatment prevents loss of vision and cognitive decline in a Wolfram syndrome rat model. *Sci Rep* 11:2275.
 522 <https://doi.org/10.1038/s41598-021-81768-6>
 523 Seppa K, Toots M, Reimets R, Jagomae T, Koppel T, Pallase M, Hasselholt S, Krogsbaek Mikkelsen M, Randel Nyengaard J, Vasar
 524 E, Terasmaa A, Plaas M (2019) GLP-1 receptor agonist liraglutide has a neuroprotective effect on an aged rat model of Wolfram
 525 syndrome. *Sci Rep* 31:15742
 526 Shang L, Hua H, Foo K, Martinez H, Watanabe K, Zimmer M, Kahler DJ, Freeby M, Chung W, LeDuc C, Goland R, Leibel RL, Egli
 527 D (2014) β -cell dysfunction due to increased ER stress in a stem cell model of Wolfram syndrome. *Diabetes* 63:923–933.
 528 <https://doi.org/10.2337/db13-0717>
 529 Smith CJ, Crock PA, King BR, Meldrum CJ, Scott RJ (2004) Phenotype-genotype correlations in a series of wolfram syndrome

families. *Diabetes Care* 27:2003–2009. <https://doi.org/10.2337/diacare.27.8.2003>

Sparsø T, Andersen G, Albrechtsen A, Jørgensen T, Borch-Johnsen K, Sandbaek A, Lauritzen T, Wasson J, Permutt MA, Glaser B, Madsbad S, Pedersen O, Hansen T (2008) Impact of polymorphisms in WFS1 on prediabetic phenotypes in a population-based sample of middle-aged people with normal and abnormal glucose regulation. *Diabetologia* 51:1646–1652. <https://doi.org/10.1007/s00125-008-1064-2>

Strom TM, Hörtnagel K, Hofmann S, Gekeler F, Scharfe C, Rabl W, Gerbitz KD, Meitinger T (1998) Diabetes insipidus, diabetes mellitus, optic atrophy and deafness (DIDMOAD) caused by mutations in a novel gene (wolframin) coding for a predicted transmembrane protein. *Hum Mol Genet* 7:2021–2028. <https://doi.org/10.1093/hmg/7.13.2021>

Takeda K, Inoue H, Tanizawa Y, Matsuzaki Y, Oba J, Watanabe Y, Shinoda K, Oka Y (2001) WFS1 (Wolfram syndrome 1) gene product: predominant subcellular localization to endoplasmic reticulum in cultured cells and neuronal expression in rat brain. *Hum Mol Genet* 10:477–484. <https://doi.org/10.1093/hmg/10.5.477>

Takei D, Ishihara H, Yamaguchi S, Yamada T, Tamura A, Katagiri H, Maruyama Y, Oka Y (2006) WFS1 protein modulates the free Ca(2+) concentration in the endoplasmic reticulum. *FEBS Lett* 580:5635–5640. <https://doi.org/10.1016/j.febslet.2006.09.007>

Toots M, Seppa K, Jagomäe T, Koppel T, Pallase M, Heinla I, Terasmaa A, Plaas M, Vasar E (2018) Preventive treatment with liraglutide protects against development of glucose intolerance in a rat model of Wolfram syndrome. *Sci Rep* 8:10183. <https://doi.org/10.1038/s41598-018-28314-z>

Ueda K, Kawano J, Takeda K, Yujiri T, Tanabe K, Anno T, Akiyama M, Nozaki J, Yoshinaga T, Koizumi A, Shinoda K, Oka Y, Tanizawa Y (2005) Endoplasmic reticulum stress induces Wfs1 gene expression in pancreatic beta-cells via transcriptional activation. *Eur J Endocrinol* 153:167–176. <https://doi.org/10.1530/eje.1.01945>

van Hoek M, Dehghan A, Witteman JC, van Duijn CM, Uitterlinden AG, Oostra BA, Hofman A, Sijbrands EJ, Janssens AC (2008) Predicting type 2 diabetes based on polymorphisms from genome-wide association studies: a population-based study. *Diabetes* 57:3122–3128. <https://doi.org/10.2337/db08-0425>

Visnapuu T, Plaas M, Reimets R, Raud S, Terasmaa A, Kõks S, Sütt S, Luuk H, Hundahl CA, Eskla KL, Altpere A, Altko A, Harro J, Vasar E (2013a) Evidence for impaired function of dopaminergic system in Wfs1-deficient mice. *Behav Brain Res* 244:90–99. <https://doi.org/10.1016/j.bbr.2013.01.046>

Visnapuu T, Raud S, Loomets M, Reimets R, Sütt S, Luuk H, Plaas M, Kõks S, Volke V, Altko A, Harro J, Vasar E (2013b) Wfs1-deficient mice display altered function of serotonergic system and increased behavioral response to antidepressants. *Front Neurosci* 7:132. <https://doi.org/10.3389/fnins.2013.00132>

Vora AJ, Lilleyman JS (1993) Wolfram syndrome: mitochondrial disorder. *Lancet* 342:1059. [https://doi.org/10.1016/0140-6736\(93\)92918-J](https://doi.org/10.1016/0140-6736(93)92918-J)

Wang L, Liu H, Zhang X, Song E, Wang Y, Xu T, Li Z (2021) WFS1 functions in ER export of vesicular cargo proteins in pancreatic beta-cells. *Nat Commun* 12:6996. <https://doi.org/10.1038/s41467-021-27344-y>

Waszczykowska A, Zmysłowska A, Braun M, Ivask M, Koks S, Jurowski P, Młynarski W (2020) Multiple retinal anomalies in Wfs1-deficient mice. *Diagnostics (Basel)* 10. <https://doi.org/10.3390/diagnostics10090607>

Westermarck P, Neuman RJ, Wasson J, Atzmon G, Wainstein J, Yerushalmi Y, Cohen J, Barzilai N, Blech I, Glaser B, Permutt MA (2010) Gene-gene interactions lead to higher risk for development of Type 2 diabetes in an Ashkenazi Jewish population. *PLOS*

ONE 5

Wolfram DJ, Wagener HP (1938) Diabetes mellitus and simple optic atrophy among siblings: report of four cases. *Mayo Clin Proc* 13:715–718

Yamada T, Ishihara H, Tamura A, Takahashi R, Yamaguchi S, Takei D, Tokita A, Satake C, Tashiro F, Katagiri H, Aburatani H, Miyazaki J, Oka Y (2006) WFS1-deficiency increases endoplasmic reticulum stress, impairs cell cycle progression and triggers the apoptotic pathway specifically in pancreatic beta-cells. *Hum Mol Genet* 15:1600–1609. <https://doi.org/10.1093/hmg/ddl081>

Young TL, Ives E, Lynch E, Person R, Snook S, MacLaren L, Cater T, Griffin A, Fernandez B, Lee MK, King MC (2001) Non-syndromic progressive hearing loss DFNA38 is caused by heterozygous missense mutation in the Wolfram syndrome gene WFS1. *Hum Mol Genet* 10:2509–2514. <https://doi.org/10.1093/hmg/10.22.2509>

Zatyka M, Da Silva Xavier G, Bellomo EA, Leadbeater W, Astuti D, Smith J, Michelangeli F, Rutter GA, Barrett TG (2015) Sarco(endo)plasmic reticulum ATPase is a molecular partner of Wolfram syndrome 1 protein, which negatively regulates its expression. *Hum Mol Genet* 24:814–827. <https://doi.org/10.1093/hmg/ddu499>

Zatyka M, Ricketts C, da Silva Xavier G, Minton J, Fenton S, Hofmann-Thiel S, Rutter GA, Barrett TG (2008) Sodium-potassium ATPase 1 subunit is a molecular partner of Wolframin, an endoplasmic reticulum protein involved in ER stress. *Hum Mol Genet* 17:190–200. <https://doi.org/10.1093/hmg/ddm296>

Zatyka M, Rosenstock TR, Sun C, Palhegyi AM, Hughes GW, Lara-Reyna S, Astuti D, di Maio A, Sciauvaud A, Korsgen ME, Stanulovic V, Kocak G, Rak M, Pourtoy-Brasselet S, Winter K, Varga T, Jarrige M, Polvèche H, Correia J, Frickel EM, Hoogenkamp M, Ward DG, Aubry L, Barrett T, Sarkar S (2023) Depletion of WFS1 compromises mitochondrial function in hiPSC-derived neuronal models of Wolfram syndrome. *Stem Cell Rep* 18:1090–1106. <https://doi.org/10.1016/j.stemcr.2023.04.002>

Abreu D, Asada R, Revilla JMP, Lavagnino Z, Kries K, Piston DW, Urano F (2020) Wolfram syndrome 1 gene regulates pathways maintaining beta-cell health and survival. *Lab Invest* 10.1038/s41374-020-0408-5

Abreu D, Stone SI, Pearson TS, Bucelli RC, Simpson AN, Hurst S, Brown CM, Kries K, Onwumere C, Gu H, Hoekel J, Tychsen L, Van Stavern GP, White NH, Marshall BA, Hershey T, Urano F (2021) A phase Ib/Ila clinical trial of dantrolene sodium in patients with Wolfram syndrome. *JCI Insight* 6

Akiyama M, Hatanaka M, Ohta Y, Ueda K, Yanai A, Uehara Y, Tanabe K, Tsuru M, Miyazaki M, Saeki S, Saito T, Shinoda K, Oka Y, Tanizawa Y (2009) Increased insulin demand promotes while pioglitazone prevents pancreatic beta cell apoptosis in Wfs1 knockout mice. *Diabetologia* 52, 653–663

Angebault C, Fauconnier J, Patergnani S, Rieusset J, Danese A, Affortit CA, Jagodzinska J, Megy C, Quiles M, Cazevielle C, Korchagina J, Bonnet-Wersinger D, Milea D, Hamel C, Pinton P, Thiry M, Lacampagne A, Delprat B, Delettre C (2018) ER-mitochondria cross-talk is regulated by the Ca(2+) sensor NCS1 and is impaired in Wolfram syndrome. *Sci Signal* 11

Bansal V, Boehm BO, Darvasi A (2018) Identification of a missense variant in the WFS1 gene that causes a mild form of Wolfram syndrome and is associated with risk for type 2 diabetes in Ashkenazi Jewish individuals. *Diabetologia* 10.1007/s00125-018-4690-3

Barrett TG, Bunday SE, Macleod AF (1995) Neurodegeneration and diabetes: UK nationwide study of Wolfram (DIDMOAD)

syndrome. *Lancet* 346, 1458-1463

Bespalova IN, Van Camp G, Bom SJ, Brown DJ, Cryns K, DeWan AT, Erson AE, Flothmann K, Kunst HP, Kurnool P, Sivakumaran TA, Cremers CW, Leal SM, Burmeister M, Lesperance MM (2001) Mutations in the Wolfram syndrome 1 gene (WFS1) are a common cause of low frequency sensorineural hearing loss. *Hum Mol Genet* 10, 2501-2508

Blackstone C, Cagalinec M, Liiv M, Hodurova Z, Hickey MA, Vaarmann A, Mandel M, Zeb A, Choubey V, Kuun M, Safiulina D, Vasar E, Veksler V, Kaasik A (2016) Role of Mitochondrial Dynamics in Neuronal Development: Mechanism for Wolfram Syndrome. *PLOS Biology* 14

Bonnet Wersinger D, Benkafadar N, Jagodzinska J, Hamel C, Tanizawa Y, Lenaers G, Delettre C (2014) Impairment of visual function and retinal ER stress activation in *Wfs1*-deficient mice. *PLoS One* 9, e97222

Bu X, Rotter JI (1993) Wolfram syndrome: a mitochondrial-mediated disorder? *Lancet* 342, 598-600

Bundey S, Fielder A, Poulton K (1993) Wolfram syndrome: mitochondrial disorder. *Lancet* 342, 1059-1060

Cairns G, Burte F, Price R, O'Connor E, Toms M, Mishra R, Moosajee M, Pyle A, Sayer JA, Yu-Wai-Man P (2021) A mutant *wfs1* zebrafish model of Wolfram syndrome manifesting visual dysfunction and developmental delay. *Sci Rep* 11, 20491

Cheng S, Wu Y, Wu W, Zhang D (2013) Association of rs734312 and rs10010131 polymorphisms in WFS1 gene with type 2 diabetes mellitus: a meta-analysis. *Endocr J* 60, 441-447

Cheurfa N, Brenner GM, Reis AF, Dubois-Laforgue D, Roussel R, Tichet J, Lantieri O, Balkau B, Fumeron F, Timsit J, Marre M, Velho G (2011) Decreased insulin secretion and increased risk of type 2 diabetes associated with allelic variations of the WFS1 gene: the Data from Epidemiological Study on the Insulin Resistance Syndrome (DESIR) prospective study. *Diabetologia* 54, 554-562

Clark AL, Kanekura K, Lavagnino Z, Spears LD, Abreu D, Mahadevan J, Yagi T, Semenkovich CF, Piston DW, Urano F (2017) Targeting Cellular Calcium Homeostasis to Prevent Cytokine-Mediated Beta Cell Death. *Sci Rep* 7, 5611

Cremers CW, Wijdeveld PG, Pinckers AJ (1977) Juvenile diabetes mellitus, optic atrophy, hearing loss, diabetes insipidus, atonia of the urinary tract and bladder, and other abnormalities (Wolfram syndrome). A review of 88 cases from the literature with personal observations on 3 new patients. *Acta Paediatr Scand Suppl* 10.1111/j.1651-2227.1977.tb15069.x, 1-16

Crouzier L, Danese A, Yasui Y, Richard EM, Lievens JC, Patergnani S, Couly S, Diez C, Denus M, Cubedo N, Rossel M, Thiry M, Su TP, Pinton P, Maurice T, Delprat B (2022a) Activation of the sigma-1 receptor chaperone alleviates symptoms of Wolfram syndrome in preclinical models. *Sci Transl Med* 14, eab3763

Crouzier L, Richard EM, Diez C, Alzaeem H, Denus M, Cubedo N, Delaunay T, Glendenning E, Baxendale S, Lievens JC, Whitfield TT, Maurice T, Delprat B (2022b) Morphological, behavioral and cellular analyses revealed different phenotypes in Wolfram syndrome *wfs1a* and *wfs1b* zebrafish mutant lines. *Hum Mol Genet* 31, 2711-2727

Cryns K, Pfister M, Pennings RJ, Bom SJ, Flothmann K, Caethoven G, Kremer H, Schatteman I, Koln KA, Toth T, Kupka S, Blin N, Nurnberg P, Thiele H, van de Heyning PH, Reardon W, Stephens D, Cremers CW, Smith RJ, Van Camp G (2002) Mutations in the WFS1 gene that cause low-frequency sensorineural hearing loss are small non-inactivating mutations. *Human genetics* 110, 389-394

De Franco E, Flanagan SE, Yagi T, Abreu D, Mahadevan J, Johnson MB, Jones G, Acosta F, Mulaudzi M, Lek N, Oh V, Petz O, Caswell R, Ellard S, Urano F, Hattersley AT (2017) Dominant ER Stress-Inducing WFS1 Mutations Underlie a Genetic Syndrome

638 of Neonatal/Infancy-Onset Diabetes, Congenital Sensorineural Deafness, and Congenital Cataracts. *Diabetes* 66, 2044-2053
 639 Elek Z, Nemeth N, Nagy G, Nemeth H, Somogyi A, Hosszufalusi N, Sasvari-Szekely M, Ronai Z (2015) Micro-RNA Binding Site
 640 Polymorphisms in the WFS1 Gene Are Risk Factors of Diabetes Mellitus. *PLoS One* 10, e0139519
 641 Fonseca SG, Fukuma M, Lipson KL, Nguyen LX, Allen JR, Oka Y, Urano F (2005) WFS1 is a novel component of the unfolded
 642 protein response and maintains homeostasis of the endoplasmic reticulum in pancreatic beta-cells. *J Biol Chem* 280, 39609-
 643 39615
 644 Fonseca SG, Ishigaki S, Osowski CM, Lu S, Lipson KL, Ghosh R, Hayashi E, Ishihara H, Oka Y, Permutt MA, Urano F (2010)
 645 Wolfram syndrome 1 gene negatively regulates ER stress signaling in rodent and human cells. *The Journal of clinical*
 646 *investigation* 120, 744-755
 647 Gharanei S, Zatyka M, Astuti D, Fenton J, Sik A, Nagy Z, Barrett TG (2013) Vacuolar-type H⁺-ATPase V1A subunit is a molecular
 648 partner of Wolfram syndrome 1 (WFS1) protein, which regulates its expression and stability. *Hum Mol Genet* 22, 203-217
 649 Gorgogietas V, Rajaei B, Heeyoung C, Santacreu BJ, Marin-Canas S, Salpea P, Sawatani T, Musuaya A, Arroyo MN, Moreno-
 650 Castro C, Benabdallah K, Demarez C, Toivonen S, Cosentino C, Pachera N, Lytrivi M, Cai Y, Cernel L, Brown C, Urano F, Marchetti
 651 P, Gilon P, Eizirik DL, Cnop M, Igoillo-Esteve M (2023) GLP-1R agonists demonstrate potential to treat Wolfram syndrome in
 652 human preclinical models. *Diabetologia* 66, 1306-1321
 653 Hatanaka M, Tanabe K, Yanai A, Ohta Y, Kondo M, Akiyama M, Shinoda K, Oka Y, Tanizawa Y (2011) Wolfram syndrome 1 gene
 654 (WFS1) product localizes to secretory granules and determines granule acidification in pancreatic beta-cells. *Hum Mol Genet*
 655 20, 1274-1284
 656 Heni M, Ketterer C, Thamer C, Herzberg-Schafer SA, Guthoff M, Stefan N, Machicao F, Staiger H, Fritsche A, Haring HU (2010)
 657 Glycemia determines the effect of type 2 diabetes risk genes on insulin secretion. *Diabetes* 59, 3247-3252
 658 Hershey T, Lugar HM, Shimony JS, Rutlin J, Koller JM, Perantie DC, Paciorkowski AR, Eisenstein SA, Permutt MA, Washington
 659 University Wolfram Study G (2012) Early brain vulnerability in Wolfram syndrome. *PLoS One* 7, e40604
 660 Hofmann S, Philbrook C, Gerbitz KD, Bauer MF (2003) Wolfram syndrome: structural and functional analyses of mutant and
 661 wild-type wolframin, the WFS1 gene product. *Hum Mol Genet* 12, 2003-2012
 662 Inoue H, Tanizawa Y, Wasson J, Behn P, Kalidas K, Bernal-Mizrachi E, Mueckler M, Marshall H, Donis-Keller H, Crock P, Rogers
 663 D, Mikuni M, Kumashiro H, Higashi K, Sobue G, Oka Y, Permutt MA (1998) A gene encoding a transmembrane protein is
 664 mutated in patients with diabetes mellitus and optic atrophy (Wolfram syndrome). *Nature Genetics* 20, 143-148
 665 Ishihara H, Takeda S, Tamura A, Takahashi R, Yamaguchi S, Takei D, Yamada T, Inoue H, Soga H, Katagiri H, Tanizawa Y, Oka Y
 666 (2004) Disruption of the WFS1 gene in mice causes progressive beta-cell loss and impaired stimulus-secretion coupling in
 667 insulin secretion. *Hum Mol Genet* 13, 1159-1170
 668 Ivask M, Volke V, Raasmaja A, Koks S (2021) High-fat diet associated sensitization to metabolic stress in Wfs1 heterozygous
 669 mice. *Mol Genet Metab* 134, 203-211
 670 Jagomae T, Seppa K, Reimets R, Pastak M, Plaas M, Hickey MA, Kukker KG, Moons L, De Groef L, Vasar E, Kaasik A, Terasmaa
 671 A, Plaas M (2021) Early Intervention and Lifelong Treatment with GLP1 Receptor Agonist Liraglutide in a Wolfram Syndrome
 672 Rat Model with an Emphasis on Visual Neurodegeneration, Sensorineural Hearing Loss and Diabetic Phenotype. *Cells* 10
 673 Kakiuchi C, Ishiwata M, Hayashi A, Kato T (2006) XBP1 induces WFS1 through an endoplasmic reticulum stress response

674 element-like motif in SH-SY5Y cells. *J Neurochem* 97, 545-555
 675 Kato T, Ishiwata M, Yamada K, Kasahara T, Kakiuchi C, Iwamoto K, Kawamura K, Ishihara H, Oka Y (2008) Behavioral and gene
 676 expression analyses of Wfs1 knockout mice as a possible animal model of mood disorder. *Neurosci Res* 61, 143-158
 677 Kinsley BT SM, Dumont RH, Swift RG, (1995) Morbidity and mortality in the Wolfram syndrome. *Diabetes Care* 18, 1566-1570
 678 Kitamura RA, Maxwell KG, Ye W, Kries K, Brown CM, Augsornworawat P, Hirsch Y, Johansson MM, Weiden T, Ekstein J, Cohen
 679 J, Klee J, Leslie K, Simeonov A, Henderson MJ, Millman JR, Urano F (2022) Multidimensional analysis and therapeutic
 680 development using patient iPSC-derived disease models of Wolfram syndrome. *JCI Insight* 7
 681 Kobayashi M, Miyagawa M, Nishio SY, Moteki H, Fujikawa T, Ohyama K, Sakaguchi H, Miyanochara I, Sugaya A, Naito Y, Morita
 682 SY, Kanda Y, Takahashi M, Ishikawa K, Nagano Y, Tono T, Oshikawa C, Kihara C, Takahashi H, Noguchi Y, Usami SI (2018) WFS1
 683 mutation screening in a large series of Japanese hearing loss patients: Massively parallel DNA sequencing-based analysis. *PLoS*
 684 *One* 13, e0193359
 685 Koks S, Soomets U, Paya-Cano JL, Fernandes C, Luuk H, Plaas M, Terasmaa A, Tillmann V, Noormets K, Vasar E, Schalkwyk LC
 686 (2009) Wfs1 gene deletion causes growth retardation in mice and interferes with the growth hormone pathway. *Physiol*
 687 *Genomics* 37, 249-259
 688 Kondo M, Tanabe K, Amo-Shiinoki K, Hatanaka M, Morii T, Takahashi H, Seino S, Yamada Y, Tanizawa Y (2018) Activation of
 689 GLP-1 receptor signalling alleviates cellular stresses and improves beta cell function in a mouse model of Wolfram syndrome.
 690 *Diabetologia* 61, 2189-2201
 691 Kunnappallil RS, Hasan G (2022) 10.1101/2022.11.10.515972
 692 Kurimoto J, Takagi H, Miyata T, Hodai Y, Kawaguchi Y, Hagiwara D, Suga H, Kobayashi T, Sugiyama M, Onoue T, Ito Y, Iwama S,
 693 Banno R, Tanabe K, Tanizawa Y, Arima H (2021) Deficiency of WFS1 leads to the impairment of AVP secretion under
 694 dehydration in male mice. *Pituitary* 24, 582-588
 695 La Morgia C, Maresca A, Amore G, Gramegna LL, Carbonelli M, Scimonelli E, Danese A, Patergnani S, Caporali L, Tagliavini F,
 696 Del Dotto V, Capristo M, Sadun F, Barboni P, Savini G, Evangelisti S, Bianchini C, Valentino ML, Liguori R, Tonon C, Giorgi C,
 697 Pinton P, Lodi R, Carelli V (2020) Calcium mishandling in absence of primary mitochondrial dysfunction drives cellular
 698 pathology in Wolfram Syndrome. *Sci Rep* 10, 4785
 699 Lee YH, Kang ES, Kim SH, Han SJ, Kim CH, Kim HJ, Ahn CW, Cha BS, Nam M, Nam CM, Lee HC (2008) Association between
 700 polymorphisms in SLC30A8, HHEX, CDKN2A/B, IGF2BP2, FTO, WFS1, CDKAL1, KCNQ1 and type 2 diabetes in the Korean
 701 population. *Journal of human genetics* 53, 991-998
 702 Lerner AG, Upton JP, Praveen PV, Ghosh R, Nakagawa Y, Igbaria A, Shen S, Nguyen V, Backes BJ, Heiman M, Heintz N, Greengard
 703 P, Hui S, Tang Q, Trusina A, Oakes SA, Papa FR (2012) IRE1alpha induces thioredoxin-interacting protein to activate the NLRP3
 704 inflammasome and promote programmed cell death under irremediable ER stress. *Cell Metab* 16, 250-264
 705 Li M, Wang S, Xu K, Chen Y, Fu Q, Gu Y, Shi Y, Zhang M, Sun M, Chen H, Han X, Li Y, Tang Z, Cai L, Li Z, Shi Y, Yang T, Polychronakos
 706 C (2020) High Prevalence of a Monogenic Cause in Han Chinese Diagnosed With Type 1 Diabetes, Partly Driven by
 707 Nonsyndromic Recessive WFS1 Mutations. *Diabetes* 69, 121-126
 708 Lipson KL, Fonseca SG, Ishigaki S, Nguyen LX, Foss E, Bortell R, Rossini AA, Urano F (2006) Regulation of insulin biosynthesis in
 709 pancreatic beta cells by an endoplasmic reticulum-resident protein kinase IRE1. *Cell Metab* 4, 245-254

710 Long J, Edwards T, Signorello LB, Cai Q, Zheng W, Shu XO, Blot WJ (2012) Evaluation of genome-wide association study-
 711 identified type 2 diabetes loci in African Americans. *Am J Epidemiol* 176, 995-1001
 712 Lu S, Kanekura K, Hara T, Mahadevan J, Spears LD, Oslowski CM, Martinez R, Yamazaki-Inoue M, Toyoda M, Neilson A, Blanner
 713 P, Brown CM, Semenkovich CF, Marshall BA, Hershey T, Umezawa A, Greer PA, Urano F (2014) A calcium-dependent protease
 714 as a potential therapeutic target for Wolfram syndrome. *Proc Natl Acad Sci U S A* 111, E5292-5301
 715 Luuk H, Koks S, Plaas M, Hannibal J, Rehfeld JF, Vasar E (2008) Distribution of Wfs1 protein in the central nervous system of
 716 the mouse and its relation to clinical symptoms of the Wolfram syndrome. *J Comp Neurol* 509, 642-660
 717 Luuk H, Plaas M, Raud S, Innos J, Sutt S, Lasner H, Abramov U, Kurrikoff K, Koks S, Vasar E (2009) Wfs1-deficient mice display
 718 impaired behavioural adaptation in stressful environment. *Behav Brain Res* 198, 334-345
 719 Lyssenko V, Jonsson A, Almgren P, Pulizzi N, Isomaa B, Tuomi T, Berglund G, Altshuler D, Nilsson P, Groop L (2008) Clinical risk
 720 factors, DNA variants, and the development of type 2 diabetes. *N Engl J Med* 359, 2220-2232
 721 Marchand L, Li M, Leblicq C, Rafique I, Alarcon-Martinez T, Lange C, Rendon L, Tam E, Courville-Le Bouyonnec A, Polychronakos
 722 C (2021) Monogenic Causes in the Type 1 Diabetes Genetics Consortium Cohort: Low Genetic Risk for Autoimmunity in Case
 723 Selection. *J Clin Endocrinol Metab* 106, 1804-1810
 724 Maxwell KG, Augsornworawat P, Velazco-Cruz L, Kim MH, Asada R, Hogrebe NJ, Morikawa S, Urano F, Millman JR (2020) Gene-
 725 edited human stem cell-derived beta cells from a patient with monogenic diabetes reverse preexisting diabetes in mice. *Sci*
 726 *Transl Med* 12
 727 Mets RB, Emery SB, Lesperance MM, Mets MB (2010) Congenital cataracts in two siblings with Wolfram syndrome. *Ophthalmic*
 728 *Genet* 31, 227-229
 729 Morikawa S, Blacher L, Onwumere C, Urano F (2022) Loss of Function of WFS1 Causes ER Stress-Mediated Inflammation in
 730 Pancreatic Beta-Cells. *Front Endocrinol (Lausanne)* 13, 849204
 731 Morikawa S, Tajima T, Nakamura A, Ishizu K, Ariga T (2017) A novel heterozygous mutation of the WFS1 gene leading to
 732 constitutive endoplasmic reticulum stress is the cause of Wolfram syndrome. *Pediatric diabetes* 18, 934-941
 733 Nguyen LD, Fischer TT, Abreu D, Arroyo A, Urano F, Ehrlich BE (2020) Calpain inhibitor and ibudilast rescue beta cell functions
 734 in a cellular model of Wolfram syndrome. *Proc Natl Acad Sci U S A* 10.1073/pnas.2007136117
 735 Noormets K, Koks S, Ivask M, Aunapuu M, Arend A, Vasar E, Tillmann V (2014) Energy metabolism and thyroid function of
 736 mice with deleted wolframin (Wfs1) gene. *Exp Clin Endocrinol Diabetes* 122, 281-286
 737 Noormets K, Koks S, Kavak A, Arend A, Aunapuu M, Keldrimaa A, Vasar E, Tillmann V (2009) Male mice with deleted Wolframin
 738 (Wfs1) gene have reduced fertility. *Reprod Biol Endocrinol* 7, 82
 739 Noormets K, Koks S, Muldmaa M, Muring L, Vasar E, Tillmann V (2011) Sex differences in the development of diabetes in
 740 mice with deleted wolframin (Wfs1) gene. *Exp Clin Endocrinol Diabetes* 119, 271-275
 741 Odisho T, Zhang L, Volchuk A (2015) ATF6 β regulates the Wfs1 gene and has a cell survival role in the ER stress response in
 742 pancreatic beta-cells. *Experimental cell research* 330, 111-122
 743 Oslowski CM, Hara T, O'Sullivan-Murphy B, Kanekura K, Lu S, Hara M, Ishigaki S, Zhu LJ, Hayashi E, Hui ST, Greiner D, Kaufman
 744 RJ, Bortell R, Urano F (2012) Thioredoxin-interacting protein mediates ER stress-induced beta cell death through initiation of
 745 the inflammasome. *Cell Metab* 16, 265-273

746 Osman AA, Saito M, Makepeace C, Permutt MA, Schlesinger P, Mueckler M (2003) Wolframin expression induces novel ion
 747 channel activity in endoplasmic reticulum membranes and increases intracellular calcium. *J Biol Chem* 278, 52755-52762
 748 Paley RG, Tunbridge RE (1956) Primary optic atrophy in diabetes mellitus. *Diabetes* 5, 295-296
 749 Panfili E, Frontino G, Pallotta MT (2023) GLP-1 receptor agonists as promising disease-modifying agents in WFS1 spectrum
 750 disorder. *Front Clin Diabetes Healthc* 4, 1171091
 751 Panfili E, Mondanelli G, Orabona C, Belladonna ML, Gargaro M, Fallarino F, Orecchini E, Prontera P, Proietti E, Frontino G, Tirelli
 752 E, Iacono A, Vacca C, Puccetti P, Grohmann U, Esposito S, Pallotta MT (2021) Novel mutations in the WFS1 gene are associated
 753 with Wolfram syndrome and systemic inflammation. *Hum Mol Genet* 30, 265-276
 754 Pickett KA, Duncan RP, Paciorkowski AR, Permutt MA, Marshall B, Hershey T, Earhart GM, Washington University Wolfram
 755 Study G (2012) Balance impairment in individuals with Wolfram syndrome. *Gait Posture* 36, 619-624
 756 Plaas M, Seppa K, Reimets R, Jagomae T, Toots M, Koppel T, Vallisoo T, Nigul M, Heinla I, Meier R, Kaasik A, Piirsoo A, Hickey
 757 MA, Terasmaa A, Vasar E (2017) Wfs1- deficient rats develop primary symptoms of Wolfram syndrome: insulin-dependent
 758 diabetes, optic nerve atrophy and medullary degeneration. *Sci Rep* 7, 10220
 759 Polymeropoulos MH, Swift RG, Swift M (1994) Linkage of the gene for Wolfram syndrome to markers on the short arm of
 760 chromosome 4. *Nature genetics* 8, 95-97
 761 Pourtoy-Brasselet S, Sciauvaud A, Boza-Moran MG, Cailleret M, Jarrige M, Polveche H, Polentes J, Chevet E, Martinat C,
 762 Peschanski M, Aubry L (2021) Human iPSC-derived neurons reveal early developmental alteration of neurite outgrowth in the
 763 late-occurring neurodegenerative Wolfram syndrome. *Am J Hum Genet* 108, 2171-2185
 764 Richard EM, Brun E, Korchagina J, Crouzier L, Affortit C, Alves S, Cazevielle C, Mausset-Bonnefont AL, Lenoir M, Puel JL,
 765 Maurice T, Thiry M, Wang J, Delprat B (2023) Wfs1(E864K) knock-in mice illuminate the fundamental role of Wfs1 in
 766 endocochlear potential production. *Cell Death Dis* 14, 387
 767 Riggs AC, Bernal-Mizrachi E, Ohsugi M, Wasson J, Fatrai S, Welling C, Murray J, Schmidt RE, Herrera PL, Permutt MA (2005)
 768 Mice conditionally lacking the Wolfram gene in pancreatic islet beta cells exhibit diabetes as a result of enhanced endoplasmic
 769 reticulum stress and apoptosis. *Diabetologia* 48, 2313-2321
 770 Rötig A, Cormier V, Chatelain P, Francois R, Saudubray JM, Rustin P, Munnich A (1993) Deletion of mitochondrial DNA in a case
 771 of early-onset diabetes mellitus, optic atrophy, and deafness (Wolfram syndrome, MIM 222300). *The Journal of clinical*
 772 *investigation* 91, 1095-1098
 773 Sakakibara Y, Sekiya M, Fujisaki N, Quan X, Iijima KM (2018) Knockdown of wfs1, a fly homolog of Wolfram syndrome 1, in the
 774 nervous system increases susceptibility to age- and stress-induced neuronal dysfunction and degeneration in *Drosophila*. *PLoS*
 775 *Genet* 14, e1007196
 776 Sandhu MS, Weedon MN, Fawcett KA, Wasson J, Debenham SL, Daly A, Lango H, Frayling TM, Neumann RJ, Sherva R, Blech I,
 777 Pharoah PD, Palmer CN, Kimber C, Tavendale R, Morris AD, McCarthy MI, Walker M, Hitman G, Glaser B, Permutt MA,
 778 Hattersley AT, Wareham NJ, Barroso I (2007) Common variants in WFS1 confer risk of type 2 diabetes. *Nature genetics* 39,
 779 951-953
 780 Sedman T, Runkorg K, Krass M, Luuk H, Plaas M, Vasar E, Volke V (2016) Exenatide Is an Effective Antihyperglycaemic Agent in
 781 a Mouse Model of Wolfram Syndrome 1. *J Diabetes Res* 2016, 9239530

782 Seppa K, Jagomae T, Kukker KG, Reimets R, Pastak M, Vasar E, Terasmaa A, Plaas M (2021) Liraglutide, 7,8-DHF and their co-
783 treatment prevents loss of vision and cognitive decline in a Wolfram syndrome rat model. *Sci Rep* 11, 2275

784 Seppa K, Toots M, Reimets R, Jagomae T, Koppel T, Pallase M, Hasselholt S, Krogsbaek Mikkelsen M, Randel Nyengaard J, Vasar
785 E, Terasmaa A, Plaas M (2019) GLP-1 receptor agonist liraglutide has a neuroprotective effect on an aged rat model of Wolfram
786 syndrome. *Sci Rep* 9, 15742

787 Shang L, Hua H, Foo K, Martinez H, Watanabe K, Zimmer M, Kahler DJ, Freeby M, Chung W, LeDuc C, Goland R, Leibel RL, Egli
788 D (2014) beta-cell dysfunction due to increased ER stress in a stem cell model of Wolfram syndrome. *Diabetes* 63, 923-933

789 Smith CJ, Crock PA, King BR, Meldrum CJ, Scott RJ (2004) Phenotype-genotype correlations in a series of wolfram syndrome
790 families. *Diabetes Care* 27, 2003-2009

791 Sparso T, Andersen G, Albrechtsen A, Jorgensen T, Borch-Johnsen K, Sandbaek A, Lauritzen T, Wasson J, Permutt MA, Glaser B,
792 Madsbad S, Pedersen O, Hansen T (2008) Impact of polymorphisms in WFS1 on prediabetic phenotypes in a population-based
793 sample of middle-aged people with normal and abnormal glucose regulation. *Diabetologia* 51, 1646-1652

794 Strom TM, Hortnagel K, Hofmann S, Gekeler F, Scharfe C, Rabl W, Gerbitz KD, Meitinger T (1998) Diabetes insipidus, diabetes
795 mellitus, optic atrophy and deafness (DIDMOAD) caused by mutations in a novel gene (wolframin) coding for a predicted
796 transmembrane protein. *Hum Mol Genet* 7, 2021-2028

797 Takeda K, Inoue H, Tanizawa Y, Matsuzaki Y, Oba J, Watanabe Y, Shinoda K, Oka Y (2001) WFS1 (Wolfram syndrome 1) gene
798 product: predominant subcellular localization to endoplasmic reticulum in cultured cells and neuronal expression in rat brain.
799 *Hum Mol Genet* 10, 477-484

800 Takei D, Ishihara H, Yamaguchi S, Yamada T, Tamura A, Katagiri H, Maruyama Y, Oka Y (2006) WFS1 protein modulates the free
801 Ca(2+) concentration in the endoplasmic reticulum. *FEBS Lett* 580, 5635-5640

802 Toots M, Seppa K, Jagomae T, Koppel T, Pallase M, Heinla I, Terasmaa A, Plaas M, Vasar E (2018) Preventive treatment with
803 liraglutide protects against development of glucose intolerance in a rat model of Wolfram syndrome. *Scientific Reports* 8

804 Ueda K, Kawano J, Takeda K, Yujiri T, Tanabe K, Anno T, Akiyama M, Nozaki J, Yoshinaga T, Koizumi A, Shinoda K, Oka Y, Tanizawa
805 Y (2005) Endoplasmic reticulum stress induces Wfs1 gene expression in pancreatic beta-cells via transcriptional activation.
806 *Eur J Endocrinol* 153, 167-176

807 van Hoek M, Dehghan A, Witteman JC, van Duijn CM, Uitterlinden AG, Oostra BA, Hofman A, Sijbrands EJ, Janssens AC (2008)
808 Predicting type 2 diabetes based on polymorphisms from genome-wide association studies: a population-based study.
809 *Diabetes* 57, 3122-3128

810 Visnapuu T, Plaas M, Reimets R, Raud S, Terasmaa A, Kõks S, Sütt S, Luuk H, Hundahl CA, Eskla K-L, Altpere A, Althoa A, Harro
811 J, Vasar E (2013a) Evidence for impaired function of dopaminergic system in Wfs1-deficient mice. *Behavioural Brain Research*
812 244, 90-99

813 Visnapuu T, Raud S, Loomets M, Reimets R, Sütt S, Luuk H, Plaas M, Koks S, Volke V, Althoa A, Harro J, Vasar E (2013b) Wfs1-
814 deficient mice display altered function of serotonergic system and increased behavioral response to antidepressants. *Front*
815 *Neurosci* 7, 132

816 Vora AJ, Lilleyman JS (1993) Wolfram syndrome: mitochondrial disorder. *Lancet* 342, 1059

817 Wang L, Liu H, Zhang X, Song E, Wang Y, Xu T, Li Z (2021) WFS1 functions in ER export of vesicular cargo proteins in pancreatic

818 beta-cells. Nat Commun 12, 6996

819 Waszczykowska A, Zmyslowska A, Braun M, Ivask M, Koks S, Jurowski P, Mlynarski W (2020) Multiple Retinal Anomalies in
820 Wfs1-Deficient Mice. Diagnostics (Basel) 10

821 Westermark P, Neuman RJ, Wasson J, Atzmon G, Wainstein J, Yerushalmi Y, Cohen J, Barzilai N, Blech I, Glaser B, Permutt MA
822 (2010) Gene-Gene Interactions Lead to Higher Risk for Development of Type 2 Diabetes in an Ashkenazi Jewish Population.
823 PLoS ONE 5

824 Wolfram DJ, Wagener HP (1938) DIABETES MELLITUS AND SIMPLE OPTIC ATROPHY AMONG SIBLINGS: REPORT OF FOUR CASES.
825 Mayo Clinic Proceedings 13, 715-718

826 Yamada T, Ishihara H, Tamura A, Takahashi R, Yamaguchi S, Takei D, Tokita A, Satake C, Tashiro F, Katagiri H, Aburatani H,
827 Miyazaki J, Oka Y (2006) WFS1-deficiency increases endoplasmic reticulum stress, impairs cell cycle progression and triggers
828 the apoptotic pathway specifically in pancreatic beta-cells. Hum Mol Genet 15, 1600-1609

829 Young TL, Ives E, Lynch E, Person R, Snook S, MacLaren L, Cater T, Griffin A, Fernandez B, Lee MK, King MC (2001) Non-
830 syndromic progressive hearing loss DFNA38 is caused by heterozygous missense mutation in the Wolfram syndrome gene
831 WFS1. Hum Mol Genet 10, 2509-2514

832 Zatyka M, Da Silva Xavier G, Bellomo EA, Leadbeater W, Astuti D, Smith J, Michelangeli F, Rutter GA, Barrett TG (2015)
833 Sarco(endo)plasmic reticulum ATPase is a molecular partner of Wolfram syndrome 1 protein, which negatively regulates its
834 expression. Hum Mol Genet 24, 814-827

835 Zatyka M, Ricketts C, da Silva Xavier G, Minton J, Fenton S, Hofmann-Thiel S, Rutter GA, Barrett TG (2007) Sodium-potassium
836 ATPase 1 subunit is a molecular partner of Wolframin, an endoplasmic reticulum protein involved in ER stress. Human
837 Molecular Genetics 17, 190-200

838 Zatyka M, Rosenstock TR, Sun C, Palhegyi AM, Hughes GW, Lara-Reyna S, Astuti D, di Maio A, Sciauvaud A, Korsgen ME,
839 Stanulovic V, Kocak G, Rak M, Pourtoy-Brasselet S, Winter K, Varga T, Jarrige M, Polveche H, Correia J, Frickel EM, Hoogenkamp
840 M, Ward DG, Aubry L, Barrett T, Sarkar S (2023) Depletion of WFS1 compromises mitochondrial function in hiPSC-derived
841 neuronal models of Wolfram syndrome. Stem Cell Reports 18, 1090-1106

842

843

844 **Figure legends**

845

846 **Fig. 1** The structure of *WFS1*. *WFS1* consists of eight exons, with the largest one being exon 8. The location of reported variants
847 is depicted in different colors. The image has been adapted from Clinvar (<https://www.ncbi.nlm.nih.gov/clinvar/>), and the
848 information was accessed on July 23, 2023

849

850 **Fig. 2** Multiple functions of WFS1. (A) WFS1 regulates Ca^{2+} uptake in the ER by modulating SERCA activity. WFS1
851 dysfunction increases ER Ca^{2+} levels and SOCE. Increased Ca^{2+} concentration activates Calpain-2, leading to cell death. (B)
852 WFS1 activates IP_3R through NCS1 and stimulates Ca^{2+} release from the ER. WFS1 dysfunction impairs IP_3R activity, followed
853 by decreased Ca^{2+} release from the ER and Ca^{2+} uptake into the mitochondria. (C) WFS1 stabilizes HRD1, an E3 ubiquitin

854 ligase, and degrades ATF6. WFS1 abnormality induces the hyperactivation of ATF6. (D) WFS1 plays a vital role in maintaining
855 the pH within secretory granules. WFS1 dysfunction leads to impaired granule acidification and insulin exocytosis. (E) The C-
856 terminal of WFS1 binds to vesicular cargo proteins. The N-terminal of WFS1 is recognized by the protein transport protein
857 SEC24, a component of coat protein complex II (COPII). WFS1 dysfunction disrupts the generation of mature COPII vesicles
858 and hinders intercellular trafficking from the endoplasmic reticulum (ER) to the Golgi complex. ATF6, activating transcription
859 factor 6; HRD1, HMG-CoA reductase degradation 1 homolog; SERCA, sarcoendoplasmic reticulum ATPase; SOCE, store-
860 operated Ca²⁺ entry; IP₃R, 1,4,5-trisphosphate receptor; NCS1, neuronal calcium sensor 1; GRP75, glucose-regulated protein
861 75; VDAC1, voltage dependent anion channel 1; MCU, mitochondrial Ca²⁺ uniporter. Created with BioRender.com

862
863

864 Table Animal models of Wolfram syndrome
865

Reference	Species (Strain)	Specificity	Targeted region of <i>Wfs1</i>	Phenotype / onset or tested age
Ishihara et al. (2004) (Ishihara et al. 2004)	Mouse [(129Sv x B6) x B6] F2	Whole body <i>Wfs1</i> KO	Exon 2 (Replaced by neomycin- resistance gene)	Non-fasted hyperglycemia /16 weeks (Ishihara et al. 2004)
				Decreased whole-pancreas insulin content / 2 weeks old (Ishihara et al. 2004)
				Impaired glucose homeostasis and insulin secretion on OGTT / 17 weeks old (Ishihara et al. 2004)
	Mouse (B6)			Decreased AVP secretion (Kurimoto et al. 2021)
				Alternation in emotionally triggered behavior, social interaction, and behavioral despair / 34 weeks old (Kato et al. 2008)
				Impaired photoreceptor and inner

				retinal functions. Elevated ER stress in the retina (Bonnet Wersinger et al. 2014)
Riggs et al. (2005) (Riggs et al. 2005)	Mouse (B6)	Conditional <i>Wfs1</i> KO (Pancreatic β -cell)	Exon 8	Glucose intolerance and insulin deficiency / 12 weeks old
				Impaired glucose homeostasis / 6 weeks old (Abreu et al. 2020)
				Lower ratio of retinal thickness/longitudinal diameter (Waszczykowska et al. 2020)
				Impaired behavioral adaptation / 2-4 months old (Luuk et al. 2009)
Luuk et al. (2008) (Luuk et al. 2008)	Mouse (B6)	Whole body <i>Wfs1</i> KO	Exon 8 (Replaced by <i>LacZ</i>)	Hearing loss / 4 months old (Richard et al. 2023)
Köks et al. (2009) (Köks et al. 2009)	(129S6)			Impaired fertility (Noormets et al. 2009)
				Reduced volume of the optic nerve, brain stem, and cortex (Blackstone et al. 2016)
				Dopaminergic system impairment (Visnapuu et al. 2013a)
				Increased sensitivity to the serotonin inhibitor (Visnapuu et al. 2013b)
				Lower body mass, food intake, and serum leptin levels than wild type

(Noormets et al. 2014)

Impaired insulin secretion In high-fat diet-fed *Wfs1* heterozygous mice
(Ivask et al. 2021)

Richard et al. (2023)	Mouse	Whole body <i>Wfs1</i>	Exon 8	Hearing loss / post-natal day 27-29
(Richard et al. 2023)	(B6)	E864K KI	(E864K KI)	Vestibular deficits / post-natal day 21 (Richard et al. 2023)
Plaas et al. (2017)	Rat	Whole body <i>Wfs1</i>	Exon 5	Impaired glucose-stimulated insulin secretion / 3 months old (Plaas et
(Plaas et al.	(Sprague-Dawley	KO	(Deleted 27 amino acids (aa 212–238) and a	al. 2017)

2017)	rats)		substitution of serine to alanine at aa 239)	Hyperglycemia, severe body weight loss / by 12 months old (Plaas et al. 2017)
				Retinal gliosis, optic nerve atrophy / 15 months old (Plaas et al. 2017)
				Decreased medullary volume (Plaas et al. 2017)
				Low-frequency, sensorineural hearing loss / 6.5 months old (Jagomäe et al. 2021)
Sakakibara et al. (2018) (Sakakibara et al. 2018)	Drosophila	Conditional <i>wfsI</i> KO (neurons, glial cells)	<i>Drosophila wfsI</i> (CG4917, <i>WFSI</i> homolog)	Behavioral deficits, age-associated neurodegeneration (Sakakibara et al. 2018)
			<i>Drosophila wfsI</i> (CG4917, <i>WFSI</i> homolog)	Flight less (<i>wfsI^{e03461}</i> / <i>wfsI^{e03461}</i>) / 5 days old (Kunnappallil and Hasan 2022)
Kunnappallil and Hasan (2022) (Kunnappallil and Hasan 2022)	Drosophila	Whole body <i>wfsI</i> KI	Homozygous null variant (<i>wfsI^{e03461}</i> / <i>wfsI^{e03461}</i>)	Flight defect (<i>wfsI^{M114041}</i>) / 20 days old (Kunnappallil and Hasan 2022)
		Conditional (neurons) <i>wfsI</i> KD	C-terminal truncated variant (<i>wfsI^{M114041}</i>)	Flight defect (pan- or dopaminergic-neuronal <i>wfsI</i> KD) / 5 days old (Kunnappallil and Hasan 2022)
			Pan- or dopaminergic-neuronal <i>wfsI</i> KD	
Cairns et al. (2021) (Cairns et al. 2021)	Zebrafish	Whole body <i>wfsIa^{W692X}</i> (sa10021) or <i>wfsIb^{W493X}</i> (sa16422)	<i>wfsIa</i> or <i>wfsIb</i> (orthologues in zebrafish)	Neurodegeneration, loss of retinal ganglion cells, and visual dysfunction in zebrafish carrying <i>wfsIb^{W493X}</i> (Cairns et al. 2021)

Crouzier et al. (2022)	Zebrafish	Whole body		Enlarged ear area and decreased rod cells in retina (<i>wfs1a</i> ^{C825X})
		<i>wfs1a</i> ^{C825X} (sa11465)	<i>wfs1a</i> or <i>wfs1b</i>	Mitochondrial dysfunction (<i>wfs1b</i> ^{W493X})
(Crouzier et al. 2022b)		or	(orthologues in zebrafish)	Decreased expression of UPR-related genes in physiological conditions and impaired visual function (<i>wfs1a</i> ^{C825X} , <i>wfs1b</i> ^{W493X})
		<i>wfs1b</i> ^{W493X} (sa16422)		(Crouzier et al. 2022b)

866 B6, C57 BL/6; KO, knock out; KI, knock-in; OGTT, oral glucose tolerance test; KD, knockdown; UPR, unfolded protein
867 response.