



Title	Miscellaneous notes on Braconidae (Hymenoptera)
Author(s)	Watanabe, Chihisa
Citation	Insecta matsumurana, 18(3-4), 66-66
Issue Date	1954-09
Doc URL	https://hdl.handle.net/2115/9541
Type	departmental bulletin paper
File Information	18(3-4)_p66-66.pdf



MISCELLANEOUS NOTES ON BRACONIDAE

(Hymenoptera)

By CHIHISA WATANABE

Entomological Institute, Hokkaido University, Sapporo

Macrocentrus gifuensis ASHMEAD, 1906

This species has been widely known as a parasite of the European corn borer, *Pyrausta nubilalis* HÜBNER. The present specimens, however, were reared by KAWASE from the bean webworm, *Sylepta ruralis* SCOPOLI, at Takada. According to SUGIYAMA and KAWASE (1952)¹⁾ this species is bred from larvae of *Sylepta ruralis*, feeding on leaves of *Boehmeria tricuspis* MAKINO.²⁾

Hosts: *Pyrausta rubilalis* HÜBNER and *Sylepta ruralis* SCOPOLI.

Specimens examined: 12 ♂♂ & 15 ♀♀ 20-vi-52, Takada, Niigata-ken, Japan, H. KAWASE leg.

Distribution: Japan, Korea, Manchuria and Europe.

Tropobracon CAMERON, 1905

On account of FAHRINGER's paper arrived belatedly in Japan, *Shirakia* VIERECK (1913) was sunken as a new synonym of *Tropobracon* CAMERON by the writer (1951)³⁾, but this synonymy had been already established by FAHRINGER (1942)⁴⁾.

1) Oyō-Koutyū, 8: 73, 1952.

2) A species of the family Urticaceae. The Japanese name is "Aka-so".

3) Ins. Mats., 17: 100, 1951.

4) Arb. morphol. taxon. Ent. Berlin-Dahlem, 9: 43, 1942.