



HOKKAIDO UNIVERSITY

Title	Discovery of <i>Oxydema longulum</i> Boheman in Saipan Island(Col., Curculionidae)
Author(s)	Konishi, Masayasu
Citation	Insecta matsumurana, 20(3-4), 104-104
Issue Date	1956-12
Doc URL	https://hdl.handle.net/2115/9595
Type	departmental bulletin paper
File Information	20(3-4)_p104-104.pdf



DISCOVERY OF *OXYDEMA LONGULUM* BOHEMAN
IN SAIPAN ISLAND

(Col., Curculionidae)

By MASAYASU KONISHI
Sapporo, Japan

Through the kindness of Mr. NODOKA HAYASHI in Yokohama, I have had the opportunity to add the following Cossonine weevil to the fauna of Saipan Island.

Oxydema longulum (BOHEMAN)

Rhyncolus longulus BOHEMAN, *Eugenies Resa. Ent.*, p. 149 (1859).

Pseudolus longulus SHARP, *Trans. Roy. Dublin Soc.*, II (3), p. 190, pl. V, Fig. 33 (1885); PERKINS, *Fauna Hawaiiensis*, II (3), p. 149 (1900); MARSHALL, *Insects of Samoa*, IV (5), p. 251, 332 (1931); CSIKI, in JUNK et SCHENKLING, *Col. Cat.*, 149, p. 191 (1936).

Oxydema longulum ZIMMERMAN, B. P. Bishop Mus., *Occ. Papers*, 15 (25), p. 287 (1940), B. P. Bishop Mus., *Bull.* 172, p. 138, pl. 4, Fig. C (1942); SWEZEY, B. P. Bishop Mus., *Spec. Publ.* 44, p. 52, 147 (1954).

Specimens examined: 2 ♂♂, Tarahoho, Saipan Island, 15. X, 1941, DENGO MATSUSHITA leg.

This species has hitherto been recored from the Hawaiian Islands, the Marquesas Island, Pitcairn Island, the Samoan Islands, Tonga Island and Guam Island.