



HOKKAIDO UNIVERSITY

Title	Detection and characterization of vector-borne parasites and Wolbachia endosymbionts in greater one-horned rhinoceros (<i>Rhinoceros unicornis</i>) in Nepal
Author(s)	Pandey, Gita Sadaula; Manandhar, Prajwol; Shrestha, Bijaya Kumar et al.
Citation	Acta tropica, 258 https://doi.org/10.1016/j.actatropica.2024.107344
Issue Date	2024-10
Doc URL	https://hdl.handle.net/2115/96162
Type	journal article
File Information	111355.pdf



Detection and characterization of vector-borne parasites and *Wolbachia* endosymbionts in greater one-horned rhinoceros (*Rhinoceros unicornis*) in Nepal

Gita Pandey Sadaula^{1,2}, Prajwol Manandhar³, Bijaya Kumar Shrestha⁴, Amir Sadaula⁵, Naoki Hayashi^{1,6}, Abdelbaset Eweda Abdelbaset^{1,7}, Pradeepa Silwal⁵, Toshio Tsubota⁸, Mackenzie L. Kwak¹, Nariaki Nonaka^{1,6}, Ryo Nakao^{1,6*}

¹Laboratory of Parasitology, Graduate School of Infectious Diseases, Faculty of Veterinary Medicine, Hokkaido University, Japan

² National Cattle Research Program, Nepal Agricultural Research Council, Rampur, Chitwan, Nepal

³Center for Molecular Dynamics Nepal, Thapathali, Kathmandu, Nepal

⁴Chitwan National Park Office, Kasara, Chitwan, Nepal

⁵National Trust for Nature Conservation – Biodiversity Conservation Center, Sauraha, Chitwan, Nepal

⁶Division of Parasitology, Veterinary Research Unit, International Institute for Zoonosis Control, Hokkaido University, Japan

⁷Clinical Laboratory Diagnosis, Department of Animal Medicine, Faculty of Veterinary Medicine, Assiut University, Assiut, 71515, Egypt

⁸Laboratory of Wildlife Biology and Medicine, Faculty of Veterinary Medicine, Hokkaido University, Japan

Abstract:

Vector-borne parasite infections affect both domestic and wild animals. They are often asymptomatic but can result in fatal outcomes under natural and human-induced stressors. Given the limited availability of molecular data on vector-borne parasites in *Rhinoceros unicornis* (greater one-horned rhinoceros), this study employed molecular tools to detect and characterize the vector-borne parasites in rescued rhinoceros in Chitwan National Park, Nepal. Whole blood samples were collected from thirty-six *R. unicornis* during rescue and treatment operations. Piroplasmida infections were first screened using nested polymerase chain reaction (PCR) targeting 18S ribosomal RNA gene. *Wolbachia* was detected by amplifying 16S rRNA gene, while filarial nematodes were detected through amplification of 28S rRNA, COI, *myoHC* and *hsp70* genes. Our results confirmed the presence of *Theileria bicornis* with a prevalence of 75% (27/36) having two previously unreported haplotypes (H8 and H9). *Wolbachia* endosymbionts were detected in 25% (9/36) of tested samples and belonged to either supergroup C or F. Filarial nematodes of

the genera *Mansonella* and *Onchocerca* were also detected. There were no significant association between *T. bicornis* infections and the age, sex, or location from which the animals were rescued. The high prevalence of *Theileria* with novel haplotypes along with filarial parasites has important ecological and conservational implications and highlights the need to implement parasite surveillance programs for wildlife in Nepal. Further studies monitoring vector-borne pathogens and interspecies transmission among wild animals, livestock and human are required.

Keywords: *Mansonella*, Nepal, *Onchocerca*, *Rhinoceros unicornis*, *Theileria*, *Wolbachia*

1 Introduction

The greater one-horned rhinoceros (*Rhinoceros unicornis*) (hereinafter rhinoceros) is the second largest extant megaherbivore in South Asia, specifically inhabiting the floodplains of the Ganges, Brahmaputra, and Sindh rivers and their large tributaries, stretching from the Indo-Burmese border in the east to the Pakistan border in the west (Dinerstein, 2003). Rhinoceros are categorized as vulnerable by IUCN red list and are currently restricted to small, isolated protected areas of less than 20,000 km² due to habitat loss, fragmentation, and poaching (Ellis and Talukdar, 2019; Jhala et al., 2021). The major population of rhinoceros in Nepal resides in Chitwan National Park (CNP) (694 out of 752), according to a recent census carried out in 2021 with an annual growth rate of less than 2.5%, which is half of that in the previous census carried out in 2015 (~ 5%). Currently, the population of rhinoceros in CNP has been rejuvenated from < 100 individuals during the 1960s with continuous conservation efforts including the establishment of national park in 1973 and strict law enforcement (Dinerstein, 2003). Since 2015, the mortality of rhinoceros has spiked with an average of ~ 25 deaths annually (Bhandari et al., 2022). Although incidences of poaching have considerably declined due to rigorous conservation efforts, a rise in mortalities due to unknown or natural causes has reduced the annual growth rate of

the population. Conservationists are increasingly concerned about the potential impact of pathogen on the health and mortality of Nepal's rhinoceros populations.

Piroplasmida infections have been associated with morbidity and mortality in rhinoceros population in various countries. *T. bicornis* was first described from the black rhinoceros in Africa (Nijhof et al., 2003). It was subsequently found in white rhinoceroses in Kruger National Park (Govender et al., 2011) and in black and white rhinos in Kenya (Otiende et al., 2016). Besides free-ranging rhinoceros, *T. bicornis* was also detected in captive rhinoceros in Australia (Yam et al., 2018). Additionally, *T. bicornis* was reported in other wildlife in South Africa (Pfitzer et al., 2011) and in cattle in Uganda (Muhanguzi et al., 2010). Three haplotypes (H1, H3 and H4) of *T. bicornis* reported from black rhinoceros and haplotype H2 reported from white rhinoceros (Otiende et al., 2016; Yam et al., 2018). Filarioids are parasitic nematodes belonging to the superfamily Filarioidea and can infect a wide range of hosts, including humans. These parasites can be transmitted by an extensive range of arthropod vectors including mosquitoes and ticks (Otranto et al., 2011; Otranto and Deplazes, 2019; Perles et al., 2023; PupiĆ-Bakrač et al., 2021; Shelley and Coscarón, 2001). Epidemiological studies of filarioids have primarily focused on domestic animals (Grácio et al., 2015; Otranto and Dantas-Torres, 2010). *Dirofilaria immitis* has been reported in wild carnivores in different regions of the world (Otranto et al., 2015), while *Mansonella* sp. has been recorded in ring-tailed coatis (Perles et al., 2023), deer (Uni et al., 2004), and bears (Masatani et al., 2021). However, the epidemiology of filarioids in megaherbivores is still largely unknown. Interestingly, some filarial parasites harbor *Wolbachia* as an endosymbiont, some of which are involved in the reproduction, development, and survival of the host nematode (Martin and Gavotte, 2010; Taylor et al., 2005).

In Nepal, several studies have examined the vector-borne parasitic infections in livestock and ticks. Of note, *Theileria* infection was commonly observed in cattle (Shrestha et al., 2017; Poudel et al., 2023) and ticks (Gupta et al., 2013). Until now, however, no information is available on *Theileria* infection in wild animals in Nepal. Additionally, there is no data available on filarial infections in livestock and wildlife in Nepal. In this study, molecular techniques were employed to identify and characterize the vector-borne parasites in free-ranging rhinoceros population in Nepal.

2 Materials and Methods

2.1 Study area and sample collection:

CNP lies in the south-central part of Nepal with an area of 952.63 km² and a buffer zone area of 729.37 km². The landscape of CNP is a matrix of grassy flood plains, and tropical evergreen Sal Forest while the buffer zone is highly settled and heavily cultivated. The climate in CNP is subtropical monsoon with three distinct seasons: monsoon (June – October), cool-dry (October – February), and hot-dry (February – June). Mean annual temperature ranges between 8°C in January and 36°C in April and rainfall of 2,036 ± SE 64 mm per year (>80% in monsoon) (Subedi, 2012). A total of 68 mammal species, 544 bird species, 56 reptile and amphibian species, and 126 species of fish are recorded in the park (CNP, 2012).

Samples from 36 rhinoceros (19 from the core area and 17 from the buffer zone) were collected opportunistically during rescue operations between 2015 and 2022. Rhinoceros were rescued when they entered human settlements and were returned to their natural habitat. Some were injured in territorial fights with conspecifics, while the floods forcibly expelled others from their original habitat. Rhinoceros were immobilized jointly by the Department of National Park and Wildlife Conservation and the National Trust for Nature

Conservation (NTNC) using a standard protocol with a combination of etorphine hydrochloride and acepromazine delivered by dart gun and reversed by diprenorphine. The blood samples were collected from an auricular vein and stored at -20°C in the NTNC– Biodiversity Conservation Center (NTNC-BCC), molecular laboratory, Sauraha, Chitwan. The chief of the CNP office granted legal permission for rhinoceros capture and handling. Ethical approval for the study was obtained from the Ethical Clearance Committee of Nepal Veterinary Council (Ref No: 56/2079.80).

2.2 DNA extraction and PCR

Extraction of genomic DNA from whole blood was carried out using a DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) following the manufacturer’s protocol. A nested polymerase chain reaction (PCR) for *Babesia*, *Theileria*, and *Hepatozoon* species (BTH) was performed to amplify almost a section of the 18S ribosomal RNA gene (rDNA) (1400-1600 bp) (Masatani et al., 2017). The primer sets of BTH 1st F/BTH 1st R and BTH 2nd F/BTH 2nd R were used for the first and second PCRs, respectively.

The presence of Anaplasmataceae was first screened with the primer set (EHR16SD and EHR16SR) amplifying 16S ribosomal RNA gene (rDNA) (Parola et al., 2000). The positive samples were further characterized by running additional PCRs amplifying heat shock protein (*groEL*) and citrate synthase (*gltA*) genes of *Anaplasma* and *Ehrlichia* and 16S rDNA of *Wolbachia* using the primers listed in Table 1.

Samples positive for *Wolbachia* were subjected to PCR to amplify 28S rDNA and cytochrome c oxidase subunit I (COI) gene of *Onchocerca* and heat shock protein 70 (*hsp70*) and myosin heavy chain (*myoHC*) genes of *Mansonella*.

2.3 In-Fusion cloning

Samples showing mixed signals in the sequencing analysis of *Wolbachia* 16S rDNA PCR products were subjected to In-Fusion cloning. First, PCR primers (InFusion_EHR16SD and InFusion_1513R) were designed based on the alignment of *Wolbachia* 16S rDNA sequences available in the database. The purified PCR products amplified by the newly designed primers were cloned into pUC19 vector (TaKaRa Bio Inc, Japan). The infusion reaction was performed in a total volume of 10 μ L, containing 2.0 μ L of 5X In-Fusion HD Enzyme Premix, 1 μ L of PCR-linearized pUC19 vector, 200 ng of each purified PCR fragment, and dH₂O from the In-Fusion HD PCR Cloning Kit (TaKaRa Bio Inc, Japan). The infusion reaction mix was incubated at 50°C for 15 min, and then placed on ice. Following incubation, the heat-shock transformation was conducted by adding 2.5 μ L of the infusion reaction mixture to the competent cells, *Escherichia coli* DH5 α (Nippon Gene, Tokyo, Japan). Transformed cells were plated on LB agar plates containing ampicillin and incubated at 37°C for 24 h. The transformants were screened with colony PCR.

2.4 Sanger sequencing

The PCR amplicons were purified using ExoSAP-IT™ Express PCR Product Cleanup Reagent (Thermo Fisher Scientific, Waltham, MA, USA). The sequencing reaction was conducted using the BigDye Terminator version 3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA) on an ABI Prism 3130xl Genetic Analyzer (Applied Biosystems). Sequence data editing and trimming of primer annealing sites was done using ATGC software (GENETYX Corporation, Tokyo, JP) and the consensus sequences were extracted for phylogenetic analysis. The sequences generated in this study were submitted to the DNA Data Bank of Japan (DDBJ) under the accession numbers: LC817751 to LC817777 for 18S rDNA of *Theileria bicornis*, LC817778 to LC817789 for

16S rDNA of *Wolbachia*, LC817841 to LC817843 for 28S rDNA of nematodes, LC817844 to LC817849 for COI of *Onchocerca* sp., LC817850 to LC817856 for *myoHC* gene of *Mansonella* sp., and LC817857 to LC817863 for *hsp70* gene of *Mansonella* sp.

2.5 Phylogenetic Analysis

We aligned the obtained sequences with those available in the database using MAFFT v7.450 software (Kato and Standley, 2013). After visual inspection, we performed the necessary trimming and editing of the sequences using MEGA7 (Kumar et al., 2016). Completely identical sequences were detected and brought together using the “pgelimdupseq” command in Phylogears2 ver. 2.0 (Tanabe, 2008), followed by the best-fit model estimation based on the Bayesian information criterion by Kakusan4 (Tanabe, 2011), choosing the K80 with gamma distribution for 16S rDNA of *Wolbachia*, the Hasegawa Kishino Yano (HKY) model with gamma distribution for 18S rDNA, and the general time-reversible (GTR) model with gamma distribution for 28S rDNA. Codon-proportional models were selected for *hsp70*, *myoHC*, and COI genes.

The phylogenetic analysis was conducted using MrBayes v3.2.6 (Ronquist et al., 2012) with 1,000,000-5,000,000 generations at a sample frequency of 500 discarding the first 10% of trees as a burn-in. The consensus tree was visualized using FigTree v1.4.4 (Rambaut, 2012).

To visualize the relationships between the *T. bicornis* haplotypes based on the 18S rDNA sequences, a median-joining haplotype network analysis was performed using PopART v1.7 (Leigh and Bryant, 2015). To compare our data with those previously reported from other regions, we added haplotypes from Australia (H2 and H4), Kenya (H1, H2 and H3), South Africa (H1 and H3).

2.6 Statistical Analysis

The association between Piroplasmida infection and age, sex, and habitat was assessed using GraphPad Prism 5.0 software (GraphPad Software, Inc., La Jolla, CA, USA). Fisher's exact test was utilized for this analysis. Odds ratios (OR) along with their corresponding 95% confidence intervals (95% CI) were computed. A significance level of $P < 0.05$ was used to determine statistical significance.

3 Results

3.1 Detection and characterization of *Theileria bicornis*

A total of 27 out of 36 (75%) rhinoceros were positive for *Theileria* infection by BTH PCR. All PCR amplicons were successfully sequenced, and two sequence types of *T. bicornis* were identified.

Out of the 27 sequences we obtained, 25 were completely identical to each other (LC817751-LC817757, LC817759-LC817771 and LC817773-LC817777) and another two were identical to each other (LC817758 and LC817772). The sequences obtained from this study were distinct from those of previously published haplotypes (H1 to H4) from African rhinoceros species and (H2 to H7) reported from *Amblyomma thollonii*. We have designated the two haplotypes of greater one-horned rhinoceroses as H8 and H9 (Figure 1). The geographic distribution of each haplotype is shown in Figure 2. The H8 haplotype was evenly distributed throughout CNP, while the H9 haplotype occurred only in one specific location.

The prevalence of infection in relation to sex, age and location is shown in Table 2. We found that the positive rate increased with age, but there was no statistically significant association observed. There was no association between *Theileria* infection and sex or sampling location.

3.2 Detection and characterization of Anaplasmataceae

A total of 9 out of 36 (25%) rhinoceros were positive for Anaplasmataceae by PCR. None of the samples were positive for *groEL* and *gltA* PCRs amplifying the genera *Anaplasma* and *Ehrlichia*. Instead, all nine samples were positive for 16S rDNA PCR for *Wolbachia*. As the direct sequencing analysis of the PCR amplicons showed mixed signals (presumably due to mixed infections), the samples were subjected to In-Fusion cloning to separate the sequences. Two distinct sequence types of *Wolbachia* were identified in two samples. Phylogenetic analysis suggested that these two *Wolbachia* sequences belonged to supergroups C and F. Specifically, one sequence type (LC817778-LC817783) is closely associated with the *Wolbachia* endosymbiont of the genus *Onchocerca* belonging to supergroup C, whereas the other (LC817784-LC817789) is closely linked to the *Wolbachia* endosymbiont of the genus *Mansonella* belonging to supergroup F. The blast result showed that one sequence type has 99% nucleotide identity with the *Wolbachia* endosymbiont of *Onchocera eberhardi* (KU255248) and the other shows 99% nucleotide identity with the *Wolbachia* endosymbiont of *Mansonella ozzardi* (AJ279034).

3.3 Detection and characterization of nematodes

All samples positive for *Wolbachia* by PCR were also positive by PCR for the 28S rDNA locus of nematodes. Sequence analysis of the amplified products identified two different sequence types. The phylogenetic tree revealed that the obtained sequences were clustered together either with the genera *Onchocerca* and *Mansonella* (Figure 4). To characterize *Onchocera* sp., the mitochondrial COI gene was amplified, and the sequence was compared with those in the database. The phylogenetic analysis showed that the sequence obtained in this study is not clustered with validated species of the genus *Onchocerca* (Figure 5). Likewise, the phylogenetic trees constructed for *hsp70* and *myoHC* genes of

Mansonella revealed that the sequences obtained from rhinoceros belong to a discrete taxon within the genus *Mansonella* clade.

4 Discussion

Considering the limited knowledge regarding vector-borne infections in the vulnerable populations of greater one-horned rhinoceroses in Nepal, we first aimed to screen for a panel of common tick-borne pathogens including Piroplasmida and Anaplasmatidae. In addition to the first detection of *T. bicornis* in Asian rhinoceroses, our efforts led to the discovery of a poorly characterized groups of filarial parasites through their association with *Wolbachia* endosymbionts.

Our study detected *T. bicornis* in greater one-horned rhinoceros with an prevalence of 75%, which is higher than those reported from Kenya with infection rate of 43% and 66% in black and white rhinoceroses, respectively (Otiende et al., 2015) and from South Africa with a prevalence rate of 23.7% and 36.4% in black and white rhinoceros, respectively (Govender et al., 2011; Zimmermann et al., 2021). Additionally, lower prevalence rate of *T. bicornis* were reported from Nyala antelopes (52.6%) in South Africa (Pfitzer et al., 2011) and from cattle (3.8%) originating from western Uganda (Muhanguzi et al., 2010). These variations in infection rate of *T. bicornis* between different studies may be attributed to species differences, geographic locations, and availability of tick vectors.

The rhinoceros included in this study were rescued from both the core area and buffer zones. Moving rhinoceroses from the buffer zone back into the core area may increase the risk of introducing new pathogens to the potentially naïve rhinoceros population in the core area. These rescued rhinoceroses often came into contact with livestock in the buffer zone, which can serve as hosts for ticks and haemoparasites. This practice raises the likelihood of spillover of *T. bicornis* from alternative hosts to naïve rhinoceros population.

The precise transmission dynamics and specific vectors involved in the spread of *T. bicornis* remain unknown. *Theileria bicornis* (H2-H7) has been identified in *Amblyomma tholloni* ticks collected from African elephants in Kenya (King'ori et al., 2019), suggesting that *Amblyomma* spp., which are widely distributed in Nepal, could act as vectors of *T. bicornis* (Bohara and Shrestha, 2015). Consequently, molecular investigations are needed to identify *T. bicornis* or its tick vectors in livestock populations within Nepal.

Subadult rhinoceros were more likely to experience stress-related changes, including reproductive maturity, courtship, mating, and territorial competition, which could increase their vulnerability to *Theileria* infections (Govender et al., 2011). Nevertheless, our study showed no significant relationship between *T. bicornis* infection and age, which is consistent with the findings from a previous study by Otiende et al. (2015). However, our results may be influenced by sampling bias because a substantial proportion of the rhinoceros enrolled in our study were adults, which may have masked the relationship between infection status and age. Our investigation revealed no significant association between *T. bicornis* infection and sex, suggesting that males and females were at approximately equal risk of acquiring the infection. This finding may be attributed to the animals sharing the same habitat and being exposed to the same vector communities, as well as experiencing similar stress factors in the study area. These results are in line with previous studies conducted by Govender et al. (2011) and Otiende et al., (2015). Other investigators, however, showed that parasitism is sometimes sex-biased with males being infected more often with TBPs than females, at least in dogs (Chatanga et al., 2021).

Globally there are five different rhinoceros species: Africa is home to two species, black and white rhinoceros, while three species, greater one-horned, Sumatran and Javan rhinoceros, are native to Asia. The network analysis revealed that the population of *T.*

bicornis in the greater one-horned rhinoceroses in Nepal had two distinct haplotypes from those reported from African rhinoceroses (Yam et al., 2018; Zimmermann et al., 2021). This fact suggests that *T. bicornis* may have co-evolved with their animal hosts, leading to an equilibrium state in which infected rhinoceroses do not develop disease from *T. bicornis* infections (Otiende et al., 2015). However, stressors that disrupt host-parasite equilibrium can result in clinical infections. For example, certain species of *Theileria*, such as *Theileria velifera* in cattle and the *Theileria orientalis* genotype Buffeli (Kamau et al., 2011; Mans et al., 2015) can establish lifelong infections within their hosts. These infections typically do not result in clinical symptoms unless the animals experience immunosuppression or undergo stressful conditions such as pregnancy, translocation, or suboptimal rearing conditions (Sugimoto and Fujisaki, 2002). While our findings suggest that haplotypes H8 and H9 are native to South Asia and are presumably avirulent in greater one-horned rhinoceroses, we cannot exclude the possibility that these haplotypes are virulent to other rhinoceros species. Asymptomatic infections with tick-borne pathogens are common in wildlife but clinical manifestations sometimes only appear when these hosts are exposed to stressors such as co-infections with other pathogens (Munson et al., 2008), translocations (Höfle et al., 2004; Nijhof et al., 2005) malnutrition, and extreme environmental conditions (Nijhof et al., 2003).

We identified, for the first time, *Mansonella* sp. and *Onchocerca* sp. in rhinoceros in Nepal. Further species identification was impossible due to lack of parasite specimens for morphological examinations. However, this is not the first report on the detection of *Onchocerca* in blood samples. The detection of *Onchocerca* sp. in blood was previously reported in humans and donkeys in Venezuela and Italy, respectively (Botto et al., 1984; Papini et al., 2020). Clinical manifestation of filarial parasites have been previously reported in other species of rhinoceros in Africa, including hemorrhagic ulcerative filarial

301 lesions in black rhinoceros (Schulz and Kluge, 1960). Similarly, Mutinda et al., (2012)
302 reported the first case of putative filariasis outbreak in white and black rhino in Kenya
303 which may have been a result of an immunocompromised condition caused by food
304 shortage. More recently, King'ori et al., (2024) reported that ulcerative filarial skin
305 lesions caused by *Stephanofilaria dinniki* are a common disease in black and white
306 rhinoceroses. Considering the detrimental impact of filarial infections on the black and
307 white rhinoceroses during immunocompromised conditions, the clinical significance of
308 filarial parasites in greater one-horned rhinoceroses in Nepal cannot be rule out.

309 Filarial parasites including *Mansonella* and *Onchocerca* may also pose a zoonotic risk
310 (Cambra-Pellejà et al., 2020; Mediannikov and Ranque, 2018). Given that several species
311 of these parasites have been identified in wild animals (Bain et al., 1993; Masatani et al.,
312 2021; Perles et al., 2023; Uni et al., 2015, 2007; Yagi et al., 1994), the close contact
313 between wild animals, livestock and humans may also favor the potential transmission of
314 filarial parasites from animals to humans. Therefore, precise species identification of
315 filarial parasites is crucial for evaluating their zoonotic potential. Characterizing the
316 lifecycle of these parasites, including their definitive hosts, vectors, and developmental
317 stages, provides invaluable insights into their transmission and host-switching potential.
318 Such knowledge is essential for developing effective control measures to mitigate the risk
319 of zoonotic filarial diseases.

320 *Wolbachia* species are bacterial endosymbionts vertically transmitted in arthropods and
321 nematodes (Correa and Ballard, 2016; Manoj et al., 2021). They can be classified into
322 different supergroups in arthropods (A, B, E, H, I, K, and S), and nematodes (C, D, and
323 J), with supergroup F being common to both arthropods and filarial nematodes (Kaur et
324 al., 2021). In our study, *Wolbachia* detected in rhinoceroses fell into two distinct
325 supergroups: C and F. Similar findings were reported by Perles et al., (2023), where

Mansonella-associated *Wolbachia* in the supergroup F were detected in a blood sample from a ring-tailed coati (*Nasua nasua*) in Brazil. The detection of *Wolbachia* of different supergroups in rhinoceros samples can be attributed to multiple acquisition events by different major lineages of filarial nematodes (Manoj et al., 2021) . *Wolbachia* can serve as a diagnostic marker to detect the presence of filarial infections as suggested previously (Akter et al., 2019; Qiu et al., 2016). Additionally, a deeper understanding of the symbiotic relationship between *Wolbachia* and their filarial hosts may provide valuable insights into the biology, and evolution, and potential therapeutic targets for treatment of these filarial infections (Turb et al., 2012).

5 Conclusion:

This is the first report detecting *T. bicornis* with two new haplotypes in the Asian rhinoceroses. High infection rates suggested that this parasite is well established in the rhinoceros population in Nepal; however, its clinical impact, especially on those individuals under the exposure to the ecological stressors, remains unclear. The current investigation revealed that rhinoceroses in Nepal are infected with two different filarial nematodes through their association with *Wolbachia* endosymbionts. Further studies are needed to determine the taxonomical position and life cycle of these filarial parasites together with their zoonotic potential. Given the frequent rescue of rhinoceroses from human settlements and their cohabitation with livestock, the disease spillover between these two host species is likely to occur. Therefore, understanding the epidemiology of vector-borne organisms necessitates a thorough investigation of the interactions between multiple hosts that share habitat.

CRediT authorship contribution statement

Gita Pandey Sadaula: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing. **Prajwol Manandhar:** Investigation, Writing – review & editing, **Bijaya Kumar Shrestha:** Resources, **Amir Sadaula:** Resources, Writing – review & editing, **Naoki Hayashi:** Formal analysis, Software, Writing – review & editing **Abdelbaset Eweda Abdelbaset:** Formal analysis, Writing – review & editing, **Pradeepa Silwal:** Investigation, **Toshio Tsubota:** Writing – review & editing, **Mackenzie L. Kwak:** Writing – review & editing. **Nariaki Nonaka:** Supervision, Writing – review & editing. **Ryo Nakao:** Conceptualization, Funding acquisition, Methodology, Supervision, Validation, Writing – review & editing

Declaration of competing Interest

The authors declare no competing interest.

Acknowledgments

The author thanks all the staff of the Chitwan National Park and National Trust for Nature Conservation for their assistance with sampling.

This study was supported in part by the Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan (JP20KK0151 and JP23K23770); the Japan Agency for Medical Research and Development (AMED), Japan (JP23wm0225034).

References:

Akter, S., Nakao, R., Imasato, Y., Alam, M.Z., Katakura, K., 2019. Potential of cell-free DNA as a screening marker for parasite infections in dog. *Genomics* 111, 906–912. <https://doi.org/10.1016/j.ygeno.2018.05.020>

- Bain, O., Wahl, G., Renz, A., 1993. *Onchocerca ramachandrini* n. sp. from the warthog in Cameroon. *Ann Parasitol Hum Comp* 68, 139–143.
<https://doi.org/10.1051/parasite/1993683139>
- Bhandari, S., Adhikari, B., Baral, K., Subedi, S.C., 2022. Greater one-horned rhino (*Rhinoceros unicornis*) mortality patterns in Nepal. *Glob Ecol Conserv* 38, e02189.
<https://doi.org/10.1016/j.gecco.2022.e02189>
- Bohara, T.P., Shrestha, S.P., 2015. A study on cattle tick and tick borne pathogens of Mid Western Nepal. *Nepalese Vet. J* 33, 23–27.
- Botto, C., Arango, M., Yarzabal, L., 1984. Onchocerciasis in Venezuela: Prevalence of microfilaraemia in Amerindians and morphological characteristics of the microfilariae from the Upper Orinoco Focus. *Tropenmed Parasitol* 35, 167–173.
- Cambra-Pellejà, M., Gandasegui, J., Balaña-Fouce, R., Muñoz, J., Martínez-Valladares, M., 2020. Zoonotic implications of *Onchocerca* species on human health. *Pathogens*. <https://doi.org/10.3390/pathogens9090761>
- Chatanga, E., Kainga, H., Razemba, T., Ssuna, R., Swennen, L., Hayashida, K., Sugimoto, C., Katakura, K., Nonaka, N., Nakao, R., 2021. Molecular detection and characterization of tick-borne hemoparasites and Anaplasmatidae in dogs in major cities of Malawi. *Parasitol Res* 120, 267–276. <https://doi.org/10.1007/s00436-020-06967-y>
- CNP, 2012. Biodiversity: Chitwan National Park, Chitwan. .
- Correa, C.C., Ballard, J.W.O., 2016. *Wolbachia* associations with insects: Winning or losing against a master manipulator. *Front Ecol Evol*.
<https://doi.org/10.3389/fevo.2015.00153>
- Dinerstein, E., 2003. The return of the unicorns: the natural history and conservation of the greater one-horned rhinoceros. Columbia University Press.
- Ellis, S., Talukdar, B.K., 2019. *Rhinoceros unicornis*: The IUCN Red List of Threatened Species: e. T19496A18494149.
- Govender, D., Oosthuisen, M.C., Penzhorn, B.L., 2011. Piroplasm parasites of white rhinoceroses (*Ceratotherium simum*) in the Kruger National Park, and their relation to anaemia. *J S Afr Vet Assoc* 82, 36–40. <https://doi.org/10.4102/jsava.v82i1.31>
- Grácio, A.J.S., Richter, J., Komnenou, A.T., Grácio, M.A., 2015. Onchocerciasis caused by *Onchocerca lupi*: an emerging zoonotic infection. Systematic review. *Parasitol Res*. <https://doi.org/10.1007/s00436-015-4535-7>
- Gupta, V., Gupta, R., Shrestha, S.P., 2013. Infectivity of *Theileria annulata* in *Hyalomma* ticks of Eastern Terai districts, Nepal. *Nepalese Journal of Zoology* 1, 15–23.
- Höfle, U., Vicente, J., Nagore, D., Hurtado, A., Peña, A., Fuente, J. de la, Gortazar, C., 2004. The risks of translocating wildlife: Pathogenic infection with *Theileria sp.* and *Elaeophora elaphi* in an imported red deer. *Vet Parasitol* 126, 387–395.
<https://doi.org/10.1016/j.vetpar.2004.07.026>
- Jhala, H., Qureshi, Q., Jhala, Y., Reports, S.B.-S., undefined, 2021. Feasibility of reintroducing grassland megaherbivores, the greater one-horned rhinoceros, and swamp buffalo within their historic global range. *Springer* 11, 4469.
<https://doi.org/10.1038/s41598-021-83174-4>
- Kamau, J., De Vos, A.J., Playford, M., Salim, B., Kinyanjui, P., Sugimoto, C., 2011. Emergence of new types of *Theileria orientalis* in Australian cattle and possible cause of theileriosis outbreaks. *Parasit Vectors* 4. <https://doi.org/10.1186/1756-3305-4-22>

- Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Mol Biol Evol* 30, 772–780. <https://doi.org/10.1093/MOLBEV/MST010>
- Kaur, R., Shropshire, J.D., Cross, K.L., Leigh, B., Mansueto, A.J., Stewart, V., Bordenstein, Sarah R., Bordenstein, Seth R., 2021. Living in the endosymbiotic world of *Wolbachia*: A centennial review. *Cell Host Microbe*. <https://doi.org/10.1016/j.chom.2021.03.006>
- King'ori, E., Obanda, V., Chiyo, P.I., Soriguer, R.C., Morrondo, P., Angelone, S., 2019. Molecular identification of *Ehrlichia*, *Anaplasma*, *Babesia* and *Theileria* in African elephants and their ticks. *PLoS One* 14. <https://doi.org/10.1371/journal.pone.0226083>
- King'ori, E., Waiguchu, G., Ruoro, M., Muriithi, K., Mumbi, C., Omondi, M., Aminga, D., Angwenyi, S., Mijele, D., Chiyo, P.I., 2024. Identifying Risk Factors for *Stephanofilaria*-Caused Ulcerative Dermal Lesions, in Black and White Rhinoceros' Meta-Population in Kenya. *Transbound Emerg Dis* 2024, 1–14. <https://doi.org/10.1155/2024/2655970>
- Kumar, S., Stecher, G., Tamura, K., 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33, 1870–1874. <https://doi.org/10.1093/MOLBEV/MSW054>
- Leigh, J.W., Bryant, D., 2015. popart: full-feature software for haplotype network construction. *Methods Ecol Evol* 6, 1110–1116. <https://doi.org/10.1111/2041-210X.12410>
- Manoj, R.R.S., Latrofa, M.S., Epis, S., Otranto, D., 2021. *Wolbachia*: endosymbiont of onchocercid nematodes and their vectors. *Parasit Vectors*. <https://doi.org/10.1186/s13071-021-04742-1>
- Mans, B.J., Pienaar, R., Latif, A.A., 2015. A review of *Theileria* diagnostics and epidemiology. *Int J Parasitol Parasites Wildl*. <https://doi.org/10.1016/j.ijppaw.2014.12.006>
- Martin, C., Gavotte, L., 2010. The bacteria *Wolbachia* in filariae, a biological Russian dolls' system: New trends in antifilarial treatments. *Parasite* 17, 79–89. <https://doi.org/10.1051/parasite/2010172079>
- Masatani, T., Hayashi, K., Andoh, M., Tateno, M., Endo, Y., Asada, M., Kusakisako, K., Tanaka, T., Gokuden, M., Hozumi, N., Nakadohono, F., Matsuo, T., 2017. Detection and molecular characterization of *Babesia*, *Theileria*, and *Hepatozoon* species in hard ticks collected from Kagoshima, the southern region in Japan. *Ticks Tick Borne Dis* 8, 581–587. <https://doi.org/10.1016/j.ttbdis.2017.03.007>
- Masatani, T., Kojima, I., Tashiro, M., Yamauchi, K., Fukui, D., Ichikawa-Seki, M., Harasawa, R., 2021. Molecular detection of filarial nematode parasites in japanese black bears (*Ursus thibetanus japonicus*) from iwate prefecture, japan. *Journal of Veterinary Medical Science*. <https://doi.org/10.1292/jvms.20-0466>
- Mediannikov, O., Ranque, S., 2018. Mansonellosis, the most neglected human filariasis. *New Microbes New Infect* 26, S19–S22.
- Muhanguzi, D., Matovu, E., Waiswa, C., 2010. Prevalence and Characterization of *Theileria* and *Babesia* Species in Cattle under Different Husbandry Systems in Western Uganda. *Int J Anim Vet Adv* 2, 51–58.
- Munson, L., Terio, K.A., Kock, R., Mlengeya, T., Roelke, M.E., Dubovi, E., Summers, B., Sinclair, A.R.E., Packer, C., 2008. Climate extremes promote fatal co-infections during canine distemper epidemics in African lions. *PLoS One* 3. <https://doi.org/10.1371/journal.pone.0002545>

- Mutinda, M., Otiende, M., Gakuya, F., Kariuki, L., Obanda, V., Ndeere, D., Ndambiri, E., Kariuki, E., Lekolool, I., Soriguer, R.C., Rossi, L., Alasaad, S., 2012. Putative filariasis outbreak in white and black rhinoceros at Meru National Park in Kenya. *Parasit Vectors* 5. <https://doi.org/10.1186/1756-3305-5-206>
- Nijhof, A.M., Penzhorn, B.L., Lynen, G., Mollel, J.O., Morkel, P., Bekker, C.P.J., Jongejan, F., 2003. *Babesia bicornis* sp. nov. and *Theileria bicornis* sp. nov.: Tick-borne parasites associated with mortality in the black rhinoceros (*Diceros bicornis*). *J Clin Microbiol* 41, 2249–2254. <https://doi.org/10.1128/jcm.41.5.2249-2254.2003>
- Nijhof, A.M., Pillay, V., Steyl, J., Prozesky, L., Stoltz, W.H., Lawrence, J.A., Penzhorn, B.L., Jongejan, F., 2005. Molecular characterization of *Theileria* species associated with mortality in four species of African antelopes. *J Clin Microbiol* 43, 5907–5911. <https://doi.org/10.1128/JCM.43.12.5907-5911.2005>
- Otiende, M.Y., Kivata, M.W., Jowers, M.J., Makumi, J.N., Runo, S., Obanda, V., Gakuya, F., Mutinda, M., Kariuki, L., Alasaad, S., 2016. Three Novel Haplotypes of *Theileria bicornis* in Black and White Rhinoceros in Kenya. *Transbound Emerg Dis* 63, e144–e150.
- Otiende, M.Y., Kivata, M.W., Makumi, J.N., Mutinda, M.N., Okun, D., Kariuki, L., Obanda, V., Gakuya, F., Mijele, D., Soriguer, R.C., Alasaad, S., 2015. Epidemiology of *Theileria bicornis* among black and white rhinoceros metapopulation in Kenya. *BMC Vet Res* 11, 1–7. <https://doi.org/10.1186/S12917-014-0316-2/FIGURES/2>
- Otranto, D., Brianti, E., Dantas-Torres, F., Weigl, S., Latrofa, M.S., Gaglio, G., Cauquil, L., Giannetto, S., Bain, O., 2011. Morphological and molecular data on the dermal microfilariae of a species of *Cercopithifilaria* from a dog in Sicily. *Vet Parasitol* 182, 221–229. <https://doi.org/10.1016/j.vetpar.2011.05.043>
- Otranto, D., Cantacessi, C., Dantas-Torres, F., Brianti, E., Pfeiffer, M., Genchi, C., Guberti, V., Capelli, G., Deplazes, P., 2015. The role of wild canids and felids in spreading parasites to dogs and cats in Europe. Part II: Helminths and arthropods. *Vet Parasitol* 213, 24–37. <https://doi.org/10.1016/j.vetpar.2015.04.020>
- Otranto, D., Dantas-Torres, F., 2010. Canine and feline vector-borne diseases in Italy: current situation and perspectives.
- Otranto, D., Deplazes, P., 2019. Zoonotic nematodes of wild carnivores. *Int J Parasitol Parasites Wildl*. <https://doi.org/10.1016/j.ijppaw.2018.12.011>
- Papini, R.A., Lubas, G., Sgorbini, M., 2020. Incidental Detection of *Onchocerca* Microfilariae in Donkeys (*Equus asinus*) in Italy: Report of Four Cases. *Front Vet Sci* 7. <https://doi.org/10.3389/fvets.2020.569916>
- Parola, P., Roux, V., Camicas, J.-L., Baradji, I., Brouqui, P., Raoult, D., 2000. Detection of *Ehrlichiae* in African ticks by polymerase chain reaction. *Trans R Soc Trop Med Hyg* 94, 707–708.
- Perles, L., Otranto, D., Barreto, W.T.G., de Macedo, G.C., Lia, R.P., Mendoza-Roldan, J.A., Herrera, H.M., de Oliveira, C.E., Machado, R.Z., André, M.R., 2023. *Mansonella* sp. and associated *Wolbachia* endosymbionts in ring-tailed coatis (*Nasua nasua*) in periurban areas from Midwestern Brazil. *Int J Parasitol Parasites Wildl* 22, 14–19. <https://doi.org/10.1016/j.ijppaw.2023.08.002>
- Pfister, S., Oosthuizen, M.C., Bosman, A.-M., Vorster, I., Penzhorn, B.L., 2011. Tick-borne blood parasites in nyala (*Tragelaphus angasii*, Gray 1849) from KwaZulu-Natal, South Africa. *Vet Parasitol* 176, 126–131.
- Poudel, S., Gurung, Y., Shrestha, S., 2023. Prevalence of tick-borne haemoparasitic diseases and haemato-biochemical changes in cattle of Kathmandu Valley. *Nepalese Journal of Agricultural Sciences* 24, 47–58.

- Pupić-Bakrač, A., Pupić-Bakrač, J., Beck, A., Jurković, D., Polkinghorne, A., Beck, R., 2021. *Dirofilaria repens* microfilaremia in humans: Case description and literature review. *One Health*. <https://doi.org/10.1016/j.onehlt.2021.100306>
- Qiu, Y., Nakao, R., Thu, M.J., Akter, S., Alam, M.Z., Kato, S., Katakura, K., Sugimoto, C., 2016. Molecular evidence of spotted fever group rickettsiae and Anaplasmataceae from ticks and stray dogs in Bangladesh. *Parasitol Res* 115, 949–955.
- Rambaut, A., 2012. FigTree v1. 4. Molecular evolution, phylogenetics and epidemiology. Edinburgh: University of Edinburgh, Institute of Evolutionary Biology.
- Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a Large Model Space. *Syst Biol* 61, 539–542. <https://doi.org/10.1093/SYSBIO/SYS029>
- Schulz, K., Kluge, E.B., 1960. Dermatitis in the black rhinoceros (*Diceros bicornis*) due to filariasis. *J.S.A.V.M.A.* 2, 265–269.
- Shelley, A.J., Coscarón, S., 2001. Simuliid Blackflies (Diptera: Simuliidae) and Ceratopogonid Midges (Diptera: Ceratopogonidae) as Vectors of *Mansonella ozzardi* (Nematoda: Onchocercidae) in Northern Argentina, Mem Inst Oswaldo Cruz, Rio de Janeiro.
- Shrestha, S.P., Prajapati, M., Shrestha, R., Kunwar, B.L., Shrestha, S.P., 2017. Study on piroplasmosis and anaplasmosis: killer diseases of dairy cattle, in: Proceeding of the 10th National Workshop on Livestock and Fisheries Research in Nepal. p. 7.
- Subedi, N., 2012. Effect of Mikania micrantha on the demography, habitat use, and nutrition of Greater One-horned Rhinoceros in Chitwan National Park, Nepal. Dehradun: Forest Research Institute University.
- Sugimoto, C., Fujisaki, K., 2002. Non-transforming *Theileria* parasites of ruminants. *Theileria* 93–106.
- Tanabe, A. s., 2008. Life is fifthdimension. - Official Web Site of Akifumi S. Tanabe [WWW Document]. URL <https://www.fifthdimension.jp/> (accessed 9.26.23).
- Tanabe, A.S., 2011. Kakusan4 and Aminosan: two programs for comparing nonpartitioned, proportional and separate models for combined molecular phylogenetic analyses of multilocus sequence data. *Mol Ecol Resour* 11, 914–921. <https://doi.org/10.1111/J.1755-0998.2011.03021.X>
- Taylor, M.J., Bandi, C., Hoerauf, A., 2005. *Wolbachia*. Bacterial Endosymbionts of Filarial Nematodes. *Adv Parasitol* 60, 245–284. [https://doi.org/10.1016/S0065-308X\(05\)60004-8](https://doi.org/10.1016/S0065-308X(05)60004-8)
- Turb, M.E., Zambon, E., Zannoni, A., Russo, S., Gentilini, F., 2012. Detection of *Wolbachia* DNA in blood for diagnosing filaria-associated syndromes in cats. *J Clin Microbiol* 50, 2624–2630. <https://doi.org/10.1128/JCM.00528-12>
- Uni, S., Bain, O., Agatsuma, T., Harada, M., Torii, H., Fukuda, M., Takaoka, H., 2007. *Onchocerca eberhardi* n. sp. (nematoda: Filarioidea) from sika deer in Japan; Relationships between species parasitic in cervids and bovids in the holarctic region. *Parasite*. <https://doi.org/10.1051/parasite2007143199>
- Uni, S., Bain, O., Takaoka, H., 2004. Affinities between Cutifilaria (Nematoda: Filarioidea), parasites of deer, and *Mansonella* as seen in a new onchocercid, *M. (C.) perforata* n. sp., from Japan. *Parasite* 11, 131–140. <https://doi.org/10.1051/parasite/2004112131>
- Uni, S., Fukuda, M., Agatsuma, T., Bain, O., Otsuka, Y., Nakatani, J., Matsubayashi, M., Harada, M., Omar, H., Ramli, R., Hashim, R., Azirun, M.S., Takaoka, H.,

2015. *Onchocerca takaokai* n. sp. (Nematoda: Filarioidea) in Japanese wild boars
(*Sus scrofa leucomystax*): Description and molecular identification of intradermal
females. *Parasitol Int* 64, 493–502. <https://doi.org/10.1016/j.parint.2015.07.001>

Yagi, K., Bain, O., Shoho, C., 1994. *Onchocerca suzukii* n. sp. et *O.skrjabini* (*O.*
tarsicola) parasites d'un bovidé relique du Japon, *Capricornis crispus*. *Parasite* 1,
349–356. <https://doi.org/10.1051/parasite/1994014349>

Yam, J., Gestier, S., Bryant, B., Campbell-Ward, M., Bogema, D., Jenkins, C., 2018.
The identification of *Theileria bicornis* in captive rhinoceros in Australia. *Int J*
Parasitol Parasites Wildl 7, 85–89. <https://doi.org/10.1016/J.IJPPAW.2017.12.003>

Zimmermann, D.E., Penzhorn, B.L., Vorster, I., Troskie, M., Oosthuizen, M.C., 2021.
Babesia bicornis, *Theileria bicornis* and *Theileria equi* in metapopulations of two
black rhinoceros (*Diceros bicornis*) subspecies in South Africa and their potential
impact on conservation. *Ticks Tick Borne Dis* 12, 101635.

Table 1. Primers used in this study.

Primers	Sequence (5'-3')	Target organism(s)	Target gene	Purpose	Product size	Reference
BTH 1st F	GTGAAACTGCGAATGGCTCATTAC	<i>Babesia/</i> <i>Theileria/</i> <i>Hepatozoon</i>	18S rDNA	1 st PCR	1400-1600	(Masatani et al., 2017)
BTH 1st R BTH 2nd F	AAGTGATAAGGTTACAAAACTTCCC GGCTCATTACAACAGTTATAGTTTATTTG	<i>Babesia/</i> <i>Theileria/</i> <i>Hepatozoon</i>	18S rDNA	2 nd PCR	1400-1600	(Masatani et al., 2017)
BTH 2nd R BTH18SinterF	CGGTCCGAATAATTCACCGGAT ATTTCCGACTCCTTCAGCA	<i>Babesia/</i> <i>Theileria/</i> <i>Hepatozoon</i>	18S rDNA	Sequencing		(Ogata et al., 2021)
BTH18SinterR EHR16SD	AACTAAGAACGGCCATGCAC GGTACCYACAGAAGAAGTCC/	Anaplasmataceae	16S rRNA	PCR	345	(Parola et al., 2000)
EHR16SR HS1-F	TAGCACTCATCGTTTACAGC CGYCAGTGGGTGGTAATGAA	Anaplasmataceae	<i>groEL</i>	1 st PCR	1300	(Rar et al., 2011)
HS6-R HS3-F	CCWCCWGGTACWACACCTTC ATAGTYATGAAGGAGAGTGAT	<i>Anaplasma</i>	<i>groEL</i>	2 nd PCR	1256	(Liz et al., 2000)
HSV-R groEL_fwd3	TCAACAGCAGCTCTAGTWG TGGCAAATGTAGTTGTAACAGG	<i>Ehrlichia</i>	<i>groEL</i>	2 nd PCR	1100	(Gofton et al., 2016)
groEL_rev2 F4b	GCCGACTTTTAGTACAGCAA CCAGGCTTTATGTCAACTGC	Anaplasmataceae	<i>gltA</i>	1 st PCR	800	(Inokuma et al., 2005)
R1b EHR-CS136F	CGATGACCAAAACCCAT TTYATGTCTYACTGCTGCKTG	<i>Anaplasma</i> and <i>Ehrlichia</i>	<i>gltA</i>	2 nd PCR	650	(Inokuma et al., 2005)
EHR-CS778R EHR16SD	GCNCCMCCATGMGCTGG GGTACCYACAGAAGAAGTCC	<i>Wolbachia</i>	16S rRNA	PCR	1300	(Moustafa et al., 2022)
1513R Nematode1	ACGGYTACCTTGTTACGACTT GCGGAGGAAAAGAACTAA	Nematode	28S rDNA	PCR	765	(Sehgal et al., 2005)
Nematode2 COLintF	ATCCGTGTTTCAAGACGGG TGATTGGTGGTTTTGGTAA	<i>Onchocerca</i>	COI	PCR	650	(Casiraghi et al., 2001)
COLintR MyManF	ATAAGTACGAGTATCAATATC GAAGCTGAGGCTCAAGCAAT	<i>Mansonella</i>	<i>myoHC</i>	PCR	554	(Moraes et al., 2022)
MyManR h70ManF	TCTGTTTTGCTCATCGCATT TGAGACAGCTGGAGGTGTTATG	<i>Mansonella</i>	<i>hsp70</i>	PCR	484	(Moraes et al., 2022)
h70ManR InFusion_EHR16SD InFusion_1513R	ATCTTTCTGTGCCTCATCATCTG AAACGACGGCCAGTGGTACTCACAGAAGAAGTCTGGC GACCATGATTACGCCACGGCTACCTTGTTACGACTTCAC	<i>Wolbachia</i>	16S rDNA	cloning	1300	This study

Note: PCR, Polymerase chain reaction; NA, not applicable.

Table 2: *Theileria bicornis* detected in greater one-horned rhinoceroses in Nepal.

Variable	Category	No. tested	Positive no. (%)	Odds ratio (95% CI)	P- value
Age	Young (0 year)	6	4 (67)	Reference	0.62
	Subadult (3-6 years)	4	3 (<1)	1.5 (0.09-25.41)	
	Adult (>6 years)	26	20 (77)	1.67 (0.24-11.45)	
Location	National park	19	13 (68)	0.46 (0.09-2.25)	0.45
	Buffer zone	17	14 (82)	2.15 (0.44-10.44)	
Sex	Female	15	11 (73)	0.86 (0.19-3.94)	1
	Male	21	16 (76)	1.16 (0.25-5.34)	
Total		36	27(75)		




Figure1: Visualization of haplotype network of *Theliera bicornis* using 18S rRNA gene sequences. Median-joining haplotype networks was constructed from 18S rDNA sequences. The circles represent the *T. bicornis* haplotypes. The color and size of each circle correspond to the location where the haplotype was detected and the haplotype number. For the haplotype network of *T. bicornis* sequences of haplotypes from Australia (H2 and H4), Kenya (H1, H2 and H3), South Africa (H1 and H3) and Nepal (H8 and H9) were included.




Figure 2: Geographical distribution of *Theileria bicornis* haplotypes H8 and H9 in Chitwan National Park of Nepal.




Figure 3. Bayesian phylogenetic tree based on 16S rRNA gene sequences of *Wolbachia*. Nodes with posterior probabilities greater than 0.75 are indicated by black dots. Tip labels display GenBank accession numbers and their host, with green indicating arthropod hosts and blue indicating filarial nematode hosts and red indicating sequence obtained in this study. The alphabets A-F represent super group of *Wolbachia*.




Fig 4. Bayesian phylogenetic tree based on 28S rRNA gene sequences of filarial nematodes. Black dots indicate nodes with posterior probability of > 0.70.




Fig 5: Bayesian phylogenetic tree based on COI gene of filarial nematodes. Black dots indicate nodes with posterior probability of > 0.70.




Fig 6. A Bayesian Inference phylogenetic tree of *hsp70* gene of *Mansonella* spp. Black dots indicate nodes with posterior probability of > 0.75.




Fig 7: A Bayesian Inference phylogenetic tree of *myoHC* gene sequences of *Mansonella* spp. Black dots indicate nodes with posterior probability of > 0.75.