



Title	Anisotropic skyrmion crystal on a centrosymmetric square lattice under an in-plane magnetic field
Author(s)	Hayami, Satoru
Citation	Journal of magnetism and magnetic materials, 604, 172293 https://doi.org/10.1016/j.jmmm.2024.172293
Issue Date	2024-08-15
Doc URL	https://hdl.handle.net/2115/97229
Rights	© 2024. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	main_resub.pdf



Anisotropic skyrmion crystal on a centrosymmetric square lattice under an in-plane magnetic field

Satoru Hayami

Graduate School of Science, Hokkaido University, Sapporo 060-0810, Japan

Abstract

We investigate the instability toward a skyrmion crystal under an in-plane magnetic field in centrosymmetric tetragonal magnets. By performing the simulated annealing for a spin model on the square lattice, we find that the interplay among the easy-plane two-spin anisotropic interaction, positive biquadratic interaction, and high-harmonic wave-vector interaction plays an important role in stabilizing the skyrmion crystal in the in-plane external magnetic field. The obtained skyrmion crystal consists of a pair of vortex and antivortex with opposite polarization in terms of the out-of-plane spin component. We also show that various multiple- Q states emerge depending on the model parameters. Our results indicate the possibility of topological spin textures in centrosymmetric tetragonal magnets with easy-plane magnetic anisotropy when the ground-state spin configuration at zero field is characterized by a single- Q in-plane cycloidal spiral state.

Keywords: multiple- Q magnetic state, skyrmion crystal, square lattice, noncoplanar spin texture, magnetic frustration

1. Introduction

A magnetic skyrmion, which is characterized by a swirling spin texture so as to have an integer topological (skyrmion) number, has attracted many researchers owing to its rich physical properties [1, 2, 3, 4, 5, 6, 7]. Through
5 extensive studies in both theory and experiments, a magnetic skyrmion crystal (SkX), which is a periodic array of the magnetic skyrmion, has been found in materials with both noncentrosymmetric and centrosymmetric lattice structures [8, 9, 10, 11, 12, 13, 14, 15, 16, 17], and their stabilization mechanisms have been elucidated; the Dzyaloshinskii-Moriya (DM) interaction becomes an impor-
10 tant source of the SkX in noncentrosymmetric magnets [18, 19, 20, 21, 22, 23, 24],

Email address: hayami@phys.sci.hokudai.ac.jp (Satoru Hayami)

while frustrated exchange interactions, multiple spin interactions, and/or symmetric anisotropic interactions play an important role in inducing the SkX in centrosymmetric magnets [25, 26, 27, 28]. Here and hereafter, we refer to the magnetic states with integer skyrmion number in the magnetic unit cell as the SkX irrespective of constituent topological defects.

In addition to the exchange interactions that are inherent to the materials, an external magnetic field is another essential ingredient to induce the SkX, where the instability toward the SkX depends on the magnetic field direction; the magnetic field perpendicular to the ordering wave vectors tends to stabilize the SkX [20, 23, 29]. Meanwhile, it was shown that the SkX and similar topological spin crystals can be stabilized even in the magnetic field parallel to the ordering wave vectors. For example, a magnetic bimeron crystal, which is formed by a periodic alignment of a pair of two merons, can appear under the in-plane magnetic field in noncentrosymmetric magnets with the DM interaction [30, 31, 32]. Another example is the SkX in centrosymmetric magnets, which is stabilized without the DM interaction by considering the easy-plane magnetic anisotropy and a triangular-lattice geometry [33, 34]. Meanwhile, it has not been clarified the possibility of the SkXs under the in-plane field on a centrosymmetric square lattice, where the effect of geometrical frustration does not appear. Although the emergence of the SkXs on such a square lattice in the out-of-plane magnetic field has been shown by the theoretical model calculations [35, 36, 37, 38] and experiments in GdRu_2Si_2 [16, 39, 40, 41, 42] and EuAl_4 [43, 44, 45, 46, 47, 48, 49], its stabilization condition under the in-plane magnetic field has been unclear yet.

In the present study, we investigate the stability of the in-plane field-induced SkX in the centrosymmetric tetragonal system, which is different from the out-of-plane field-induced SkX in previous work [41]. Through the numerical calculations based on the simulated annealing for the phenomenological spin model, we find the conditions to stabilize the SkX on the square lattice under the in-plane field. We show that the interplay among the easy-plane two-spin anisotropic interaction, positive biquadratic interaction, and high-harmonic wave-vector interaction is important in order to realize the SkX. The obtained SkX is characterized by an anisotropic double- Q spin modulation, which reflects the lack of the fourfold rotational symmetry under the in-plane field. We also find various double- Q instabilities depending on the magnitude of the magnetic field and easy-axis anisotropic interaction.

The rest of this paper is organized as follows. In Sec. 2, we present the effective spin model consisting of bilinear and biquadratic spin interactions on the square lattice. We also outline the numerical simulated annealing. In Sec. 3, we show the results of the magnetic phase diagram at low temperatures. We describe the spin and scalar chirality configurations in real and momentum spaces in each phase. Section 4 is devoted to a summary of this paper. In Appendix Appendix A, we show two magnetic phase diagrams calculated by using different model parameters in the main text.

55 2. Model and method

2.1. Model

In order to investigate the essence in terms of the stability of the in-plane field-induced SkX at low temperatures, we analyze the spin model with the momentum-resolved interaction on the two-dimensional centrosymmetric square
60 lattice. We suppose the clean system, where the effect of lattice defects or impurities is negligible. We set the lattice constant of the square lattice as unity. The model Hamiltonian is given by

$$\begin{aligned} \mathcal{H} = & 2 \sum_{\nu=1,2} \left(-J\Gamma_{\nu} + \frac{K}{N}\Gamma_{\nu}^2 \right) \\ & + 2 \sum_{\nu=3,4} \left(-J'\Gamma_{\nu} + \frac{K'}{N}\Gamma_{\nu}^2 \right) - H \sum_i S_i^x, \end{aligned} \quad (1)$$

with

$$\Gamma_{\nu} = \mathbf{S}_{\mathbf{Q}_{\nu}} \cdot \mathbf{S}_{-\mathbf{Q}_{\nu}} - I^{\text{EPA}} S_{\mathbf{Q}_{\nu}}^z S_{-\mathbf{Q}_{\nu}}^z. \quad (2)$$

The first term in Eq. (1) represents the interaction at the wave vector $\mathbf{Q}_1 = (\pi/5, 0)$ and $\mathbf{Q}_2 = (0, \pi/5)$, where $\mathbf{S}_{\mathbf{Q}_{\nu}} = (S_{\mathbf{Q}_{\nu}}^x, S_{\mathbf{Q}_{\nu}}^y, S_{\mathbf{Q}_{\nu}}^z)$ in Γ_{ν} corresponds to the spin moment at wave vector $\mathbf{Q}_{\nu}$; $\mathbf{S}_{\mathbf{Q}_{\nu}}$ is related to the classical localized spin $\mathbf{S}_i = (S_i^x, S_i^y, S_i^z)$ with fixed length $|\mathbf{S}_i| = 1$ via the Fourier transformation. N represents the number of sites in the system. The first term consists of the bilinear exchange interaction with the coupling constant $J > 0$ and the positive
70 biquadratic interaction with the coupling constant $K > 0$, the latter of which often appears when the Fermi surface is strongly nested by $\mathbf{Q}_{\nu}$ in the itinerant electron systems [50, 51, 52]. The factor 2 stands for the contributions from $-\mathbf{Q}_1$ and $-\mathbf{Q}_2$. The interaction J tends to favor the single- Q spiral state with $\mathbf{Q}_1$ or $\mathbf{Q}_2$, while the interaction K tends to favor the multiple- Q state composed
75 of the $\mathbf{Q}_1$ and $\mathbf{Q}_2$ modulations. In addition, we introduce the effect of the easy-plane two-spin anisotropic interaction by setting the different form factors for the xy and z spin components, as shown by Γ_{ν} in Eq. (2); $I^{\text{EPA}} > 0$ denotes the degree of easy-plane magnetic anisotropy; the instability toward the SkX in the case of $I^{\text{EPA}} < 0$ has been investigated in [41]. We do not consider the
80 effect of the DM interaction, which often leads to the SkX [23, 53], owing to the centrosymmetric lattice structure.

The second term in Eq. (1) represents the momentum-resolved interaction at the wave vector $\mathbf{Q}_3 = (\pi/5, \pi/5)$ and $\mathbf{Q}_4 = (-\pi/5, \pi/5)$. We set the different interaction constants J' and K' for the bilinear and biquadratic exchange interactions, respectively. The third term represents the effect of an external in-plane
85 magnetic field along the x direction, where H represents the magnitude of the magnetic field. The spin model in Eq. (1) can be used to investigate the instability toward the SkXs in other lattice structures by appropriately considering the ordering wave vectors satisfying the lattice symmetry.

90 In the following calculations, we parametrize the model parameters as $J = 1$, $K = 0.2$, $J' = 0.6$, and $K' = (J'/J)^2 K$, whereas we change I^{EPA} and H ; J is the energy unit of the spin model in Eq. (1). The relation $J > J'$ means that the interactions at $\mathbf{Q}_1$ and $\mathbf{Q}_2$ are dominant, and those at $\mathbf{Q}_3$ and $\mathbf{Q}_4$ give the energy correction. Especially, the latter interaction J' plays an
 95 important role in inducing the multiple- Q states, since $\mathbf{Q}_3$ and $\mathbf{Q}_4$ correspond to the high-harmonic wave vectors of $\mathbf{Q}_1$ and $\mathbf{Q}_2$, i.e., $\mathbf{Q}_3 = \mathbf{Q}_1 + \mathbf{Q}_2$ and $\mathbf{Q}_4 = -\mathbf{Q}_1 + \mathbf{Q}_2$ [38, 54, 41]. As will be discussed in Sec. 3, both K and J' are important to stabilize the SkX under the in-plane magnetic field.

2.2. Method

100 We calculate the stable spin configurations of the model in Eq. (1) by performing the numerical simulated annealing following the manner in Ref. [52]. First, we start the simulations from a random spin configuration at a high temperature $T_0 = 1.5$. Then, we gradually reduce the temperature by the ratio of $T_{n+1} = \alpha T_n$ with $\alpha = 0.999999$ up to the final temperature $T = 10^{-5}$, where T_n
 105 is the n th temperature. We update all the spins one by one in real space based on the single-spin-flip Metropolis algorithm at each temperature. When we reach the final temperature, we perform 10^5 - 10^6 Monte Carlo sweeps for thermalization and measurements. We perform the above procedure while changing I^{EPA} and H . When we optimize the spin configuration near the phase boundary, we
 110 also start the simulations from the spin patterns obtained at low temperatures. We set the system size with $N = 10^2$ under the periodic boundary condition. We confirm that the following results are not altered for $N = 60^2$.

2.3. Physical quantities

We calculate the $\mathbf{Q}_\nu$ component of magnetic moments, which is given by

$$m_{\mathbf{Q}_\nu}^\eta = \sqrt{\frac{S_s^{\eta\eta}(\mathbf{Q}_\nu)}{N}}, \quad (3)$$

115 where $\eta = x, y, z$ and $S_s^{\eta\eta}(\mathbf{Q}_\nu)$ is the spin structure factor defined by

$$S_s^{\eta\eta}(\mathbf{Q}_\nu) = \frac{1}{N} \sum_{i,j} S_i^\eta S_j^\eta e^{i\mathbf{Q}_\nu \cdot (\mathbf{r}_i - \mathbf{r}_j)}. \quad (4)$$

Here, $\mathbf{r}_i$ is the position vector at site i on the square lattice. The net magnetization is given by

$$M^\eta = \frac{1}{N} \sum_i S_i^\eta. \quad (5)$$

We also calculate the scalar chirality, which is given by

$$\chi^{\text{sc}} = \frac{1}{2N} \sum_i \sum_{\delta, \delta' = \pm 1} \delta\delta' \mathbf{S}_i \cdot (\mathbf{S}_{i+\delta\hat{x}} \times \mathbf{S}_{i+\delta'\hat{y}}), \quad (6)$$

where $\hat{x}$ ($\hat{y}$) represents a shift by lattice constant in the x (y) direction on the
 120 square lattice. The nonzero χ^{sc} indicates the emergence of the topological spin textures, which includes the SkX.




Figure 1: (a) Magnetic phase diagram of the model in Eq. (1) on the square lattice by changing the easy-plane anisotropy I^{EPA} and the in-plane magnetic field H , which is obtained by the simulated annealing at $K = 0.2$ and $J' = 0.6$. $1Q$ denotes the single- Q states, and $2Q'$ and $2Q$ represent the anisotropic and isotropic double- Q states, respectively. IS, OS, CS, F, VC, and SkX stand for the in-plane spiral, out-of-plane spiral, chiral stripe, fan, vortex crystal, and skyrmion crystal, respectively. Real-space spin configurations in (b) the $2Q'$ CS I state at $I^{\text{EPA}} = 0$ and $H = 0.05$, (c) the $1Q$ OS state at $I^{\text{EPA}} = 0$ and $H = 1.2$, (d) the $2Q'$ CS II state at $I^{\text{EPA}} = 0.06$ and $H = 0$, (e) the SkX at $I^{\text{EPA}} = 0.06$ and $H = 0.7$, (f) the $2Q$ F state at $I^{\text{EPA}} = 0.06$ and $H = 1.85$, (g) the $2Q'$ I state at $I^{\text{EPA}} = 0.12$ and $H = 0.5$, (h) the $1Q$ IS state at $I^{\text{EPA}} = 0.4$ and $H = 0$, and (i) the $2Q'$ VC state at $I^{\text{EPA}} = 0.4$ and $H = 0.75$. In (b)–(i), the arrows represent the direction of the in-plane spin and the color shows its z component. In (e), the triangle and square stand for the vortex and antivortex, respectively.

3. Results

By performing the simulated annealing for the model in Eq. (1), we obtain the magnetic phase diagram in the plane of I^{EPA} and H at low temperature in Fig. 1(a). Various phases appear depending on the model parameters; the SkX appears in the intermediate I^{EPA} and H . In the following, we distinguish the obtained states by the Q_1 and Q_2 components of the magnetic moments without referring to the Q_3 and Q_4 components, since the former amplitude is larger than the latter one; we term a state with nonzero $(m_{Q_1}^\eta)^2$ or $(m_{Q_2}^\eta)^2$ as a single- Q ($1Q$) state and that with both nonzero $(m_{Q_1}^\eta)^2$ and $(m_{Q_2}^\eta)^2$ as a




Figure 2: H dependence of (a) the magnetization M^η , (b) the scalar chirality χ^{sc} , and (c) Q_1 and Q_2 components of squared magnetic moments $(m_{Q_{1,2}}^\eta)^2$ for $\eta = x, y, z$ at $I^{\text{EPA}} = 0$. The vertical dashed line represents the phase boundary between different spin states; the corresponding phases are presented above in (a).

double- Q ($2Q$) state. We present the data in terms of the uniform magnetization, scalar chirality, and the Q_1 and Q_2 components of the magnetic moments in Figs. 2–5 for different I^{EPA} .

For $I^{\text{EPA}} = 0$, the zero-field to low-field state corresponds to the $2Q'$ CS I state, where CS stands for the chiral stripe and Q' means the different amplitudes of $(m_{Q_1}^x)^2 + (m_{Q_1}^y)^2 + (m_{Q_1}^z)^2$ and $(m_{Q_2}^x)^2 + (m_{Q_2}^y)^2 + (m_{Q_2}^z)^2$. The spin configuration in this state is characterized by $(m_{Q_1}^y)^2$ and $(m_{Q_1}^z)^2$ as the dominant contribution and $(m_{Q_2}^x)^2$ as the subdominant contribution, as shown in Fig. 2(c). In other words, this spin configuration is represented by a superposi-

tion of the out-of-plane spiral wave at the $\mathbf{Q}_1$ component and the sinusoidal wave at the $\mathbf{Q}_2$ component. The orthonormal relation of the spin oscillations between the $\mathbf{Q}_1$ and $\mathbf{Q}_2$ components is due to preventing the energy loss under the spin length constraint with $|\mathbf{S}_i| = 1$ [55]. Reflecting the double- Q superposition in different spin components, the $2Q'$ CS I state exhibits the noncoplanar spin texture, as shown by the real-space spin configuration in Fig. 1(b), although they do not show a net scalar chirality in Fig. 2(b). Instead, this state accompanies the chirality density wave along the $\mathbf{Q}_2$ direction, as shown in Fig. 6(a) [56].

The emergence of the $2Q'$ CS I state at $I^{\text{EPA}} = 0$ is owing to the positive biquadratic interaction K rather than J' ; K tends to enhance the stability of multiple- Q states [52]. Indeed, this state remains stable when $J' = 0$, as will be discussed in Appendix A.

As H increases, M^x gradually increases and $(m_{\mathbf{Q}_2}^x)^2$ decreases, as shown in Figs. 2(a) and 2(c), respectively. Then, the $2Q'$ CS I state shows the phase transition to the $1Q$ OS state at $H \simeq 0.98$. The $1Q$ OS state corresponds to the single- Q out-of-plane spiral state, whose spiral plane lies in the yz plane in order to gain the Zeeman energy; it is noted that the spiral plane is always fixed in the yz plane irrespective of $\mathbf{Q}_{1,2}$ owing to the absence of the $\mathbf{Q}$ -dependent anisotropy. The real-space spin configuration of the $1Q$ OS state is shown in Fig. 1(c). The $1Q$ OS state turns into the fully polarized state at $H = 2$.

When the effect of I^{EPA} is considered, the $2Q'$ CS I state at zero field is replaced by the $2Q'$ CS II state, as shown in Fig. 1(a). We show the H dependence of physical quantities at $I^{\text{EPA}} = 0.06$ in Fig. 3. The spin texture of the $2Q'$ CS II state is characterized by $(m_{\mathbf{Q}_1}^x)^2$ and $(m_{\mathbf{Q}_1}^y)^2$ as the dominant contribution and $(m_{\mathbf{Q}_2}^z)^2$ as the subdominant contribution, as shown in Fig. 3(c). In contrast to the $2Q'$ CS I state, the spiral plane changes from the yz plane to the xy plane in order to gain the energy by I^{EPA} . Accordingly, the sinusoidal modulation occurs for the z spin component along the $\mathbf{Q}_2$ direction. The real-space spin and scalar chirality configurations are shown in Figs. 1(d) and 6(b), respectively, where the chirality density wave occurs along the $\mathbf{Q}_2$ direction similar to the $2Q'$ CS I state. It is noted that the uniform component of the scalar chirality is zero in the $2Q'$ CS II state [Fig. 3(b)].

The $2Q'$ CS II state changes into the $2Q'$ CS I state with jumps of M^x and $m_{\mathbf{Q}_2}^z$ at $H \simeq 0.23$, as shown in Figs. 3(a) and 3(c). This phase transition is regarded as the spin-flop phase transition, where the spiral plane changes from the xy plane parallel to the field direction to the yz plane perpendicular to the field direction.

While further increasing H , we find that the SkX is stabilized for $0.63 \lesssim H \lesssim 0.83$, which is characterized by nonzero scalar chirality χ^{sc} in Fig. 3(b). The spin configuration of the SkX is expressed as the linear combination of the spiral wave on the xy plane along the $\mathbf{Q}_1$ direction and that on the yz plane along the $\mathbf{Q}_2$ direction, as shown in Fig. 3(c), where the amplitudes of the magnetic moments at the $\mathbf{Q}_1$ and $\mathbf{Q}_2$ components are different owing to the lack of the fourfold rotational symmetry under the in-plane field; this feature is different from the SkX stabilized in the out-of-plane field, where the fourfold rotational




Figure 3: H dependence of (a) M^η , (b) χ^{sc} , and (c) $(m_{\mathbf{Q}_{1,2}}^\eta)^2$ for $\eta = x, y, z$ at $I^{\text{EPA}} = 0.06$. The vertical dashed line represents the phase boundary between different spin states; the corresponding phases are presented above in (a).

185 spin texture appears.

As shown by the real-space spin configuration in Fig. 1(e), the SkX phase consists of two vortices: One is the vortex with the positive z -spin polarization, which is denoted as the triangle, and the other is the antivortex with the negative z -spin polarization, which is denoted as the square. Since both the winding number and z -spin polarization are opposite between the vortex and antivortex, these vortices lead to the same sign of the scalar chirality, as shown in Fig. 6(c).




Figure 4: H dependence of (a) M^η , (b) χ^{sc} , and (c) $(m_{\mathbf{Q}}^\eta)^\eta$ for $\eta = x, y, z$ at $I^{\text{EPA}} = 0.12$. The vertical dashed line represents the phase boundary between different spin states; the corresponding phases are presented above in (a).

Although the SkX in Fig. 1(e) exhibits the positive scalar chirality, the state with the negative scalar chirality is degenerate; the degeneracy between them is lifted when the effect of bond-dependent magnetic anisotropy in $\mathbf{Q}_1$ - $\mathbf{Q}_4$ is taken into account [57]. The skyrmion core with $S_i^x = -1$ is found between the vortex and antivortex, which is located at the interstitial site rather than the lattice site.

From the energetic viewpoint, the SkX is stabilized by the interplay among




Figure 5: H dependence of (a) M^η , (b) χ^{sc} , and (c) $(m_{Q_{1,2}}^\eta)^2$ for $\eta = x, y, z$ at $I^{\text{EPA}} = 0.4$. The vertical dashed line represents the phase boundary between different spin states; the corresponding phases are presented above in (a).

200 K , J' , I^{EPA} , and H in the model Hamiltonian in Eq. (1). Among them, we find that J' , I^{EPA} , and H play an important role in inducing the SkX, while K enhances the stability of the SkX region. In other words, the SkX remains stable for $K = 0$, while it disappears for $J' = 0$, as shown in Appendix Appendix A. This means that the SkX can be realized in both itinerant magnets and insulating magnets, where the contribution from K is negligible for the latter.

205 Thus, the application of the in-plane external magnetic field to materials with easy-plane magnetic anisotropy is one of the possibilities to induce the SkX in




Figure 6: Real-space scalar chirality configurations in (a) the $2Q'$ CS I state at $I^{\text{EPA}} = 0$ and $H = 0.05$, (b) the $2Q'$ CS II state at $I^{\text{EPA}} = 0.06$ and $H = 0$, (c) the SkX at $I^{\text{EPA}} = 0.06$ and $H = 0.7$, and (d) the $2Q'$ I state at $I^{\text{EPA}} = 0.12$ and $H = 0.5$.

centrosymmetric systems irrespective of metals and insulators.

210 The SkX turns into the $2Q'$ CS I state again by increasing H . Then, the $2Q'$ CS I state changes into the $1Q$ OS state similar to the result at $I^{\text{EPA}} = 0$. The increase of H in the $1Q$ OS state leads to two additional phases before entering into the fully polarized state: One is the $2Q'$ II state and the other is the $2Q$ F state, where F means the fan. In the $2Q'$ II state, $(m_{Q_2}^y)^2$ becomes nonzero in addition to $(m_{Q_1}^y)^2$ and $(m_{Q_1}^z)^2$. Meanwhile, the $2Q$ F state is characterized

by the double- Q structure in the y component with the same intensity, i.e. $(m_{\mathbf{Q}_1}^y)^2 = (m_{\mathbf{Q}_2}^y)^2$, as shown in Fig. 3(c), where the spin configuration is shown in Fig. 1(f).

By increasing I^{EPA} , the regions of $2Q'$ CS I and $1Q$ OS phases become narrower, while those of the SkX, $2Q'$ CS II, and $2Q$ F phases become wider, as shown in Fig. 1(a). For $I^{\text{EPA}} \gtrsim 0.1$, the $2Q'$ I state appears between the SkX and $2Q'$ CS II state; the phase sequence in the case of $I^{\text{EPA}} = 0.12$ is shown in Fig. 4. The $2Q'$ I state is characterized by nonzero $(m_{\mathbf{Q}_1}^x)^2$, $(m_{\mathbf{Q}_1}^y)^2$, $(m_{\mathbf{Q}_1}^z)^2$, $(m_{\mathbf{Q}_2}^x)^2$, and $(m_{\mathbf{Q}_2}^z)^2$, as shown in Fig. 4(c); the spiral plane in the $\mathbf{Q}_1$ component lies between the xy and yz plane. Accordingly, this state shows nonzero uniform magnetization in the z component, M^z , as shown in Fig. 4(a). The real-space spin configuration of the $2Q'$ I state is shown in Fig. 1(g). Although this state accompanies the noncoplanar spin texture similar to the SkX, the scalar chirality becomes zero in the whole system in Fig. 4(b); only the chirality density wave is found, as shown in Fig. 6(d).

When I^{EPA} further increases, the SkX vanishes around $I^{\text{EPA}} \simeq 0.33$, and all the phases are characterized by the coplanar spin textures in the xy plane. In the case of $I^{\text{EPA}} = 0.4$, the $1Q$ IS state is stabilized in the low-field region instead of the $2Q'$ CS II state; the spin configuration in the former state is obtained by eliminating the $\mathbf{Q}_2$ component in the z spin in the latter state. Since I^{EPA} favors the in-plane spin texture, the phase transition from the $2Q'$ CS II state to the $1Q$ IS state in increasing I^{EPA} is reasonable. The spin configuration in real space is shown in Fig. 1(h).

The increase of H for $I^{\text{EPA}} = 0.4$ leads to the phase transition from the $1Q$ IS state to the $2Q'$ VC state with jumps of M^x and $m_{\mathbf{Q}_1}^y$, as shown in Figs. 5(a) and 5(c); VC means the vortex crystal. The $2Q'$ VC state is characterized by the spiral wave in the xy plane along the $\mathbf{Q}_1$ direction and the sinusoidal wave in the x component along the $\mathbf{Q}_2$ direction. Owing to the double- Q superposition in terms of the x -spin component, the real-space spin configuration consists of the vortex and antivortex, as shown in Fig. 1(i), which is similar to that in the SkX. Similar coplanar spin textures accompanying the vortex and antivortex have been found in triangular magnets with the easy-plane magnetic anisotropy [58], such as $\text{Y}_3\text{Co}_8\text{Sn}_4$ [59]. Our results indicate that the coplanar vortex crystal can be realized even in the centrosymmetric tetragonal host when the in-plane magnetic field is applied.

4. Summary

To summarize, we have investigated the stability of the SkX in centrosymmetric tetragonal magnets with easy-plane magnetic anisotropy by applying the in-plane external magnetic field. The data were obtained by performing the simulated annealing for the spin model including the biquadratic interaction, high-harmonic wave-vector interaction, and easy-plane two-spin anisotropic interaction. As a result, we have found that their synergy tends to favor the SkX in the in-plane field. The SkX is stabilized in the intermediate-field region when

the strength of easy-plane anisotropy is moderate. We have also found that the high-harmonic wave-vector interaction plays an important role in inducing the SkX, while the biquadratic interaction enhances its stability. Our results suggest the importance of the applied field direction; the in-plane magnetic field is essential to realize the SkX in centrosymmetric magnets with easy-plane magnetic anisotropy. Since the SkX in the out-of-plane magnetic field has been found in centrosymmetric tetragonal itinerant magnets, such as GdRu_2Si_2 with the easy-axis magnetic anisotropy, the Gd-based compounds are typical candidate materials. When the magnetic anisotropy is tuned from the easy axis to the easy plane by the chemical doping, the emergence of the SkX induced by the in-plane magnetic field is expected.

This research was supported by JSPS KAKENHI Grants Numbers JP21H01037, JP22H04468, JP22H00101, JP22H01183, JP23H04869, JP23K03288, and by JST PRESTO (JPMJPR20L8) and JST CREST (JPMJCR23O4). Parts of the numerical calculations were performed in the supercomputing systems in ISSP, the University of Tokyo.

Appendix A. Phase diagrams for other model parameters

We briefly discuss the magnetic phase diagrams when the effect of the biquadratic interaction K or high-harmonic wave-vector interaction J' is ignored. Figure A.7(a) shows the phase diagram in the case of $K = 0$. As shown in the phase diagram, the SkX is stabilized for intermediate I^{EPA} and H , although its stability region becomes narrower compared to the phase diagram at $K = 0.2$ in Fig. 1(a). This result indicates that the high-harmonic wave-vector interaction J' is significant in stabilizing the SkX in Fig. 1(a), but its stability region is narrow. Meanwhile, some of the other phases, $2Q'$ CS II, $2Q'$ II, and $2Q$ F, vanish in Fig. A.7(a), which means that solely J' is not enough to induce these phases.

Next, Fig. A.7(b) shows the phase diagram in the case of $J' = 0$. In contrast to the phase diagram in Fig. A.7(a), the SkX and $2Q'$ VC state do not appear in the phase diagram. In other words, the biquadratic interaction K is not enough to induce these two phases within the present model parameters.

References

- [1] A. Bogdanov, A. Hubert, Thermodynamically stable magnetic vortex states in magnetic crystals, *J. Magn. Magn. Mater.* 138 (3) (1994) 255 – 269. doi:[http://dx.doi.org/10.1016/0304-8853\(94\)90046-9](http://dx.doi.org/10.1016/0304-8853(94)90046-9).
- [2] A. Bogdanov, A. Hubert, The properties of isolated magnetic vortices, *Phys. Stat. Sol. (b)* 186 (2) (1994) 527–543.
- [3] A. Bogdanov, A. Hubert, The stability of vortex-like structures in uniaxial ferromagnets, *J. Magn. Magn. Mater.* 195 (1) (1999) 182–192.




Figure A.7: Magnetic phase diagrams by setting (a) $K = 0$ and (b) $J' = 0$. The other model parameters are the same as those used in Fig. 1(a).

- [4] N. Nagaosa, Y. Tokura, Topological properties and dynamics of magnetic skyrmions, Nat. Nanotechnol. 8 (12) (2013) 899–911. doi:10.1038/nnano.2013.243.
- [5] G. Finocchio, F. Büttner, R. Tomasello, M. Carpentieri, M. Kläui, Magnetic skyrmions: from fundamental to applications, J. Phys. D: Appl. Phys. 49 (42) (2016) 423001. doi:10.1088/0022-3727/49/42/423001.

- [6] A. Fert, N. Reyren, V. Cros, Magnetic skyrmions: advances in physics and potential applications, *Nat. Rev. Mater.* 2 (7) (2017) 17031. doi:10.1038/natrevmats.2017.31.
- 305 [7] X. Zhang, Y. Zhou, K. M. Song, T.-E. Park, J. Xia, M. Ezawa, X. Liu, W. Zhao, G. Zhao, S. Woo, Skyrmion-electronics: writing, deleting, reading and processing magnetic skyrmions toward spintronic applications, *J. Phys.: Condens. Matter* 32 (14) (2020) 143001. doi:10.1088/1361-648X/ab5488.
- 310 [8] S. Mühlbauer, B. Binz, F. Jonietz, C. Pfleiderer, A. Rosch, A. Neubauer, R. Georgii, P. Böni, Skyrmion lattice in a chiral magnet, *Science* 323 (5916) (2009) 915–919. doi:10.1126/science.1166767.
- [9] X. Z. Yu, Y. Onose, N. Kanazawa, J. H. Park, J. H. Han, Y. Matsui, N. Nagaosa, Y. Tokura, Real-space observation of a two-dimensional skyrmion crystal, *Nature* 465 (7300) (2010) 901–904. doi:10.1038/nature09124.
- 315 [10] X. Z. Yu, N. Kanazawa, Y. Onose, K. Kimoto, W. Zhang, S. Ishiwata, Y. Matsui, Y. Tokura, Near room-temperature formation of a skyrmion crystal in thin-films of the helimagnet FeGe, *Nat. Mater.* 10 (2) (2011) 106–109. doi:10.1038/nmat2916.
- 320 [11] S. Seki, X. Z. Yu, S. Ishiwata, Y. Tokura, Observation of skyrmions in a multiferroic material, *Science* 336 (6078) (2012) 198–201. doi:10.1126/science.1214143.
- [12] S. Heinze, K. von Bergmann, M. Menzel, J. Brede, A. Kubetzka, R. Wiesendanger, G. Bihlmayer, S. Blügel, Spontaneous atomic-scale magnetic skyrmion lattice in two dimensions, *Nat. Phys.* 7 (9) (2011) 713–718. doi:10.1038/nphys2045.
- 325 [13] T. Adams, A. Chacon, M. Wagner, A. Bauer, G. Brandl, B. Pedersen, H. Berger, P. Lemmens, C. Pfleiderer, Long-wavelength helimagnetic order and skyrmion lattice phase in Cu_2OSeO_3 , *Phys. Rev. Lett.* 108 (23) (2012) 237204. doi:10.1103/PhysRevLett.108.237204.
- 330 [14] Z. Hou, W. Ren, B. Ding, G. Xu, Y. Wang, B. Yang, Q. Zhang, Y. Zhang, E. Liu, F. Xu, et al., Observation of various and spontaneous magnetic skyrmionic bubbles at room temperature in a frustrated kagome magnet with uniaxial magnetic anisotropy, *Adv. Mater.* 29 (29) (2017) 1701144. doi:https://doi.org/10.1002/adma.201701144.
- 335 [15] T. Kurumaji, T. Nakajima, M. Hirschberger, A. Kikkawa, Y. Yamasaki, H. Sagayama, H. Nakao, Y. Taguchi, T.-h. Arima, Y. Tokura, Skyrmion lattice with a giant topological Hall effect in a frustrated triangular-lattice magnet, *Science* 365 (6456) (2019) 914–918. doi:10.1126/science.aau0968.
- 340

- [16] N. D. Khanh, T. Nakajima, X. Yu, S. Gao, K. Shibata, M. Hirschberger, Y. Yamasaki, H. Sagayama, H. Nakao, L. Peng, K. Nakajima, R. Takagi, T.-h. Arima, Y. Tokura, S. Seki, Nanometric square skyrmion lattice in a centrosymmetric tetragonal magnet, *Nat. Nanotechnol.* 15 (2020) 444. doi:10.1038/s41565-020-0684-7.
- [17] Y. Tokura, N. Kanazawa, Magnetic skyrmion materials, *Chem. Rev.* 121 (2021) 2857. doi:10.1021/acs.chemrev.0c00297.
- [18] I. Dzyaloshinsky, A thermodynamic theory of “weak” ferromagnetism of antiferromagnetics, *J. Phys. Chem. Solids* 4 (4) (1958) 241–255.
- [19] T. Moriya, Anisotropic superexchange interaction and weak ferromagnetism, *Phys. Rev.* 120 (1) (1960) 91. doi:https://doi.org/10.1103/PhysRev.120.91.
- [20] U. K. Rößler, A. N. Bogdanov, C. Pfleiderer, Spontaneous skyrmion ground states in magnetic metals, *Nature* 442 (7104) (2006) 797–801. doi:10.1038/nature05056.
- [21] B. Binz, A. Vishwanath, V. Aji, Theory of the helical spin crystal: A candidate for the partially ordered state of MnSi, *Phys. Rev. Lett.* 96 (2006) 207202. doi:10.1103/PhysRevLett.96.207202.
- [22] B. Binz, A. Vishwanath, Theory of helical spin crystals: Phases, textures, and properties, *Phys. Rev. B* 74 (2006) 214408. doi:10.1103/PhysRevB.74.214408.
- [23] S. D. Yi, S. Onoda, N. Nagaosa, J. H. Han, Skyrmions and anomalous Hall effect in a Dzyaloshinskii-Moriya spiral magnet, *Phys. Rev. B* 80 (2009) 054416. doi:10.1103/PhysRevB.80.054416.
- [24] A. B. Butenko, A. A. Leonov, U. K. Rößler, A. N. Bogdanov, Stabilization of skyrmion textures by uniaxial distortions in noncentrosymmetric cubic helimagnets, *Phys. Rev. B* 82 (2010) 052403. doi:10.1103/PhysRevB.82.052403.
- [25] T. Okubo, S. Chung, H. Kawamura, Multiple- q states and the skyrmion lattice of the triangular-lattice Heisenberg antiferromagnet under magnetic fields, *Phys. Rev. Lett.* 108 (2012) 017206. doi:10.1103/PhysRevLett.108.017206.
- [26] A. O. Leonov, M. Mostovoy, Multiply periodic states and isolated skyrmions in an anisotropic frustrated magnet, *Nat. Commun.* 6 (2015) 8275. doi:10.1038/ncomms9275.
- [27] J. Liang, J. Yu, J. Chen, M. Qin, M. Zeng, X. Lu, X. Gao, J.-M. Liu, Magnetic field gradient driven dynamics of isolated skyrmions and anti-skyrmions in frustrated magnets, *New J. of Phys.* 20 (5) (2018) 053037. doi:10.1088/1367-2630/aac24c.

- [28] R. Yambe, S. Hayami, Effective spin model in momentum space: Toward a systematic understanding of multiple- Q instability by momentum-resolved anisotropic exchange interactions, *Phys. Rev. B* 106 (2022) 174437. doi:10.1103/PhysRevB.106.174437.
- [29] S. Hayami, R. Yambe, Field-direction sensitive skyrmion crystals in cubic chiral systems: Implication to $4f$ -electron compound EuPtSi, *J. Phys. Soc. Jpn.* 90 (7) (2021) 073705. doi:10.7566/JPSJ.90.073705.
- [30] B. Göbel, A. Mook, J. Henk, I. Mertig, O. A. Tretiakov, Magnetic bimerons as skyrmion analogues in in-plane magnets, *Phys. Rev. B* 99 (2019) 060407(R). doi:10.1103/PhysRevB.99.060407.
- [31] K.-W. Moon, J. Yoon, C. Kim, C. Hwang, Existence of in-plane magnetic skyrmion and its motion under current flow, *Phys. Rev. Appl.* 12 (2019) 064054. doi:10.1103/PhysRevApplied.12.064054.
- [32] L. Flacke, V. Ahrens, S. Mendisch, L. Körber, T. Böttcher, E. Meidinger, M. Yaqoob, M. Müller, L. Liensberger, A. Kákay, M. Becherer, P. Pirro, M. Althammer, S. Geprägs, H. Huebl, R. Gross, M. Weiler, Robust formation of nanoscale magnetic skyrmions in easy-plane anisotropy thin film multilayers with low damping, *Phys. Rev. B* 104 (2021) L100417. doi:10.1103/PhysRevB.104.L100417.
- [33] S. Hayami, In-plane magnetic field-induced skyrmion crystal in frustrated magnets with easy-plane anisotropy, *Phys. Rev. B* 103 (2021) 224418. doi:10.1103/PhysRevB.103.224418.
- [34] S. Hayami, Y. Motome, Noncoplanar multiple- Q spin textures by itinerant frustration: Effects of single-ion anisotropy and bond-dependent anisotropy, *Phys. Rev. B* 103 (2021) 054422. doi:10.1103/PhysRevB.103.054422.
- [35] S. Hayami, Y. Motome, Square skyrmion crystal in centrosymmetric itinerant magnets, *Phys. Rev. B* 103 (2021) 024439. doi:10.1103/PhysRevB.103.024439.
- [36] O. I. Utesov, Thermodynamically stable skyrmion lattice in a tetragonal frustrated antiferromagnet with dipolar interaction, *Phys. Rev. B* 103 (2021) 064414. doi:10.1103/PhysRevB.103.064414.
- [37] Z. Wang, Y. Su, S.-Z. Lin, C. D. Batista, Meron, skyrmion, and vortex crystals in centrosymmetric tetragonal magnets, *Phys. Rev. B* 103 (2021) 104408. doi:10.1103/PhysRevB.103.104408.
- [38] S. Hayami, Rectangular and square skyrmion crystals on a centrosymmetric square lattice with easy-axis anisotropy, *Phys. Rev. B* 105 (2022) 174437. doi:10.1103/PhysRevB.105.174437.

- [39] N. D. Khanh, T. Nakajima, S. Hayami, S. Gao, Y. Yamasaki, H. Sagayama, H. Nakao, R. Takagi, Y. Motome, Y. Tokura, T.-h. Arima, S. Seki, Zool-
420 ogy of Multiple- Q Spin Textures in a Centrosymmetric Tetragonal Magnet with Itinerant Electrons, *Adv. Sci.* 9 (2022) 2105452. doi:10.1002/advs.202105452.
- [40] N. Matsuyama, T. Nomura, S. Imajo, T. Nomoto, R. Arita, K. Sudo, M. Kimata, N. D. Khanh, R. Takagi, Y. Tokura, S. Seki, K. Kindo, Y. Ko-
425 hama, Quantum oscillations in the centrosymmetric skyrmion-hosting magnet GdRu_2Si_2 , *Phys. Rev. B* 107 (2023) 104421. doi:10.1103/PhysRevB.107.104421.
- [41] S. Hayami, Y. Kato, Widely-sweeping magnetic field–temperature phase diagrams for skyrmion-hosting centrosymmetric tetragonal magnets, *J. Magn. Magn. Mater.* 571 (2023) 170547. doi:https://doi.org/10.1016/j.jmmm.2023.170547.
430
- [42] G. D. A. Wood, D. D. Khalyavin, D. A. Mayoh, J. Bouaziz, A. E. Hall, S. J. R. Holt, F. Orlandi, P. Manuel, S. Blügel, J. B. Staunton, O. A. Petrenko, M. R. Lees, G. Balakrishnan, Double- q ground state with topological charge stripes in the centrosymmetric skyrmion candidate GdRu_2Si_2 ,
435 *Phys. Rev. B* 107 (2023) L180402. doi:10.1103/PhysRevB.107.L180402.
- [43] K. Kaneko, T. Kawasaki, A. Nakamura, K. Munakata, A. Nakao, T. Hanashima, R. Kiyonagi, T. Ohhara, M. Hedo, T. Nakama, Y. Onuki, Charge-density-wave order and multiple magnetic transitions in divalent europium compound EuAl_4 , *J. Phys. Soc. Jpn.* 90 (6) (2021) 064704.
440 doi:10.7566/JPSJ.90.064704.
- [44] T. Shang, Y. Xu, D. J. Gawryluk, J. Z. Ma, T. Shiroka, M. Shi, E. Pomjakushina, Anomalous Hall resistivity and possible topological Hall effect in the EuAl_4 antiferromagnet, *Phys. Rev. B* 103 (2021) L020405.
445 doi:10.1103/PhysRevB.103.L020405.
- [45] R. Takagi, N. Matsuyama, V. Ukleev, L. Yu, J. S. White, S. Francoual, J. R. L. Mardegan, S. Hayami, H. Saito, K. Kaneko, K. Ohishi, Y. Onuki, T.-h. Arima, Y. Tokura, T. Nakajima, S. Seki, Square and rhombic lattices of magnetic skyrmions in a centrosymmetric binary compound, *Nat. Commun.* 13 (2022) 1472. doi:10.1038/s41467-022-29131-9.
450
- [46] X. Y. Zhu, H. Zhang, D. J. Gawryluk, Z. X. Zhen, B. C. Yu, S. L. Ju, W. Xie, D. M. Jiang, W. J. Cheng, Y. Xu, M. Shi, E. Pomjakushina, Q. F. Zhan, T. Shiroka, T. Shang, Spin order and fluctuations in the EuAl_4 and EuGa_4 topological antiferromagnets: A μSR study, *Phys. Rev. B* 105 (2022) 014423. doi:10.1103/PhysRevB.105.014423.
455
- [47] W. R. Meier, J. R. Torres, R. P. Hermann, J. Zhao, B. Lavina, B. C. Sales, A. F. May, Thermodynamic insights into the intricate magnetic phase diagram of EuAl_4 , *Phys. Rev. B* 106 (2022) 094421. doi:10.1103/PhysRevB.106.094421.

- [48] M. Gen, R. Takagi, Y. Watanabe, S. Kitou, H. Sagayama, N. Matsuyama, Y. Kohama, A. Ikeda, Y. Ōnuki, T. Kurumaji, T.-h. Arima, S. Seki, Rhombic skyrmion lattice coupled with orthorhombic structural distortion in EuAl_4 , Phys. Rev. B 107 (2023) L020410. doi:10.1103/PhysRevB.107.L020410.
- [49] R. Yang, C.-C. Le, P. Zhu, Z.-W. Wang, T. Shang, Y.-M. Dai, J.-P. Hu, M. Dressel, Charge density wave transition in the magnetic topological semimetal EuAl_4 , Phys. Rev. B 109 (2024) L041113. doi:10.1103/PhysRevB.109.L041113.
- [50] Y. Akagi, M. Udagawa, Y. Motome, Hidden multiple-spin interactions as an origin of spin scalar chiral order in frustrated Kondo lattice models, Phys. Rev. Lett. 108 (2012) 096401. doi:10.1103/PhysRevLett.108.096401.
- [51] S. Hayami, Y. Motome, Multiple- Q instability by $(d-2)$ -dimensional connections of Fermi surfaces, Phys. Rev. B 90 (2014) 060402(R). doi:10.1103/PhysRevB.90.060402.
- [52] S. Hayami, R. Ozawa, Y. Motome, Effective bilinear-biquadratic model for noncoplanar ordering in itinerant magnets, Phys. Rev. B 95 (2017) 224424. doi:10.1103/PhysRevB.95.224424.
- [53] S. Hayami, Y. Motome, Néel- and Bloch-type magnetic vortices in Rashba metals, Phys. Rev. Lett. 121 (2018) 137202. doi:10.1103/PhysRevLett.121.137202.
- [54] S. Hayami, Multiple skyrmion crystal phases by itinerant frustration in centrosymmetric tetragonal magnets, J. Phys. Soc. Jpn. 91 (2) (2022) 023705. doi:10.7566/JPSJ.91.023705.
- [55] D. Solenov, D. Mozyrsky, I. Martin, Chirality waves in two-dimensional magnets, Phys. Rev. Lett. 108 (2012) 096403. doi:10.1103/PhysRevLett.108.096403.
- [56] S. Hayami, S.-Z. Lin, Y. Kamiya, C. D. Batista, Vortices, skyrmions, and chirality waves in frustrated Mott insulators with a quenched periodic array of impurities, Phys. Rev. B 94 (2016) 174420. doi:10.1103/PhysRevB.94.174420.
- [57] S. Hayami, R. Yambe, Degeneracy lifting of Néel, Bloch, and anti-skyrmion crystals in centrosymmetric tetragonal systems, J. Phys. Soc. Jpn. 89 (10) (2020) 103702. doi:10.7566/JPSJ.89.103702.
- [58] S. Hayami, Multiple- Q magnetism by anisotropic bilinear-biquadratic interactions in momentum space, J. Magn. Magn. Mater. 513 (2020) 167181. doi:10.1016/j.jmmm.2020.167181.

- [59] R. Takagi, J. White, S. Hayami, R. Arita, D. Honecker, H. Rønnow, Y. Tokura, S. Seki, Multiple- q noncollinear magnetism in an itinerant hexagonal magnet, *Sci. Adv.* 4 (11) (2018) eaau3402. doi:10.1126/sciadv.aau3402.

500