



HOKKAIDO UNIVERSITY

Title	Phenotypic plasticity drives the development of laterality in the scale-eating cichlid fish <i>Perissodus microlepis</i>
Author(s)	Marubayashi, Nanako; Yasugi, Masaki; Takeuchi, Yuichi
Citation	Evolution, 79(10), 2035–2046 https://doi.org/10.1093/evolut/qpaf131
Issue Date	2025-07-04
Doc URL	https://hdl.handle.net/2115/97969
Rights	This is the peer reviewed version of the following article: Nanako Marubayashi, Masaki Yasugi, Yuichi Takeuchi, Phenotypic plasticity drives the development of laterality in the scale-eating cichlid fish <i>Perissodus microlepis</i> , Evolution, 2025;, qpaf131, https://doi.org/10.1093/evolut/qpaf131 , which has been published in final form at DOI:10.1093/evolut/qpaf131. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.
Type	journal article
File Information	Phenotypic_plasticity_250605_Title.pdf



Phenotypic plasticity drives the development of laterality in the scale-eating cichlid fish *Perissodus microlepis*

Nanako Marubayashi ¹, Masaki Yasugi ², Yuichi Takeuchi ^{3*}

¹Department of Anatomy and Neuroscience, Faculty of Medicine, University of Toyama, Toyama, Japan.

²Faculty of Marine Science and Technology, Fukui Prefectural University, Fukui, Japan.

³Graduate School of Life Science, Hokkaido University, Sapporo, Japan.

***Corresponding author:** Yuichi Takeuchi

Email: ytake@sci.hokudai.ac.jp

Data availability

The data and accompanying description are available from: [http://10.5061/dryad.q83bk3jv8](http://dx.doi.org/10.5061/dryad.q83bk3jv8). Any additional information required to reanalyse the data reported in this paper is available from the corresponding author upon request.

Author contributions

N.M. and Y.T. conceived the ideas; N.M. and Y.T. conducted behavioural experiments; N.M., M.Y. and Y.T. performed the morphological analysis; M.Y. supervised the behavioural analysis using image-processing and deep-learning methods; N.M., M.Y. and Y.T. led the writing of the manuscript. All authors gave final approval for publication and agreed to be held accountable for the work performed.

Funding

This study was supported by the JSPS KAKENHI programme (Y.T. – JP23K05960, JP23KK0131, and JP24H01418; M.Y. – JP18KK0208, JP20H05702, and JP23KK0131) and by grants from the Mitsubishi Foundation (Y.T. – 202210033), the Ohsumi Frontier Science Foundation, Toray Science and Technology Grant, the Asahi Glass Foundation (Y.T.), and the Chubu Branch of the Ecological Society of Japan (N.M.).

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

We are grateful to the staff at World Freshwater Aquarium Aquatotto Gifu for arranging access to the fish. We thank M. Matsuyama, Y. Higuchi, M. Nakagawa, M. Imai, and C. Tsukada (University of Toyama) and T. Taga, Y. Shibata, H. Goto, N. Yamashima, T. Sato, K. Koike, H. Nagasaka, and A. Inuta (Hokkaido University) for their help with fish

36 care. We also thank Prof. Hiroki Hata (Ehime University) for supporting the geographic

37 morphometric work.

38