



HOKKAIDO UNIVERSITY

Title	On the spectral synthesis for the unit circle in $FLsq(R_2)$
Author(s)	Kobayashi, Masaharu; Sato, Enji
Citation	Journal of Pseudo-Differential Operators and Applications, 16(3) https://doi.org/10.1007/s11868-025-00728-0
Issue Date	2025-08-20
Doc URL	https://hdl.handle.net/2115/98041
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: http://dx.doi.org/10.1007/s11868-025-00728-0
Type	journal article
File Information	KoSa4 (2).pdf



On the spectral synthesis for the unit circle in $\mathcal{FL}_s^q(\mathbf{R}^2)$

Masaharu Kobayashi^{1*} and Enji Sato^{2†}

^{1*}Department of Mathematics, Hokkaido University, Kita 10, Nishi 8,
Kita-Ku, Sapporo, 060-0810, Hokkaido, Japan.

²Faculty of Science, Yamagata University, Kojirakawa 1-4-12,
Yamagata-City, 990-8560, Yamagata, Japan.

*Corresponding author(s). E-mail(s):

m-kobayashi@math.sci.hokudai.ac.jp;

Contributing authors: esato@sci.kj.yamagata-u.ac.jp;

[†]These authors contributed equally to this work.

Abstract

Let $\mathcal{FL}_s^q(\mathbf{R}^2)$ denote the set of all tempered distributions $f \in \mathcal{S}'(\mathbf{R}^2)$ such that the norm $\|f\|_{\mathcal{FL}_s^q} := (\int_{\mathbf{R}^2} |\mathcal{F}[f](\xi)| (1 + |\xi|)^s)^q d\xi)^{1/q}$ is finite, where $\mathcal{F}[f]$ denotes the Fourier transform of f . We investigate the spectral synthesis for the unit circle $S^1 \subset \mathbf{R}^2$ in $\mathcal{FL}_s^q(\mathbf{R}^2)$ with $1 < q < \infty$ and $s > 2/q'$, where q' denotes the conjugate exponent of q .

Keywords: Fourier-Wermer algebra, Fourier-Lebesgue spaces, Banach algebra, Set of spectral synthesis

MSC Classification: 42B35 , 43A45 , 43A15

1 Introduction

In commutative harmonic analysis, there is an important problem of determining whether the spectral synthesis holds for a given Banach space B and a given subset E of $\mathbf{R}^n$. To be more precise, let $(B, \|\cdot\|_B)$ be a Banach space of bounded continuous, complex-valued functions, i.e., $B \subset C_b(\mathbf{R}^n)$, and assume that $C_c^\infty(\mathbf{R}^n)$ is dense in B and convergence in B implies pointwise convergence. For every closed subset E of $\mathbf{R}^n$,

we set $I(E) := \{f \in B \mid f|_E = 0\}$ and define $J(E)$ as the closure in B of the set

$$\{f \in C_c^\infty(\mathbf{R}^n) \mid f = 0 \text{ in a neighborhood of } E\}.$$

Here, $f|_E = 0$ means that $f(x) = 0$ for all $x \in E$. Since $I(E) \supset J(E)$, it is natural to ask whether $I(E) \subset J(E)$ holds. If $I(E) = J(E)$, then E is called a *set of spectral synthesis* in B . Let $A(\mathbf{R}^n)$ be the Fourier algebra of all functions on $\mathbf{R}^n$ which are the Fourier transform of functions in $L^1(\mathbf{R}^n)$, also denoted by $\mathcal{FL}_0^1(\mathbf{R}^n)$ elsewhere. It is well known that $A(\mathbf{R}^n) \subset C_0(\mathbf{R}^n)$ by the Riemann-Lebesgue lemma and $A(\mathbf{R}^n)$ is a Banach algebra with respect to pointwise multiplication and the norm $\|f\|_A := \|\mathcal{F}[f]\|_{L^1}$, where $\mathcal{F}[f](\xi) := \int_{\mathbf{R}^n} f(x)e^{-ix\xi}dx$. The first counterexample of the spectral synthesis was given by Schwartz [25] and he showed that the unit sphere $S^{n-1} := \{x \in \mathbf{R}^n \mid |x| = 1\}$ is not a set of spectral synthesis in $A(\mathbf{R}^n)$, if $n \geq 3$. Surprisingly, Herz [17] showed that the spectral synthesis holds for S^1 in $A(\mathbf{R}^2)$. Motivated by these results, Varopoulos [29] extended Herz's method to higher dimensions to study this gap and showed that S^{n-1} ($n \geq 3$) is of weak synthesis, i.e., the closure in $A(\mathbf{R}^n)$ of the set $\{f \in C_c^\infty(\mathbf{R}^n) \mid f|_E = 0\}$ is equal to $I(S^{n-1})$. On the other hand, Reiter [22] showed similar results in the Fourier-Beurling algebras $\mathcal{FL}_s^1(\mathbf{R}^n)$ and proved that S^1 is a set of spectral synthesis in $\mathcal{FL}_s^1(\mathbf{R}^2)$ if $0 < s < 1/2$, and S^1 is not a set of spectral synthesis in $\mathcal{FL}_s^1(\mathbf{R}^2)$ if $s \geq 1/2$. The study of spectral synthesis is also related to a uniqueness property for PDEs (see for example [16] and Section 6 below). The reader is referred to [5], [6], [7], [23], [24], [27] for more details on the spectral synthesis.

In this paper, we investigate the spectral synthesis for S^1 in the Fourier-Wermer algebra $\mathcal{FL}_s^q(\mathbf{R}^2)$ with $1 < q < \infty$ and $s > 2/q'$, which gives an answer to the problem proposed by Domar [6]. The precise definition of $\mathcal{FL}_s^q(\mathbf{R}^n)$ will be given in Section 2.2, but we remark that $\mathcal{FL}_s^q(\mathbf{R}^n) \subset C_0(\mathbf{R}^n)$ and $\mathcal{FL}_s^q(\mathbf{R}^n)$ is a Banach algebra with respect to pointwise multiplication if $1 < q < \infty$ and $s > n/q'$. "Spectral synthesis in $\mathcal{FL}_s^q$ " is a classical topic, but recently it has been found to be useful in determining "Spectral synthesis in modulation spaces $M_s^{p,q}$ " (see [11], [12]). Our results are as follows:

Theorem 1 *Let $1 < q < \infty$ and $s > 2/q' + 1/2$. Then S^1 is not a set of spectral synthesis in $\mathcal{FL}_s^q(\mathbf{R}^2)$.*

Theorem 2 *Let $1 < q < 2$ and $2/q' < s < 2/q' + 1/2$. Then S^1 is a set of spectral synthesis in $\mathcal{FL}_s^q(\mathbf{R}^2)$.*

Moreover, we have the following result for $n \geq 3$.

Theorem 3 *Let $1 < q < \infty$ and $s > n/q'$. Then S^{n-1} ($n \geq 3$) is not a set of spectral synthesis in $\mathcal{FL}_s^q(\mathbf{R}^n)$.*

We explain the organization of this paper. After the next preliminary section devoted to the definitions and basic properties of $\mathcal{FL}_s^q(\mathbf{R}^n)$ and a set of spectral synthesis, we prove Theorem 1 in Section 3. Theorems 2 and 3 are treated in Sections 4 and 5, respectively. Finally, we give an application of spectral synthesis to PDEs and some remarks on closed ideals in $\mathcal{FL}_s^q(\mathbf{R}^2)$ in Sections 6 and 7.

2 Preliminaries

The following notation will be used throughout this paper. The notation $I \lesssim J$ means that I is bounded by a constant times J and we write $I \approx J$ if $I \lesssim J$ and $J \lesssim I$. We use C to denote various positive constants which may change from line to line. The closed ball with center $x_0 \in \mathbf{R}^n$ and radius $R > 0$ is defined by $B_n(x_0, R) := \{x \in \mathbf{R}^n \mid |x - x_0| \leq R\}$ and its Lebesgue measure is denoted by $|B_n(x_0, R)|$. For $x \in \mathbf{R}^n$ and $E \subset \mathbf{R}^n$, we set $d(x, E) := \inf\{|x - y| \mid y \in E\}$. We define for $1 \leq q < \infty$ and $s \in \mathbf{R}$

$$\|f\|_{L_s^q} := \left(\int_{\mathbf{R}^n} (|f(x)| \langle x \rangle^s)^q dx \right)^{1/q}$$

and $\|f\|_{L_s^\infty} := \text{ess. sup}_{x \in \mathbf{R}^n} \langle x \rangle^s |f(x)|$ with $\langle x \rangle := 1 + |x|$. We simply write $L^q(\mathbf{R}^n)$ instead of $L_0^q(\mathbf{R}^n)$. For $1 \leq q < \infty$, q' denotes the conjugate exponent of q , i.e., $1/q + 1/q' = 1$. Let $C_c(\mathbf{R}^n)$ be the set of all $f \in C(\mathbf{R}^n)$ such that $\text{supp } f$ is compact. Also let $C_c^\infty(\mathbf{R}^n)$ be the set of all $f \in C^\infty(\mathbf{R}^n)$ such that $\text{supp } f$ is compact. If $1 \leq q < \infty$ and $s \in \mathbf{R}$, then $C_c^\infty(\mathbf{R}^n)$ is dense in $L_s^q(\mathbf{R}^n)$. As in the proof of [23, Proposition 1.6.11], we can easily see

$$\lim_{|t| \rightarrow 0} \|f(\cdot - t) - f(\cdot)\|_{L_s^q} = 0 \quad (1)$$

for $1 \leq q < \infty$ and $f \in L_s^q(\mathbf{R}^n)$ (see [8], [15], [26] for more details on the weighted function spaces). Let $\mathcal{S}(\mathbf{R}^n)$ be the Schwartz space of all complex-valued rapidly decreasing infinitely differentiable functions on $\mathbf{R}^n$ and $\mathcal{S}'(\mathbf{R}^n)$ be the set of all tempered distributions, i.e., the topological dual of $\mathcal{S}(\mathbf{R}^n)$. We use $\langle f, g \rangle$ to denote the extension of the inner product $\langle f, g \rangle_{L^2(\mathbf{R}^n)} = \int_{\mathbf{R}^n} f(t) \overline{g(t)} dt$ on $L^2(\mathbf{R}^n)$ to $\mathcal{FL}_s^q(\mathbf{R}^n) \times \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$ (see Section 2.2 below). The Fourier transform of $f \in L^1(\mathbf{R}^n)$ is $\mathcal{F}[f](\xi) := \int_{\mathbf{R}^n} f(x) e^{-ix\xi} dx$, and the inverse Fourier transform of $h \in L^1(\mathbf{R}^n)$ is $\mathcal{F}^{-1}[h](x) := (2\pi)^{-n} \mathcal{F}[h](-x)$. We note that $\langle f, g \rangle_{L^2(\mathbf{R}^n)} = (2\pi)^{-n} \langle \mathcal{F}[f], \mathcal{F}[g] \rangle_{L^2(\mathbf{R}^n)}$, $\mathcal{F}[f * g] = \mathcal{F}[f] \cdot \mathcal{F}[g]$ and $\mathcal{F}[fg] = (2\pi)^{-n} (\mathcal{F}[f] * \mathcal{F}[g])$, where the convolution $f * g$ is defined by $(f * g)(x) = \int_{\mathbf{R}^n} f(x - y) g(y) dy$. Moreover, χ_E denotes the characteristic function of a set $E \subset \mathbf{R}^n$, i.e., $\chi_E(x) = 1$ (if $x \in E$), $= 0$ (if $x \notin E$).

2.1 The Bessel functions

Let $k \in \mathbf{Z}$. Then the *Bessel function* of the first kind of order k is defined by

$$J_k(t) := \frac{1}{2\pi} \int_0^{2\pi} e^{it \sin \theta} e^{-ik\theta} d\theta \quad (t > 0). \quad (2)$$

We also note

$$J_0(t) = \frac{1}{2\pi} \int_0^{2\pi} e^{it \cos \theta} d\theta = \frac{1}{\pi} \int_0^\pi e^{it \cos \theta} d\theta.$$

The Bessel function has the following interesting properties (see for instance [13, Appendix B], [28, p.153, p.158], [30, p.18]).

Lemma 4 We have the following.

- (i) $J_k(t) = O(t^{-1/2})$ as $t \rightarrow \infty$ ($k = 0, 1, 2, \dots$).
- (ii) $J_k(t) = (-1)^k J_{-k}(t)$ ($k \in \mathbf{Z}$).
- (iii) $\frac{d}{dt} \{t^{-k} J_k(t)\} = -t^{-k} J_{k+1}(t)$ ($k = 0, 1, 2, \dots$).
- (iv) $\frac{d}{dt} \{t^k J_k(t)\} = t^k J_{k-1}(t)$ ($k = 1, 2, \dots$).

2.2 The Fourier-Wermer algebra

Let $1 \leq q < \infty$ and $s \in \mathbf{R}$. Then $\mathcal{FL}_s^q(\mathbf{R}^n)$ is the set of all $f \in \mathcal{S}'(\mathbf{R}^n)$ such that the norm

$$\|f\|_{\mathcal{FL}_s^q} := \left(\int_{\mathbf{R}^n} (|\mathcal{F}[f](\xi)| \langle \xi \rangle^s)^q d\xi \right)^{1/q}$$

is finite. It is well known that $(\mathcal{FL}_s^q(\mathbf{R}^n), \|\cdot\|_{\mathcal{FL}_s^q})$ is a Banach space and its dual space is $\mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$, i.e., $\ell \in (\mathcal{FL}_s^q(\mathbf{R}^n))'$ if and only if

$$\ell(f) = \frac{1}{(2\pi)^n} \int_{\mathbf{R}^n} \mathcal{F}[f](\xi) \overline{\mathcal{F}[\phi](\xi)} d\xi =: \langle f, \phi \rangle \quad (f \in \mathcal{FL}_s^q(\mathbf{R}^n))$$

for some $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$ with $\|\ell\|_{op} = (2\pi)^{-n} \|\phi\|_{\mathcal{FL}_{-s}^{q'}}$. We note that if $1 < q < \infty$, $s > n/q'$ and $f \in \mathcal{FL}_s^q(\mathbf{R}^n)$, then $\mathcal{F}[f] \in L^1(\mathbf{R}^n)$ by the Hölder inequality, and thus f is continuous and vanishes at infinity, i.e., $f \in C_0(\mathbf{R}^n)$, by the Riemann-Lebesgue lemma. Thus the inversion formula $f(x) = \mathcal{F}^{-1}[\mathcal{F}[f]](x)$ holds for all $x \in \mathbf{R}^n$. Moreover since $\mathcal{F}[f](\xi) = \overline{\mathcal{F}[f](-\xi)}$, we have $\bar{f} \in \mathcal{FL}_s^q(\mathbf{R}^n)$ whenever $f \in \mathcal{FL}_s^q(\mathbf{R}^n)$. Therefore for $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$, we can define $\bar{\phi} \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$ by

$$\langle f, \bar{\phi} \rangle := \overline{\langle \bar{f}, \phi \rangle} \quad (f \in \mathcal{FL}_s^q(\mathbf{R}^n))$$

and obtain

$$\langle f, \bar{\phi} \rangle = (2\pi)^{-n} \int_{\mathbf{R}^n} \mathcal{F}[f](-\xi) \mathcal{F}[\phi](\xi) d\xi = (2\pi)^{-n} (\mathcal{F}[f] * \mathcal{F}[\phi])(0).$$

We also note that if $1 < q < \infty$, then $\mathcal{FL}_s^q(\mathbf{R}^n)$ is a multiplication algebra if and only if $s > n/q'$ (see for example [3], [19, Proposition B.2]). In some literature, $\mathcal{FL}_s^q(\mathbf{R}^n)$ are called the *Fourier-Lebesgue spaces*, but following a suggestion of Feichtinger [10], we call $\mathcal{FL}_s^q(\mathbf{R}^n)$ the *Fourier-Wermer algebra* because they are the Fourier images of convolution algebras $L_s^q(\mathbf{R}^n)$ that were studied in the early paper of Wermer [31] (see also [5]).

In the following, we often use the following lemma.

Lemma 5 Let $1 < q < \infty$ and $s > n/q'$. Then for $f \in \mathcal{F}L_s^q(\mathbf{R}^n)$ and $\Phi \in L_{-s}^{q'}(\mathbf{R}^n)$, we have the following:

- (i) $\mathcal{F}[f] * \Phi$ is a continuous function on $\mathbf{R}^n$.
- (ii) $\mathcal{F}[f] * \Phi \in L_{-s}^{q'}(\mathbf{R}^n)$.

Proof (i) Let $\xi_0 \in \mathbf{R}^n$ and $\varepsilon > 0$. Since $\mathcal{F}[f] \in L_s^q(\mathbf{R}^n)$ and $C_c^\infty(\mathbf{R}^n)$ is dense in $L_s^q(\mathbf{R}^n)$, we can find $\Psi \in C_c^\infty(\mathbf{R}^n)$ with $\|\mathcal{F}[f] - \Psi\|_{L_s^q} < \varepsilon$ and $\text{supp } \Psi \subset B_n(0, R)$ for some $R > 0$. Moreover, since Ψ is uniformly continuous on $\mathbf{R}^n$, there exists $0 < \delta < 1$ such that

$$\sup_{|\eta_1 - \eta_2| < \delta} |\Psi(\eta_1) - \Psi(\eta_2)| < \varepsilon(1 + |\xi_0| + R)^{-(s+n/q)}. \quad (3)$$

Now we show $\mathcal{F}[f] * \Phi$ is continuous at ξ_0 . Suppose $|\xi_0 - \xi| < \delta$. Then the Hölder inequality implies

$$\begin{aligned} |(\mathcal{F}[f] * \Phi)(\xi_0) - (\mathcal{F}[f] * \Phi)(\xi)| &= \left| \int_{\mathbf{R}^n} (\mathcal{F}[f](\xi_0 - \eta) - \mathcal{F}[f](\xi - \eta)) \Phi(\eta) d\eta \right| \\ &\leq \|\Phi\|_{L_{-s}^{q'}} \|\mathcal{F}[f](\xi_0 - \cdot) - \mathcal{F}[f](\xi - \cdot)\|_{L_s^q}. \end{aligned}$$

On the other hand, we note that

$$\begin{aligned} \|\mathcal{F}[f](\xi_0 - \cdot) - \Psi(\xi_0 - \cdot)\|_{L_s^q} &= \left(\int_{\mathbf{R}^n} (|\mathcal{F}[f](\eta) - \Psi(\eta)| \langle \xi_0 - \eta \rangle^s)^q d\xi \right)^{1/q} \\ &\leq \langle \xi_0 \rangle^s \|\mathcal{F}[f] - \Psi\|_{L_s^q} < \varepsilon \langle \xi_0 \rangle^s \end{aligned}$$

and similarly

$$\|\Psi(\xi - \cdot) - \mathcal{F}[f](\xi - \cdot)\|_{L_s^q} \leq \langle \xi \rangle^s \|\mathcal{F}[f] - \Psi\|_{L_s^q} < 2^s \varepsilon \langle \xi_0 \rangle^s.$$

Therefore, we have

$$\begin{aligned} &\|\mathcal{F}[f](\xi_0 - \cdot) - \mathcal{F}[f](\xi - \cdot)\|_{L_s^q} \\ &\leq \|\mathcal{F}[f](\xi_0 - \cdot) - \Psi(\xi_0 - \cdot)\|_{L_s^q} + \|\Psi(\xi_0 - \cdot) - \Psi(\xi - \cdot)\|_{L_s^q} + \|\Psi(\xi - \cdot) - \mathcal{F}[f](\xi - \cdot)\|_{L_s^q} \\ &\leq (1 + 2^s) \varepsilon \langle \xi_0 \rangle^s + \|\Psi(\xi_0 - \cdot) - \Psi(\xi - \cdot)\|_{L_s^q}. \end{aligned}$$

Moreover, since $\text{supp } \Psi(\xi_0 - \cdot) \subset B_n(0, |\xi_0| + R)$ and $\text{supp } \Psi(\xi - \cdot) \subset B_n(0, 1 + |\xi_0| + R)$, we have

$$\text{supp } |\Psi(\xi_0 - \cdot) - \Psi(\xi - \cdot)| \subset B_n(0, R_0)$$

with $R_0 := 1 + |\xi_0| + R$. Combining these estimates with (3), we have

$$\begin{aligned} \|\Psi(\xi_0 - \cdot) - \Psi(\xi - \cdot)\|_{L_s^q} &= \left(\int_{B_n(0, R_0)} (|\Psi(\xi_0 - \eta) - \Psi(\xi - \eta)| \langle \eta \rangle^s)^q d\eta \right)^{1/q} \\ &\leq \varepsilon R_0^{-(s+n/q)} \left(\int_{B_n(0, R_0)} \langle \eta \rangle^{sq} d\eta \right)^{1/q} \leq 2^s \varepsilon |B_n(0, 1)|^{1/q}, \end{aligned}$$

which implies the desired result.

(ii) Let $g \in L_s^q(\mathbf{R}^n)$. Then the Hölder and Young inequalities imply

$$\left| \int_{\mathbf{R}^n} (\mathcal{F}[f] * \Phi)(\xi) g(\xi) d\xi \right| \leq \int_{\mathbf{R}^n} \left(\int_{\mathbf{R}^n} |\mathcal{F}[f](\xi - \eta)| |g(\xi)| d\xi \right) |\Phi(\eta)| d\eta$$

$$\begin{aligned}
&\leq \|\Phi\|_{L_{-s}^{q'}} \left\| \int_{\mathbf{R}^n} |\mathcal{F}[f](\xi - \cdot)| |g(\xi)| d\xi \right\|_{L_s^q} \\
&\leq \|\Phi\|_{L_{-s}^{q'}} (\|\mathcal{F}[f]\|_{L_s^q} \|g\|_{L^1} + \|\mathcal{F}[f]\|_{L^1} \|g\|_{L_s^q}) \\
&\lesssim \|\Phi\|_{L_{-s}^q} \|f\|_{\mathcal{F}L_s^q} \|g\|_{L_s^q}
\end{aligned}$$

(cf. [19, Proposition 3.4]). Hence $\mathcal{F}[f] * \Phi \in L_{-s}^{q'}(\mathbf{R}^n)$. □

Remark 1 We can also prove Lemma 5 as follows: Since $L_s^q(\mathbf{R}^n)$ is a convolution algebra and $\langle \mathcal{F}[f] * \Phi, g \rangle = \langle \Phi, \mathcal{F}[\bar{f}] * g \rangle$ (at least for test functions), we have

$$|\langle \mathcal{F}[f] * \Phi, g \rangle| \leq \|\Phi\|_{L_{-s}^{q'}} \|\mathcal{F}[\bar{f}] * g\|_{L_s^q} \lesssim \|\Phi\|_{L_{-s}^{q'}} \|f\|_{\mathcal{F}L_s^q} \|g\|_{L_s^q}.$$

Hence $\mathcal{F}[f] * \Phi \in L_{-s}^{q'}(\mathbf{R}^n)$. The continuous translation property (1) then also gives continuity.

We also consider the differentiability of f in $\mathcal{F}L_s^q(\mathbf{R}^2)$.

Lemma 6 Let $1 < q < \infty$ and $s > 2/q' + 1/2$. Suppose that $f \in \mathcal{F}L_s^q(\mathbf{R}^2)$ be radial, i.e., $f(x) = f^+(|x|)$ for some $f^+(\rho)$ ($\rho \geq 0$). Then $f^+(\rho)$ is differentiable and satisfies

$$|(f^+)'(\rho)| \leq C_\rho \|f\|_{\mathcal{F}L_s^q} \quad (\rho > 0). \quad (4)$$

Proof Let $\rho > 0$. To see this, we use the polar coordinates $\xi = (r \cos \theta, r \sin \theta)$. Since $\mathcal{F}[f]$ is also radial, $\mathcal{F}[f](\xi) = \mathcal{F}[f](|\xi|, 0)$ and thus

$$\begin{aligned}
(f^+)(\rho) &= f(\rho, 0) = \frac{1}{(2\pi)^2} \int_{\mathbf{R}^2} \mathcal{F}[f](\xi) e^{i\xi \cdot (\rho, 0)} d\xi \\
&= \frac{1}{(2\pi)^2} \int_0^{2\pi} \left(\int_0^\infty \mathcal{F}[f](r, 0) e^{i\rho r \cos \theta} r dr \right) d\theta \\
&= \frac{1}{2\pi} \int_0^\infty \mathcal{F}[f](r, 0) J_0(\rho r) r dr,
\end{aligned}$$

where J_0 denotes the Bessel function defined in (2). Then by the mean value theorem and Lemma 4 (iii), we have for $h \in \mathbf{R}$ with $|h| < \rho/2$

$$\begin{aligned}
\frac{f^+(\rho + h) - f^+(\rho)}{h} &= \frac{1}{2\pi} \int_0^\infty \mathcal{F}[f](r, 0) \frac{J_0((\rho + h)r) - J_0(\rho r)}{rh} r^2 dr \\
&= \frac{1}{2\pi} \int_0^\infty \mathcal{F}[f](r, 0) (\operatorname{Re} J_0'((\rho + \theta_1 h)r) + i \operatorname{Im} J_0'((\rho + \theta_2 h)r)) r^2 dr \\
&= \frac{1}{2\pi} \int_0^\infty \mathcal{F}[f](r, 0) (\operatorname{Re} J_1((\rho + \theta_1 h)r) + i \operatorname{Im} J_1((\rho + \theta_2 h)r)) r^2 dr
\end{aligned}$$

for some $\theta_1, \theta_2 \in (0, 1)$. We note $|J_1(t)| \leq 1$ ($t \in \mathbf{R}$). Moreover, since $J_1(r) = O(r^{-1/2})$ ($r \rightarrow \infty$) and $(\rho + \theta_i h)r > \rho r/2$ ($i = 1, 2$), we have

$$\left| \int_1^\infty \mathcal{F}[f](r, 0) r^2 r^{-1/2} dr \right| \leq C_\rho \int_0^{2\pi} \int_0^\infty |\mathcal{F}[f](r, 0)| \langle r \rangle^{1/2} r dr d\theta$$

$$= \int_{\mathbf{R}^2} |\mathcal{F}[f](\xi)| \langle \xi \rangle^{1/2} d\xi \lesssim \|f\|_{\mathcal{F}L_s^q}.$$

Therefore, the Lebesgue dominated convergence theorem implies that $f^+(\rho)$ is differentiable and (4) holds. $\square$

The following lemma will be used in Section 7.

Lemma 7 Let $1 \leq q \leq 2$ and $2/q' < s < 2/q' + 1/2$. For $h \in C_c(\mathbf{R}^2) \cap \mathcal{F}L_s^1(\mathbf{R}^2)$ with $h(0) = 1$ and $\mathcal{F}[h](\xi) \geq 0$, we define

$$S_\rho(\mathcal{F}[h])(\xi) := \rho^2 \mathcal{F}[h](\rho\xi) \quad (\xi \in \mathbf{R}^2).$$

Then we have

$$\lim_{\rho \rightarrow \infty} \|S_\rho(\mathcal{F}[h]) * \mathcal{F}[f] - \mathcal{F}[f]\|_{L_s^q} = 0 \quad (5)$$

for all $f \in \mathcal{F}L_s^q(\mathbf{R}^2)$. Equivalently, we have

$$\lim_{\rho \rightarrow \infty} \left\| h\left(\frac{\cdot}{\rho}\right) f(\cdot) - f(\cdot) \right\|_{\mathcal{F}L_s^q} = 0 \quad (6)$$

for all $f \in \mathcal{F}L_s^q(\mathbf{R}^2)$.

Remark 2 Let $g = \frac{1}{2}\chi_{[-1,1]}(t)$ ($t \in \mathbf{R}$) and define a B-spline type function h by

$$h(x) := c((g * g * g * g)(x_1) \cdot (g * g * g * g)(x_2))$$

with $x = (x_1, x_2) \in \mathbf{R}^2$ or equivalently $\mathcal{F}[h](\xi) = c(\frac{\sin \xi_1}{\xi_1})^4 (\frac{\sin \xi_2}{\xi_2})^4$ with $\xi = (\xi_1, \xi_2) \in \mathbf{R}^2$. Here c is so chosen that $h(0) = \int_{\mathbf{R}^2} \mathcal{F}[h](\xi) d\xi = 1$. Since $\mathcal{F}[\frac{1}{2}\chi_{[-1,1]}](t) = \frac{\sin t}{t}$ and $s < 2$, we can easily see $h \in C_c(\mathbf{R}^2) \cap \mathcal{F}L_s^1(\mathbf{R}^2)$.

The proof of Lemma 7 We note $h(0) = \int_{\mathbf{R}^2} \mathcal{F}[h](\eta) d\eta = 1$ and

$$(S_\rho(\mathcal{F}[h]) * \mathcal{F}[f])(\xi) = \int_{\mathbf{R}^2} \mathcal{F}[h](\eta) \mathcal{F}[f]\left(\xi - \frac{\eta}{\rho}\right) d\eta.$$

By the Minkowski inequality for integral we have

$$\|S_\rho(\mathcal{F}[h]) * \mathcal{F}[f] - \mathcal{F}[f]\|_{L_s^q} \leq \int_{\mathbf{R}^2} |\mathcal{F}[h](\eta)| \left\| \mathcal{F}[f]\left(\cdot - \frac{\eta}{\rho}\right) - \mathcal{F}[f](\cdot) \right\|_{L_s^q} d\eta.$$

On the other hand, for all $\eta \in \mathbf{R}^n$, we have

$$\lim_{\rho \rightarrow \infty} \left\| \mathcal{F}[f]\left(\cdot - \frac{\eta}{\rho}\right) - \mathcal{F}[f](\cdot) \right\|_{L_s^q} = 0$$

and

$$\left\| \mathcal{F}[f]\left(\cdot - \frac{\eta}{\rho}\right) - \mathcal{F}[f](\cdot) \right\|_{L_s^q} \leq \left\langle \frac{\eta}{\rho} \right\rangle^s \|f\|_{\mathcal{F}L_s^q} + \|f\|_{\mathcal{F}L_s^q} \leq C_f \langle \eta \rangle^s$$

for $\rho \geq 1$. Since $h \in \mathcal{F}L_s^1(\mathbf{R}^2)$, the Lebesgue convergence theorem shows (5), and thus (6). $\square$

2.3 Closed Ideals in $\mathcal{FL}_s^q(\mathbf{R}^n)$

Throughout Section 2.3, we assume $1 < q < \infty$ and $s > n/q'$. Let I be a linear subspace of $\mathcal{FL}_s^q(\mathbf{R}^n)$. Then I is called an *ideal* in $\mathcal{FL}_s^q(\mathbf{R}^n)$ if $fg \in I$ whenever $f \in \mathcal{FL}_s^q(\mathbf{R}^n)$ and $g \in I$. Moreover, if an ideal I in $\mathcal{FL}_s^q(\mathbf{R}^n)$ is a closed subset of $\mathcal{FL}_s^q(\mathbf{R}^n)$, then I is called a *closed ideal* in $\mathcal{FL}_s^q(\mathbf{R}^n)$. For a subset S of $\mathcal{FL}_s^q(\mathbf{R}^n)$, let $\{I_\lambda\}_{\lambda \in \Lambda}$ be the set of all ideals containing S . Then the set $\bigcap_{\lambda \in \Lambda} I_\lambda$ is called the *ideal generated by S* . We note that the closed ideal generated by S coincides with the closure of

$$\left\{ \sum_{j=1}^n f_j g_j \mid f_j \in \mathcal{FL}_s^q(\mathbf{R}^n), g_j \in S, n \in \mathbf{N} \right\}.$$

Definition 1 Let I be a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$. Then the *zero-set* of I is defined by $Z(I) := \bigcap_{f \in I} f^{-1}(\{0\})$ with $f^{-1}(\{0\}) := \{x \in \mathbf{R}^n \mid f(x) = 0\}$.

We note that $x \in Z(I)$ if and only if $f(x) = 0$ for all $f \in I$. Moreover, $Z(I)$ is a closed subset of $\mathbf{R}^n$ whenever I is a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$. In fact, if $f \in \mathcal{FL}_s^q(\mathbf{R}^n)$, then f is continuous on $\mathbf{R}^n$ and thus $f^{-1}(\{0\})$ is a closed subset of $\mathbf{R}^n$.

The following lemma seems to be known to many people, but for the reader's convenience, we give the proof.

Lemma 8 Let E be a closed subset of $\mathbf{R}^n$ and set

$$I(E) := \{f \in \mathcal{FL}_s^q(\mathbf{R}^n) \mid f|_E = 0\}.$$

Then $I(E)$ is a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$ with $E = Z(I(E))$.

Proof It is clear that $I(E)$ is an ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$. To see $I(E)$ is closed, let $f \in \mathcal{FL}_s^q(\mathbf{R}^n)$, $\{f_n\}_{n=1}^\infty \subset I(E)$ and $\|f_n - f\|_{\mathcal{FL}_s^q} \rightarrow 0$ ($n \rightarrow \infty$). Since

$$\|f_n - f\|_{L^\infty} \lesssim \|\mathcal{F}[f_n] - \mathcal{F}[f]\|_{L^1} \lesssim \|f_n - f\|_{\mathcal{FL}_s^q},$$

we see that $\{f_n\}_{n=1}^\infty$ converges pointwise to f on $\mathbf{R}^n$. Since $f_n|_E = 0$, we have $f|_E = 0$, and thus $f \in I(E)$. Hence $I(E)$ is closed. Next, we prove $E = Z(I(E))$. Since $E \subset Z(I(E))$ is clear, we show $Z(I(E)) \subset E$. Suppose $x_0 \notin E$. Since E is closed and $C_c^\infty(\mathbf{R}^n) \subset \mathcal{FL}_s^q(\mathbf{R}^n)$, there exists $f \in \mathcal{FL}_s^q(\mathbf{R}^n)$ such that $f(x_0) = 1$ and $f|_E = 0$. Then $f \in I(E)$ and $f(x_0) \neq 0$. Thus $x_0 \notin Z(I(E))$, which implies the desired result. $\square$

Definition 2 Let E be a closed set $\mathbf{R}^n$. Then E is called a *set of spectral synthesis* for $\mathcal{FL}_s^q(\mathbf{R}^n)$ if we have $I_1 = I_2$ when $Z(I_1) = Z(I_2) = E$ for two closed ideals I_1 and I_2 (i.e., a closed ideal I with $Z(I) = E$ is a unique closed ideal).

We also prepare the following lemma, which also characterizes the closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$ (cf. [24, Theorem 7.1.2]). In the following, we call a closed ideal I in $\mathcal{FL}_s^q(\mathbf{R}^n)$ *modulation invariant* if $[M_\eta f](x) := e^{ix\eta} f(x) \in I$ whenever $f \in I$ and $\eta \in \mathbf{R}^n$.

Lemma 9 Let I be a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$ and $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$. Then

$$\langle M_{-\eta}f, \phi \rangle = 0 \quad (7)$$

for all $f \in I$ and $\eta \in \mathbf{R}^n$ if and only if

$$\langle f, \phi \rangle = 0 \quad (8)$$

for all $f \in I$. Moreover, every closed modulation invariant subspace of $\mathcal{FL}_s^q(\mathbf{R}^n)$ is an ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$. Conversely, every closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$ is modulation invariant.

Proof We give the proof only for (8) implies (7); the other case is clear. Assume that (8) holds for all $f \in I$. By Lemma 5, $\mathcal{F}[f] * \mathcal{F}[\bar{\phi}]$ is continuous and $f\bar{\phi} \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$. Moreover, since

$$\langle M_{-\eta}f, \phi \rangle = (2\pi)^{-n} (\mathcal{F}[f] * \mathcal{F}[\bar{\phi}])(\eta) = \mathcal{F}[f\bar{\phi}](\eta),$$

we need to prove $f\bar{\phi} = 0$. By duality, it suffices to prove $\langle g, f\bar{\phi} \rangle = 0$ for all $f \in I$ and $g \in \mathcal{FL}_s^q(\mathbf{R}^n)$. Let $f \in I$ and $g \in \mathcal{FL}_s^q(\mathbf{R}^n)$. As in the proof of Lemma 5 (ii), we have

$$\iint_{\mathbf{R}^n} |\mathcal{F}[f](\xi - \eta)| |\mathcal{F}[\bar{\phi}](\eta)| |\overline{\mathcal{F}[g](\xi)}| d\xi d\eta \lesssim \|f\|_{\mathcal{FL}_s^q} \|g\|_{\mathcal{FL}_s^q} \|\bar{\phi}\|_{\mathcal{FL}_{-s}^{q'}}.$$

Thus it follows from the Fubini Theorem that

$$\begin{aligned} \overline{\langle g, f\bar{\phi} \rangle} &= (2\pi)^{-2n} \int_{\mathbf{R}^n} (\mathcal{F}[f] * \mathcal{F}[\bar{\phi}])(\xi) \overline{\mathcal{F}[g](\xi)} d\xi \\ &= (2\pi)^{-2n} \int_{\mathbf{R}^n} (\mathcal{F}[f] * \mathcal{F}[\bar{g}])(\xi) \overline{\mathcal{F}[\phi](\xi)} d\xi = \langle f\bar{g}, \phi \rangle. \end{aligned}$$

Since $\bar{g} \in \mathcal{FL}_s^q(\mathbf{R}^n)$ and I is an ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$, we have $f\bar{g} \in I$ and thus $\langle f\bar{g}, \phi \rangle = 0$. Hence we have $\langle g, f\bar{\phi} \rangle = 0$.

Finally, we prove the characterization of a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$. Let I be a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$. Suppose that $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$ and (8) holds for all $f \in I$. Then by (7)

$$\langle M_{-\eta}f, \phi \rangle = 0 \quad (9)$$

for all $f \in I$ and $\eta \in \mathbf{R}^n$. If I is not modulation invariant, then there exist $f_0 \in I$ and $\eta_0 \in \mathbf{R}^n$ such that $M_{-\eta_0}f_0 \notin I$. Then the Hahn-Banach theorem and $(\mathcal{FL}_s^q(\mathbf{R}^n))' = \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$ imply that there exists $\phi_0 \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$ such that $\langle f, \phi_0 \rangle = 0$ for all $f \in I$ and $\langle M_{-\eta_0}f_0, \phi_0 \rangle = 1$. This contradicts (9). Conversely, we assume that I is a closed modulation invariant subspace of $\mathcal{FL}_s^q(\mathbf{R}^n)$. If I is not an ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$, then there exist $f_0 \in I$ and $g_0 \in \mathcal{FL}_s^q(\mathbf{R}^n)$ such that $f_0\bar{g}_0 \notin I$. Then the Hahn-Banach theorem implies that there exists $\phi_0 \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^n)$ such that

$$\langle f, \phi_0 \rangle = 0 \quad (10)$$

for all $f \in I$, and

$$\langle f_0\bar{g}_0, \phi_0 \rangle = 1. \quad (11)$$

On the other hand, since I is modulation invariant, we have $M_{-\xi}f_0 \in I$ for all $\xi \in \mathbf{R}^n$. Thus we have by (10)

$$\mathcal{F}[f_0\bar{\phi}_0](\xi) = (2\pi)^{-n} (\mathcal{F}[f_0] * \mathcal{F}[\bar{\phi}_0])(\xi) = \langle M_{-\xi}f_0, \phi_0 \rangle = 0. \quad (12)$$

Hence $f_0\bar{\phi}_0 = 0$ and thus $\langle f_0\bar{g}_0, \phi_0 \rangle = \overline{\langle g_0, f_0\bar{\phi}_0 \rangle} = 0$. However, this contradicts (11). $\square$

3 The proof of Theorem 1

Throughout Section 3, we simply write $I(S^1)$ in Lemma 8 as I . Let I_0 be the set of all functions g on $\mathbf{R}^2$ with $g = f^2$, where $f \in \mathcal{FL}_s^q(\mathbf{R}^2)$, $f|_{S^1} = 0$ and f is the inverse Fourier transform for some radial function on $\mathbf{R}^2$, and define I_1 as the closed ideal generated by I_0 in $\mathcal{FL}_s^q(\mathbf{R}^2)$. We note $Z(I) = S^1$ by Lemma 8. Moreover, by Lemma 9, I_1 is the smallest closed modulation invariant subspace of $\mathcal{FL}_s^q(\mathbf{R}^2)$ with $I_0 \subset I_1$. Thus I_1 coincides with the closure of the set

$$H := \left\{ \sum_{j=1}^m e^{ix\eta_j} g_j(x) \mid \eta_j \in \mathbf{R}^2, g_j \in I_0, m \in \mathbf{N} \right\}. \quad (13)$$

In the following, we show $Z(I_1) = S^1$ and $I \neq I_1$, which implies S^1 is not a set of spectral synthesis in $\mathcal{FL}_s^q(\mathbf{R}^2)$. To prove $I \neq I_1$, we construct $\ell_0 \in (\mathcal{FL}_s^q(\mathbf{R}^2))'$ and $f_0 \in I \setminus I_1$ such that $\ell_0 = 0$ on I_1 and $\ell_0(f_0) \neq 0$.

3.1 Technical Lemma

In the following lemma, μ denotes the Radon measure in $\mathbf{R}^2$ corresponding to a mass distributed homogeneously over the unit circle S^1 , i.e.,

$$\int_{\mathbf{R}^2} f(x) d\mu(x) := \frac{1}{2\pi} \int_0^{2\pi} \left(\int_0^\infty f(r \cos \theta, r \sin \theta) r d\delta_1(r) \right) d\theta,$$

where δ_1 is defined by $\int_0^\infty h(r) d\delta_1(r) := h(1)$ for $h \in C([0, \infty))$ (see for instance [21], [25]).

Lemma 10 Let $1 < q < \infty$ and $s > 2/q' + 1/2$. Define the function k_0 defined by

$$k_0(\xi) := \int_{\mathbf{R}^2} e^{ix\xi} d\mu(x) = \frac{1}{2\pi} \int_0^{2\pi} e^{i|\xi| \cos \theta} d\theta \quad (\xi \in \mathbf{R}^2)$$

and set $K_0(\xi) := i\xi_2 k_0(\xi)$ with $\xi = (\xi_1, \xi_2) \in \mathbf{R}^2$. Then

$$\ell_0(f) := \int_{\mathbf{R}^2} \mathcal{F}[f](\xi) K_0(\xi) d\xi \quad (f \in \mathcal{FL}_s^q(\mathbf{R}^2)) \quad (14)$$

defines a bounded linear functional on $\mathcal{FL}_s^q(\mathbf{R}^2)$.

Proof We first note that $|k_0(\xi)| \leq 1$ ($\xi \in \mathbf{R}^2$) and $k_0(\xi) = O(|\xi|^{-1/2})$ as $|\xi| \rightarrow \infty$ (see Lemma 4 (i)). Moreover, since $|\xi_2| \leq |\xi| \leq |\xi|^{1/2}$ whenever $|\xi| \leq 1$, the Hölder inequality show that

$$\begin{aligned} |\ell_0(f)| &\leq \int_{\mathbf{R}^2} |\mathcal{F}[f](\xi)| |\xi_2| |k_0(\xi)| d\xi \\ &\lesssim \int_{|\xi| \leq 1} |\mathcal{F}[f](\xi)| \langle \xi \rangle^{1/2} d\xi + \int_{|\xi| \geq 1} |\mathcal{F}[f](\xi)| \langle \xi \rangle \langle \xi \rangle^{-1/2} d\xi \\ &\leq 2 \|\langle \cdot \rangle^{-s+1/2}\|_{L^{q'}} \|f\|_{\mathcal{FL}_s^q}. \end{aligned}$$

We note that $s > 1/2 + 2/q'$ implies $(-s + 1/2)q' < -2$. Thus ℓ_0 defines a bounded linear functional on $\mathcal{FL}_s^q(\mathbf{R}^2)$. $\square$

Remark 3 Let $1 < q < \infty$, $s = 1/2 + 2/q'$ and define f_0 by

$$\mathcal{F}[f_0](\xi) = \chi_{|\xi| \leq 2}(\xi) |\xi|^{-5/2} (\log |\xi|)^{-1} \operatorname{sgn}(K_0(\xi)).$$

Then $f_0 \in \mathcal{FL}_s^q(\mathbf{R}^2)$ and $|\ell_0(f_0)| = \infty$. Thus ℓ_0 is not bounded on $\mathcal{FL}_s^q(\mathbf{R}^2)$.

3.2 The proof of Theorem 1

We divide the proof into three parts.

Step 1. We first prove $Z(I_1) = S^1$. Let $x_0 \in S^1$ and $f \in I_1$. Since I_1 is the closure of H defined in (13), there exists $\{f_n\}_{n=1}^\infty \subset H$ such that $\|f - f_n\|_{\mathcal{FL}_s^q} \rightarrow 0$ ($n \rightarrow \infty$). We note that $f_n(x_0) = 0$, and thus

$$|f(x_0)| = |f(x_0) - f_n(x_0)| \lesssim \|\mathcal{F}[f] - \mathcal{F}[f_n]\|_{L^1} \lesssim \|f - f_n\|_{\mathcal{FL}_s^q},$$

which implies $f(x_0) = 0$. Therefore $S^1 \subset Z(I_1)$. On the other hand, if $x_0 \notin S^1$, then there exists a radial function $f \in C_c^\infty(\mathbf{R}^2)$ such that $f|_{S^1} = 0$ and $f(x_0) \neq 0$. Thus $g := f^2 \in I_0$ and $x_0 \notin g^{-1}(\{0\})$, which implies $x_0 \notin Z(I_1)$.

Step 2. To prove $I \neq I_1$, we first show $\ell_0 = 0$ on I_1 , where ℓ_0 denotes the bounded linear functional on $\mathcal{FL}_s^q(\mathbf{R}^2)$ defined in Lemma 10. We note that since I_1 is the closure of H , it suffices to show $\ell_0(M_\eta g) = 0$ for all $g \in I_0$ and $\eta \in \mathbf{R}$. Let $g \in I_0$ and $\eta \in \mathbf{R}$. Then

$$\begin{aligned} \ell_0(M_\eta g) &= \int_{\mathbf{R}^2} \mathcal{F}_{x \rightarrow \xi}[e^{ix\eta}g(x)](\xi) K_0(\xi) d\xi \\ &= \int_{\mathbf{R}^2} \mathcal{F}[g](\xi - \eta) \left(i\xi_2 \int_{\mathbf{R}^2} e^{iy\xi} d\mu(y) \right) d\xi \\ &= \int_{\mathbf{R}^2} \left(\int_{\mathbf{R}^2} \mathcal{F}[g](\xi - \eta) \frac{\partial}{\partial y_2} [e^{iy\xi}] d\xi \right) d\mu(y) \\ &= \int_{\mathbf{R}^2} \left(\frac{\partial}{\partial y_2} \int_{\mathbf{R}^2} \mathcal{F}[g](\xi - \eta) e^{iy\xi} d\xi \right) d\mu(y) \\ &= (2\pi)^2 \int_{\mathbf{R}^2} \left(\frac{\partial}{\partial y_2} [e^{iy\eta}g(y)] \right) d\mu(y). \end{aligned}$$

On the other hand, since $g = f^2$ for some radial $f \in \mathcal{FL}_s^q(\mathbf{R}^2)$ with $f|_{S^1} = 0$, we have

$$\frac{\partial}{\partial y_2} [e^{iy\eta}g(y)] = \left(\frac{\partial}{\partial y_2} [e^{iy\eta}] \right) g(y) + e^{iy\eta} \left(\frac{\partial g}{\partial y_2} \right) (y)$$

$$= \left(\frac{\partial}{\partial y_2} [e^{iy\eta}] \right) (f(y))^2 + 2e^{iy\eta} f(y) \frac{\partial f}{\partial y_2}(y) = 0$$

for all $y \in S^1$. Therefore, we obtain $\ell_0(M_\eta g) = 0$.

Step 3. Finally, we show $I \neq I_1$. The proof is completed by showing that there exists $f_0 \in I$ such that $\ell_0(f_0) \neq 0$. Let τ be a nonnegative function in $C_c^\infty(\mathbf{R}^2)$ such that $\tau(x) = 1$ on $B_2(x^{(0)}, 10^{-5})$ and $\text{supp } \tau \subset B_2(x^{(0)}, 10^{-4})$ with $x^{(0)} := (1/2, \sqrt{3}/2)$, and define $f_0 \in C_c^\infty(\mathbf{R}^2)$ by

$$f_0(x) = \tau(x) \left(x_2 - \sqrt{1 - x_1^2} \right), \quad x = (x_1, x_2) \in \mathbf{R}^2.$$

We note that $f_0|_{S^1} = 0$ and $\frac{\partial f_0}{\partial x_2}(x) = \frac{\partial \tau}{\partial x_2}(x) (x_2 - \sqrt{1 - x_1^2}) + \tau(x)$, and thus $\frac{\partial f_0}{\partial x_2} \geq 0$ on S^1 . Therefore in the same way as Step 2, we obtain

$$\begin{aligned} \ell_0(f_0) &= \int_{\mathbf{R}^2} \mathcal{F}[f_0](\xi) K_0(\xi) d\xi \\ &= \int_{\mathbf{R}^2} \left(\frac{\partial}{\partial x_2} \int_{\mathbf{R}^2} \mathcal{F}[f_0](\xi) e^{ix\xi} d\xi \right) d\mu(x) \\ &= (2\pi)^2 \int_{\mathbf{R}^2} \frac{\partial f_0}{\partial x_2}(x) d\mu(x) \\ &= (2\pi)^2 \int_{\mathbf{R}^2} \left(\frac{\partial \tau}{\partial x_2}(x) \left(x_2 - \sqrt{1 - x_1^2} \right) + \tau(x) \right) d\mu(x) \\ &= (2\pi)^2 \int_{\mathbf{R}^2} \tau(x) d\mu(x) = 2\pi \int_0^{2\pi} \tau(\cos \theta, \sin \theta) d\theta > 0, \end{aligned}$$

which implies the desired result.

4 The proof of Theorem 2

Throughout Section 4, we assume $1 < q < 2$ and $2/q' < s < 2/q' + 1/2$. Moreover, we simply write I instead of $I(S^1)$, and J denotes the closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^2)$ generated by

$$J^0 := \{f \in \mathcal{FL}_s^q(\mathbf{R}^2) \mid f = 0 \text{ in a neighborhood of } S^1\}. \quad (15)$$

We note that I is the biggest closed ideal I' in $\mathcal{FL}_s^q(\mathbf{R}^2)$ with $Z(I') = S^1$, and J is the smallest closed ideal I'' in $\mathcal{FL}_s^q(\mathbf{R}^2)$ with $Z(I'') = S^1$ (see Theorem 17 below). In the following, we show $I = J$, which implies S^1 is a set of spectral synthesis in $\mathcal{FL}_s^q(\mathbf{R}^2)$. Since $J \subset I$ is clear, we need to prove $I \subset J$. The proof is completed by showing the following:

Proposition 11 *Let I' be a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^2)$ and $f_0 \in \mathcal{FL}_s^q(\mathbf{R}^2)$. If every $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^2)$ such that $\mathcal{F}[\phi] \in C(\mathbf{R}^2)$ and $\langle f, \phi \rangle = 0$ for all $f \in I'$ satisfies $\langle f_0, \phi \rangle = 0$, then f_0 is in I' .*

We use Proposition 11 with $I' = J$ and $f_0 \in I$. Moreover, to prove “every $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^2)$ such that $\mathcal{F}[\phi] \in C(\mathbf{R}^2)$ and $\langle f, \phi \rangle = 0$ for all $f \in J$ satisfies $\langle f_0, \phi \rangle = 0$ ”, we use an idea derived from Herz [17]. In the following, $\mathcal{FM}(\mathbf{R}^2)$ denotes the set of all Fourier-Stieltjes transform of the bounded regular Borel measures $M(\mathbf{R}^2)$, i.e.,

$$\mathcal{FM}(\mathbf{R}^2) = \left\{ \mathcal{F}[\mu] \mid \mathcal{F}[\mu](x) = \int_{\mathbf{R}^2} e^{-ix\xi} d\mu(\xi), \mu \in M(\mathbf{R}^2) \right\}.$$

Proposition 12 *Let $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^2)$, $\mathcal{F}[\phi] \in C(\mathbf{R}^2)$ and $\langle f, \phi \rangle = 0$ for all $f \in J$. Then there exists $\{\sigma_n[\phi]\}_n \subset \mathcal{FM}(\mathbf{R}^2)$ such that*

- (i) $\sigma_n[\phi] = \mathcal{F}[\mu_n]$ for some $\mu_n \in M(\mathbf{R}^2)$ with $\text{supp } \mu_n \subset S^1$.
- (ii) $\|\sigma_n[\phi]\|_{L_{-s}^{q'}} \leq C\|\phi\|_{\mathcal{FL}_{-s}^{q'}}$.
- (iii) $\{\sigma_n[\phi]\}_n$ converges to $\mathcal{F}[\phi]$ uniformly on any compact set of $\mathbf{R}^2$.

In fact, if Propositions 11 and 12 are true, then we can prove Theorem 2 as follows: Let $f_0 \in I$ and $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^2)$ be such that $\mathcal{F}[\phi] \in C(\mathbf{R}^2)$ and $\langle f, \phi \rangle = 0$ for all $f \in J$. Then there exists $\{\sigma_n[\phi]\}_n \subset \mathcal{FM}(\mathbf{R}^2)$ satisfying (i)-(iii) in Proposition 12. Since $\mathcal{F}[f_0] \cdot \mathcal{F}[\phi] \in L^1(\mathbf{R}^2)$ and $\mathcal{F}[f_0] \in L_s^q(\mathbf{R}^2)$, for any $\varepsilon > 0$, there exists a compact set K such that

$$\int_{\mathbf{R}^2 \setminus K} |\mathcal{F}[f_0](\xi) \mathcal{F}[\phi](\xi)| d\xi < \frac{\varepsilon}{4}$$

and

$$\left(\int_{\mathbf{R}^2 \setminus K} (|\mathcal{F}[f_0](\xi)| \langle \xi \rangle^s)^q d\xi \right)^{1/q} < \frac{\varepsilon}{4C\|\phi\|_{\mathcal{FL}_{-s}^{q'}}},$$

where C denotes the constant in Proposition 12 (ii). Then we have

$$\int_{\mathbf{R}^2 \setminus K} |\mathcal{F}[f_0](\xi) \sigma_n[\phi](\xi)| d\xi \leq \|\mathcal{F}[f_0]\|_{L_s^q(\mathbf{R}^2 \setminus K)} \|\sigma_n[\phi]\|_{L_{-s}^{q'}(\mathbf{R}^2 \setminus K)} < \frac{\varepsilon}{4}.$$

Therefore,

$$\begin{aligned} |\langle f_0, \phi \rangle - \langle f_0, \mu_n \rangle| &\lesssim \left| \int_K \mathcal{F}[f_0](\xi) \left(\overline{\mathcal{F}[\phi](\xi)} - \overline{\sigma_n[\phi](\xi)} \right) d\xi \right| \\ &\quad + \int_{\mathbf{R}^2 \setminus K} |\mathcal{F}[f_0](\xi)| (|\mathcal{F}[\phi](\xi)| + |\sigma_n[\phi](\xi)|) d\xi \\ &\leq \sup_{\xi \in K} |\mathcal{F}[\phi](\xi) - \sigma_n[\phi](\xi)| \int_K |\mathcal{F}[f_0](\xi)| d\xi + \frac{\varepsilon}{2}. \end{aligned}$$

On the other hand, since $f_0|_{S^1} = 0$ and $\text{supp } \mu_n \subset S^1$, we have

$$\begin{aligned} \langle f_0, \mu_n \rangle &= (2\pi)^{-2} \int_{\mathbf{R}^2} \mathcal{F}[f_0](\xi) \overline{\left(\int_{\mathbf{R}^2} e^{-ix\xi} d\mu_n(x) \right)} d\xi \\ &= (2\pi)^{-2} \int_{\mathbf{R}^2} \overline{\left(\int_{\mathbf{R}^2} \mathcal{F}[f_0](\xi) e^{ix\xi} d\xi \right)} d\mu_n(x) = \int_{\mathbf{R}^2} \overline{f_0(x)} d\mu_n(x) = 0. \end{aligned}$$

For this reason $\langle f_0, \phi \rangle = 0$, and we conclude $f_0 \in J$ by Proposition 11.

4.1 The proof of Proposition 11

Let $\varphi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^2)$ be such that $\langle f, \varphi \rangle = 0$ for all $f \in I'$. To show $f_0 \in I'$, it suffices to prove $\langle f_0, \varphi \rangle = 0$. In fact, if $f_0 \notin I'$, then the Hahn-Banach theorem yields a contradiction. Moreover, since $\langle f_0, \varphi \rangle = \mathcal{F}[f_0 \bar{\varphi}](0)$, it suffices to prove $f_0 \bar{\varphi} = 0$. We prove this by the duality argument. Let $g \in \mathcal{FL}_s^q(\mathbf{R}^2)$. Since $\bar{g} \in \mathcal{FL}_s^q(\mathbf{R}^2)$ and I' is an ideal in $\mathcal{FL}_s^q(\mathbf{R}^2)$, we see $f \bar{g} \in I'$ whenever $f \in I'$. Thus $\langle f, g\varphi \rangle = \langle f \bar{g}, \varphi \rangle = 0$ for all $f \in I'$. On the other hand, by Lemma 5 we have $g\varphi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^2)$ and $\mathcal{F}[g\varphi] \in C(\mathbf{R}^2)$. Thus by our assumption, $\langle f_0, g\varphi \rangle = 0$. Therefore $\overline{\langle g, f_0 \bar{\varphi} \rangle} = \langle f_0, g\varphi \rangle = 0$, and thus $f_0 \bar{\varphi} = 0$.

4.2 Technical lemmas

Throughout Section 4.2, we assume that $\phi \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^2)$ satisfies $\mathcal{F}[\phi] \in C(\mathbf{R}^2)$ and $\langle f, \phi \rangle = 0$ for all $f \in J$. In the following, we show that ϕ satisfies $\mathcal{F}[\phi] + \Delta[\mathcal{F}[\phi]] = 0$, where Δ stands for the Laplace operator. For this, we introduce auxiliary functions G_0 and Φ_h in Lemmas 13 and 14.

Lemma 13 Define $F_0(\xi) := \mathcal{F}^{-1}[\chi_{B_2(0,1/2)}](\xi)$ and $G_0(\xi) := (F_0(\xi))^2$ for $\xi \in \mathbf{R}^2$. Then we have

- (i) $G_0(\xi) = O(|\xi|^{-3})$ ($|\xi| \rightarrow \infty$).
- (ii) $\mathcal{F}[G_0](x) = 0$ ($|x| \geq 1$).
- (iii) $G_0 \in L_s^q(\mathbf{R}^2)$.

Proof (i) We use the polar coordinates $\xi = (R \cos \eta, R \sin \eta)$ and $x = (r \cos \theta, r \sin \theta)$. Since $x\xi = Rr \cos(\theta - \eta)$, we have

$$F_0(\xi) = \frac{1}{(2\pi)^2} \int_0^{1/2} r \left(\int_0^{2\pi} e^{iRr \cos(\theta - \eta)} d\theta \right) dr = \frac{1}{2\pi} \int_0^{1/2} r J_0(Rr) dr$$

by the periodicity of the cosine function and the definition of the Bessel function J_0 . Moreover, it follows from Lemma 4 (i) and (iv) that

$$F_0(\xi) = \frac{1}{2\pi R^2} \int_0^{R/2} r J_0(r) dr = \frac{1}{2\pi R^2} [r J_1(r)]_{r=0}^{r=R/2} = O(R^{-3/2})$$

as $R = |\xi| \rightarrow \infty$. Hence $G_0(\xi) = (F_0(\xi))^2 = O(|\xi|^{-3})$ as $|\xi| \rightarrow \infty$.

(ii) Since $G_0 \in C(\mathbf{R}^2)$, (i) implies $G_0 \in L^1(\mathbf{R}^2)$. Therefore

$$\mathcal{F}[G_0](x) = (2\pi)^{-2} \left(\chi_{B_2(0,1/2)} * \chi_{B_2(0,1/2)} \right) (x).$$

Hence we obtain $\text{supp } \mathcal{F}[G_0] \subset B_2(0, 1/2) + B_2(0, 1/2) = B_2(0, 1)$.

(iii) By (i) and the continuity of G_0 , we see that $G_0 \in L_s^q(\mathbf{R}^2)$ if and only if $\int_{|\xi| \geq 1} (|G_0(\xi)| \langle \xi \rangle^s)^q d\xi < \infty$, which is equivalent to $(3-s)q > 2$. Hence, we get $G_0 \in L_s^q(\mathbf{R}^2)$ by our assumption on s . $\square$

Lemma 14 Let $h > 0$ and G_0 be the function defined in Lemma 13. Define

$$\Phi_h(x) := \int_{\mathbf{R}^2} e^{-ix\xi} G_0(h\xi) \overline{\mathcal{F}[\phi](\xi)} d\xi \quad (x \in \mathbf{R}^2).$$

Then $\text{supp } \Phi_h \subset K_h := \{x \in \mathbf{R}^2 \mid d(x, S^1) \leq 2h\}$.

Proof We first note

$$\mathcal{F}_{\xi \rightarrow u}^{-1}[e^{-ix\xi} G_0(h\xi)](u) = \frac{1}{(2\pi h)^2} \mathcal{F}[G_0] \left(\frac{x-u}{h} \right)$$

and thus

$$\Phi_h(x) = \frac{1}{(2\pi h)^2} \int_{\mathbf{R}^2} \mathcal{F} \left[\mathcal{F}[G_0] \left(\frac{x-\cdot}{h} \right) \right] (\xi) \overline{\mathcal{F}[\phi](\xi)} d\xi.$$

By our assumption on ϕ , it suffices to show $\mathcal{F}[G_0] \left(\frac{x-\cdot}{h} \right) \in J^0 \subset J$ if $d(x, S^1) > 2h$. Suppose $d(x, S^1) > 2h$ and let $y \in \mathbf{R}^2$ be such that $d(y, S^1) < h$. Since $|x-y| \geq d(x, S^1) - d(y, S^1) > h$, we have $\mathcal{F}[G_0] \left(\frac{x-y}{h} \right) = 0$ by Lemma 13 (ii). Hence, $\mathcal{F}[G_0] \left(\frac{x-\cdot}{h} \right) \in J^0$. $\square$

Lemma 15 ϕ satisfies $\mathcal{F}[\phi] + \Delta[\mathcal{F}[\phi]] = 0$.

Proof We will divide the proof into four parts.

Step 1. Let $f \in \mathcal{F}L_s^q(\mathbf{R}^n)$ and Φ_h be the function in Lemma 14. Firstly, we show

$$|\langle M_{-\eta} f, \phi \rangle| \lesssim \liminf_{h \downarrow 0} \left(\int_{K_h} |f(x)|^2 dx \right)^{1/2} \left(\int_{K_h} |\Phi_h(x)|^2 dx \right)^{1/2}.$$

To see this, we first note that $\Phi_h \in L^1(\mathbf{R}^2)$ and $G_0(h\xi) \overline{\mathcal{F}[\phi](\xi)} \in L^1(\mathbf{R}_\xi^2)$, since

$$|\Phi_h(x)| \leq \int_{\mathbf{R}^2} |G_0(h\xi)| |\mathcal{F}[\phi](\xi)| d\xi \leq C_h \|G_0\|_{L_s^q} \|\phi\|_{\mathcal{F}L_{-s}^{q'}}$$

and $\text{supp } \Phi_h \subset K_h$. Then the Fourier inversion formula yields

$$G_0(h\xi) \overline{\mathcal{F}[\phi](\xi)} = (2\pi)^{-2} \mathcal{F}^{-1}[\Phi_h](\xi) = (2\pi)^{-2} \mathcal{F}[\Phi_h](-\xi)$$

for all $\xi \in \mathbf{R}^2$. From the above equality with $G_0(0) = 1$, we obtain

$$(2\pi)^2 \langle M_{-\eta} f, \phi \rangle = \langle \mathcal{F}[f], \mathcal{F}[M_\eta \phi] \rangle_{L^2(\mathbf{R}^2)}$$

$$\begin{aligned}
&= \langle \mathcal{F}[f](\xi), \overline{G_0(0)\mathcal{F}[\phi](\xi - \eta)} \rangle_{L^2(\mathbf{R}^2)} \\
&= \liminf_{h \downarrow 0} \langle \mathcal{F}[f](\xi), \overline{G_0(h(\xi - \eta))\mathcal{F}[\phi](\xi - \eta)} \rangle_{L^2(\mathbf{R}^2)} \\
&= (2\pi)^{-2} \liminf_{h \downarrow 0} \langle \mathcal{F}[f](\xi), \overline{\mathcal{F}[\Phi_h](\eta - \xi)} \rangle_{L^2(\mathbf{R}^2)} \\
&= (2\pi)^{-2} \liminf_{h \downarrow 0} \langle \mathcal{F}[f](\xi), \mathcal{F}[\overline{M_\eta \Phi_h}] \rangle_{L^2(\mathbf{R}^2)} \\
&= \liminf_{h \downarrow 0} \langle f, \overline{M_\eta \Phi_h} \rangle.
\end{aligned}$$

Hence, the Cauchy-Schwarz inequality and the Plancherel theorem yield the desired result.

Step 2. Secondly, we prove

$$\left(\int_{K_h} |\Phi_h(x)|^2 dx \right)^{1/2} \leq Ch^{-1-s+2/q'} \|\phi\|_{\mathcal{FL}_{-s}^{q'}} \quad (16)$$

for $0 < h < 1/2$. Since $\Phi_h(x) = \mathcal{F}_{\xi \rightarrow x}[G_0(h\xi)\overline{\mathcal{F}[\phi](\xi)}](x)$ and $\text{supp } \Phi_h \subset K_h$, we obtain by the Plancherel theorem and the Hölder inequality with $\frac{1}{q'-2} + \frac{1}{q'} = 1$,

$$\begin{aligned}
\int_{K_h} |\Phi_h(x)|^2 dx &\approx \int_{\mathbf{R}^2} (|G_0(h\xi)|\langle \xi \rangle^s)^2 \left(|\mathcal{F}[\phi](\xi)|\langle \xi \rangle^{-s} \right)^2 d\xi \\
&\leq \left(\int_{\mathbf{R}^2} (|G_0(h\xi)|\langle \xi \rangle^s)^{2q'/(q'-2)} d\xi \right)^{(q'-2)/q'} \|\phi\|_{\mathcal{FL}_{-s}^{q'}}^2.
\end{aligned}$$

On the other hand, we note $\langle \xi/h \rangle \leq \langle \xi \rangle/h$ for $0 < h < 1/2$, and $2q'(-3+s)/(q'-2) < -2$ if and only if $s < 2 + 2/q'$. Thus we obtain

$$\int_{\mathbf{R}^2} (|G_0(h\xi)|\langle \xi \rangle^s)^{2q'/(q'-2)} d\xi = \frac{1}{h^2} \int_{\mathbf{R}^2} \left(|G_0(\xi)| \left\langle \frac{\xi}{h} \right\rangle^s \right)^{2q'/(q'-2)} d\xi \lesssim h^{-2-\frac{2sq'}{q'-2}},$$

since G_0 is continuous and satisfies $G_0(\xi) = O(|\xi|^{-3})$ as $|\xi| \rightarrow \infty$. Combining these inequalities, we obtain (16).

Step 3. Thirdly, we show that there exists $f_0 \in \mathcal{FL}_s^q(\mathbf{R}^n)$ such that

$$\int_{K_h} |f_0(x)|^2 dx = O(h^3) \quad (\text{as } h \rightarrow 0).$$

Let $k \in C_c^\infty(\mathbf{R})$ be such that $\text{supp } k \subset B_1(0, 3/2)$, $k(t) = 1$ on $B_1(0, 5/4)$ and $0 \leq k(t) \leq 1$ ($t \in \mathbf{R}$), and define real-valued $\tau \in C_c^\infty(\mathbf{R}^2)$ by $\tau(x) := k(|x|)$. We first show $\phi = \tau\phi$ by the duality argument. Let $g \in C_c^\infty(\mathbf{R}^2) \subset \mathcal{FL}_s^q(\mathbf{R}^2)$ and $\psi \in C_c^\infty(\mathbf{R}^2)$ be such that $\psi(x) = 1$ on $\text{supp } g$. Moreover, define real-valued $\tau_0 \in C_c^\infty(\mathbf{R}^2)$ by

$$\tau_0(x) := (1 - \tau(x))\psi(x).$$

Then $\text{supp } \tau_0 \subset \mathbf{R}^2 \setminus S^1$, $\tau(x) + \tau_0(x) = 1$ on $\text{supp } g$ and $S^1 \subset \text{int}(\text{supp } \tau)$. Since $g = (\tau + \tau_0)g$, we obtain

$$\langle g, \phi \rangle = \langle g, \tau\phi \rangle + \langle g, \tau_0\phi \rangle.$$

We note that $\tau_0 = 0$ in a neighborhood of S^1 , and thus $\tau_0 \in J$. Moreover, since J is closed modulation invariant subspace of $\mathcal{FL}_s^q(\mathbf{R}^n)$, we have $M_\eta \tau_0 \in J$ for all $\eta \in \mathbf{R}^2$. Thus, we obtain

$$\mathcal{F}[\tau_0\phi](\eta) = \overline{\langle M_\eta \tau_0, \phi \rangle} = 0,$$

which implies $\langle g, \tau_0 \phi \rangle = 0$, and thus $\langle g, \phi \rangle = \langle g, \tau \phi \rangle$. Hence $\phi = \tau \phi$. From this we get

$$\mathcal{F}[\phi](\eta) = \mathcal{F}[\tau \phi](\eta) = \overline{\langle M_\eta \tau, \phi \rangle} \quad (\eta \in \mathbf{R}^2).$$

Moreover, since $\mathcal{F}[\tau] \in \mathcal{S}(\mathbf{R}^2)$, we have

$$\mathcal{F}[\phi](\eta) + \Delta[\mathcal{F}[\phi]](\eta) = \overline{\langle (1 + \Delta_\eta)[M_\eta \tau], \phi \rangle} = \overline{\langle M_\eta f_0, \phi \rangle},$$

where we denote $f_0(x) := (1 - |x|^2)\tau(x)$. Therefore,

$$\int_{K_h} |f_0(x)|^2 dx = \int_{K_h} (1 - |x|^2)^2 |\tau(x)|^2 dx \leq C \|\tau\|_{L^\infty} (1 - |1 + 2h|^2)^2 \int_{1-2h}^{1+2h} r dr \leq C_\tau h^3$$

for $0 < h < 1/2$. Hence, we obtain the desired result.

Step 4. Finally, we prove $\mathcal{F}[\phi] + \Delta[\mathcal{F}[\phi]] = 0$. Let f_0 be the function defined in Step 3. Then by Step 1, we have

$$|\langle M_\eta f_0, \phi \rangle| \lesssim \liminf_{h \downarrow 0} \left(\int_{K_h} |f_0(\xi)|^2 d\xi \right) \left(\int_{K_h} |\Phi_h(\xi)|^2 d\xi \right) \lesssim \liminf_{h \downarrow 0} h^{3-2-2s+\frac{4}{q'}} = 0$$

by our assumption $s < 1/2 + 2/q'$, and thus

$$\mathcal{F}[\phi](\eta) + \Delta[\mathcal{F}[\phi]](\eta) = \overline{\langle M_\eta f_0, \phi \rangle} = 0.$$

□

4.3 The proof of Proposition 12

Let $k \in \mathbf{Z}$. We consider the polar coordinates $\xi_1 = r \cos \theta$ and $\xi_2 = r \sin \theta$ in $\mathbf{R}^2$, and define a function ϕ_k on $\mathbf{R}^2$ by

$$\mathcal{F}[\phi_k](\xi) := C_k(\phi; r) e^{ik\theta} \quad (\xi = (\xi_1, \xi_2) \in \mathbf{R}^2)$$

with

$$C_k(\phi; r) := \frac{1}{2\pi} \int_0^{2\pi} \mathcal{F}[\phi](r \cos \omega, r \sin \omega) e^{-ik\omega} d\omega.$$

Step 1. We first show that ϕ_k satisfies $\mathcal{F}[\phi_k] \in C(\mathbf{R}^2)$, $\phi_k \in \mathcal{F}L_{-s}^{q'}(\mathbf{R}^2)$ and $\langle f, \phi_k \rangle = 0$ for all $f \in J$. The continuity of $\mathcal{F}[\phi_k]$ follows from

$$\begin{aligned} \mathcal{F}[\phi_k](\xi) &= \frac{1}{2\pi} \int_{-\theta}^{2\pi-\theta} \mathcal{F}[\phi](r \cos(\omega + \theta), r \sin(\omega + \theta)) e^{-ik\omega} d\omega \\ &= \frac{1}{2\pi} \int_0^{2\pi} \mathcal{F}[\phi](r \cos(\omega + \theta), r \sin(\omega + \theta)) e^{-ik\omega} d\omega \end{aligned}$$

and the continuity of $\mathcal{F}[\phi]$ with the Lebesgue convergence theorem. To prove $\phi \in \mathcal{F}L_{-s}^{q'}(\mathbf{R}^2)$, we note that by the Hölder inequality

$$\begin{aligned} |\mathcal{F}[\phi_k](\xi)| &\leq \frac{1}{2\pi} \int_0^{2\pi} |\mathcal{F}[\phi](r \cos \omega, r \sin \omega)| d\omega \\ &\leq (2\pi)^{1/q-1} \left(\int_0^{2\pi} |\mathcal{F}[\phi](r \cos \omega, r \sin \omega)|^{q'} d\omega \right)^{1/q'} \end{aligned}$$

and also we have

$$\|\phi\|_{\mathcal{FL}_{-s}^{q'}}^{q'} = \int_0^\infty \left(\int_0^{2\pi} |\mathcal{F}[\phi](r \cos \omega, r \sin \omega)|^{q'} d\omega \right) \langle r \rangle^{-sq'} r dr.$$

Therefore,

$$\begin{aligned} \|\phi_k\|_{\mathcal{FL}_{-s}^{q'}}^{q'} &= \int_0^{2\pi} \left(\int_0^\infty |\mathcal{F}[\phi_k](r \cos \theta, r \sin \theta)|^{q'} \langle r \rangle^{-sq'} r dr \right) d\theta \\ &\lesssim \int_0^{2\pi} \left(\int_0^\infty \left(\int_0^{2\pi} |\mathcal{F}[\phi](r \cos \omega, r \sin \omega)|^{q'} d\omega \right) \langle r \rangle^{-sq'} r dr \right) d\theta \lesssim \|\phi\|_{\mathcal{FL}_{-s}^{q'}}^{q'}, \end{aligned}$$

which gives $\phi_k \in \mathcal{FL}_{-s}^{q'}(\mathbf{R}^2)$. Finally, we prove $\langle f, \phi_k \rangle = 0$ for all $f \in J$. Since J is the closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^2)$ generated by J^0 , it is enough to prove $\langle M_{-\eta} f_0, \phi_k \rangle = 0$ for all $f_0 \in J^0$ and $\eta = (\eta_1, \eta_2) \in \mathbf{R}^2$ (cf. (13)). By the polar coordinates

$$\begin{aligned} (2\pi)^2 \langle M_{-\eta} f_0, \phi_k \rangle &= \langle \mathcal{F}[M_{-\eta} f_0], \mathcal{F}[\phi_k] \rangle_{L^2(\mathbf{R}^2)} \\ &= \int_0^\infty \left(\int_0^{2\pi} \mathcal{F}[f_0](r \cos \theta + \eta_1, r \sin \theta + \eta_2) \overline{\mathcal{F}[\phi_k](r \cos \theta, r \sin \theta)} d\theta \right) r dr \end{aligned}$$

and

$$\overline{\mathcal{F}[\phi_k](r \cos \theta, r \sin \theta)} = \frac{1}{2\pi} \int_0^{2\pi} \overline{\mathcal{F}[\phi](r \cos(\omega + \theta), r \sin(\omega + \theta))} e^{ik\omega} d\omega.$$

Combining these estimates with the Fubini theorem and the change of variables, we obtain

$$\begin{aligned} \langle M_{-\eta} f_0, \phi_k \rangle &= \frac{1}{(2\pi)^3} \int_0^{2\pi} \left(\int_0^\infty \left(\int_0^{2\pi} \mathcal{F}[f_0](r \cos(\theta - \omega) + \eta_1, r \sin(\theta - \omega) + \eta_2) \right. \right. \\ &\quad \left. \left. \times \overline{\mathcal{F}[\phi](r \cos \theta, r \sin \theta)} d\theta \right) r dr \right) e^{ik\omega} d\omega. \end{aligned}$$

We note $r \cos(\theta - \omega) = \xi_1 \cos \omega + \xi_2 \sin \omega$, $r \sin(\theta - \omega) = \xi_2 \cos \omega - \xi_1 \sin \omega$. Then we can easily see

$$\begin{aligned} &\mathcal{F}[f_0](\xi_1 \cos \omega + \xi_2 \sin \omega + \eta_1, -\xi_1 \sin \omega + \xi_2 \cos \omega + \eta_2) \\ &= \mathcal{F}[f_0]((\xi_1, \xi_2)^t A + (\eta_1, \eta_2)) \\ &= \iint_{\mathbf{R}^2} f_0(x_1, x_2) e^{-i((\xi_1, \xi_2)^t A + (\eta_1, \eta_2)) \cdot (x_1, x_2)} dx_1 dx_2 \\ &= \iint_{\mathbf{R}^2} e^{-i(\eta_1, \eta_2) \cdot (y_1, y_2)^t A} f_0((y_1, y_2)^t A) e^{-i(\xi_1, \xi_2) \cdot (y_1, y_2)} dy_1 dy_2 = \mathcal{F}[F_0](\xi) \end{aligned}$$

with $A = \begin{pmatrix} \cos \omega & \sin \omega \\ -\sin \omega & \cos \omega \end{pmatrix}$ and $F_0(y) = e^{-i\eta \cdot y^t A} f_0(y^t A)$, where ${}^t A$ denotes the transpose of A . Therefore we have

$$\langle M_{-\eta} f_0, \phi_k \rangle = \frac{1}{(2\pi)^3} \int_0^{2\pi} \left(\int_{\mathbf{R}^2} \mathcal{F}[F_0](\xi) \overline{\mathcal{F}[\phi](\xi)} d\xi \right) e^{ik\omega} d\omega.$$

Moreover, since $f_0 \in J^0$ and A is a rotation matrix in $\mathbf{R}^2$, and also J is a closed modulation invariant subspace, we have $F_0 \in J$ and thus $\langle F_0, \phi \rangle = 0$. Hence $\langle M_{-\eta} f_0, \phi_k \rangle = 0$ for all $f_0 \in J^0$ and $\eta \in \mathbf{R}^2$.

Step 2. We note that ϕ_k satisfies $\mathcal{F}[\phi_k] + \Delta[\mathcal{F}[\phi_k]] = 0$. In fact, as we have already seen in Step 1, ϕ_k satisfies the same condition to ϕ . Thus we get the desired result.

Step 3. Now we give the proof of Proposition 12. By Step 2 and the polar coordinates, we have

$$\begin{aligned} 0 &= \mathcal{F}[\phi_k] + \Delta[\mathcal{F}[\phi_k]] = \mathcal{F}[\phi_k] + \left(\frac{\partial^2}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} + \frac{1}{r} \frac{\partial}{\partial r} \right) \mathcal{F}[\phi_k] \\ &= C_k(\phi; r) e^{ik\theta} + \left(\frac{\partial^2}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} + \frac{1}{r} \frac{\partial}{\partial r} \right) (C_k(\phi; r) e^{ik\theta}) \\ &= \left(C_k(\phi; r) + \frac{\partial^2}{\partial r^2} C_k(\phi; r) + \frac{1}{r^2} C_k(\phi; r) (-k^2) + \frac{1}{r} \frac{\partial}{\partial r} C_k(\phi; r) \right) e^{ik\theta}. \end{aligned}$$

Then, we get

$$\left(\frac{\partial^2}{\partial r^2} + \left(1 - \frac{k^2}{r^2} \right) + \frac{1}{r} \frac{\partial}{\partial r} \right) C_k(\phi; r) = 0.$$

We note that the above equality is just the Bessel differential equation, and we can state $C_k(\phi; r) = C_k(\phi) J_k(r)$ for some $C_k(\phi) \in \mathbf{R}$, where J_k is the Bessel function defined in (2).

The proof of Proposition 12 (i)

We consider the polar coordinates in $\mathbf{R}^2$, i.e., $\xi_1 = r \cos \theta$ and $\xi_2 = r \sin \theta$, and define a modified version of the Fejér kernel by

$$\sigma_N[\phi](\xi) := \sum_{k=-N}^N \left(1 - \frac{|k|}{N+1} \right) C_k(\phi; r) e^{ik\theta}.$$

Moreover, let ν_k be the measure on $\mathbf{R}^2$ defined by

$$\int_{\mathbf{R}^2} f(x) d\nu_k(x) := \frac{1}{2\pi} \int_0^{2\pi} \left(\int_0^\infty f(r \cos \omega, r \sin \omega) r d\delta_1(r) \right) e^{ik\omega} d\omega$$

(cf. Section 3.1). Then we have

$$\begin{aligned}
\mathcal{F}[\nu_k](\xi) &= \mathcal{F}[\nu_k](r \cos \theta, r \sin \theta) = \int_{\mathbf{R}^2} e^{ix \cdot (r \cos \theta, r \sin \theta)} d\nu_k(x) \\
&= \frac{1}{2\pi} \int_0^{2\pi} e^{ir \cos(\omega - \theta)} e^{ik\omega} d\omega = \frac{1}{2\pi} \int_{-\theta - \frac{\pi}{2}}^{2\pi - \theta - \frac{\pi}{2}} e^{ir \cos(\omega + \frac{\pi}{2})} e^{ik(\omega + \theta + \frac{\pi}{2})} d\omega \\
&= \frac{e^{ik(\theta + \frac{\pi}{2})}}{2\pi} \int_{-\theta - \frac{\pi}{2}}^{2\pi - \theta - \frac{\pi}{2}} e^{-ir \sin \omega} e^{ik\omega} d\omega = i^k J_k(r) e^{ik\theta}
\end{aligned}$$

and thus $C_k(\phi; r) e^{ik\theta} = C_k(\phi) J_k(r) e^{ik\theta} = i^{-k} \mathcal{F}[\nu_k](\xi)$. Hence

$$\mu_k(x) := \sum_{k=-N}^N i^{-k} \left(1 - \frac{|k|}{N+1}\right) C_k(\phi) \nu_k(x)$$

satisfies $\sigma_N[\phi] = \mathcal{F}[\mu_k]$ and $\text{supp } \mu_N \subset S^1$.

The proof of Proposition 12 (ii)

We first note that

$$\begin{aligned}
\sigma_N[\phi](\xi) &= \sum_{k=-N}^N \left(1 - \frac{|k|}{N+1}\right) \left(\frac{1}{2\pi} \int_0^{2\pi} \mathcal{F}[\phi](r \cos \omega, r \sin \omega) e^{-ik\omega} d\omega \right) e^{ik\theta} \\
&= \int_0^{2\pi} \mathcal{F}[\phi](r \cos \omega, r \sin \omega) \left(\frac{1}{2\pi} \sum_{k=-N}^N \left(1 - \frac{|k|}{N+1}\right) e^{ik(\theta - \omega)} \right) d\omega \\
&= (\mathcal{F}[\phi]_r \star K_N)(\theta),
\end{aligned}$$

where $\mathcal{F}[\phi]_r(\theta) := \mathcal{F}[\phi](r \cos \theta, r \sin \theta)$, $K_N(\theta) := \frac{1}{2\pi} \sum_{k=-N}^N \left(1 - \frac{|k|}{N+1}\right) e^{ik\theta}$ (i.e., K_N is the Fejér kernel) and $\star$ is the operator defined by $(G \star H)(\theta) := \int_0^{2\pi} G(\omega) H(\theta - \omega) d\omega$. Then it is well-known that $K_N(\theta) \geq 0$ and $\int_0^{2\pi} K_N(\theta) d\theta = 1$. Therefore we have by the Hölder inequality

$$\begin{aligned}
\|\sigma_N[\phi]\|_{L_{-s}^{q'}}^{q'} &= \int_0^{2\pi} \left(\int_0^\infty (|(\mathcal{F}[\phi]_r \star K_N)(\theta)| \langle r \rangle^{-s})^{q'} r dr \right) d\theta \\
&\lesssim \int_0^{2\pi} \left(\int_0^\infty \left(\int_0^{2\pi} |\mathcal{F}[\phi]_r(\theta - \omega)| K_N(\omega) d\omega \right)^{q'} \langle r \rangle^{-sq'} r dr \right) d\theta \\
&\leq \int_0^{2\pi} \left(\int_0^\infty \left(\int_0^{2\pi} |\mathcal{F}[\phi]_r(\theta - \omega)|^{q'} K_N(\omega) d\omega \right) \left(\int_0^{2\pi} K_N(\omega) d\omega \right)^{\frac{q'}{q}} \langle r \rangle^{-sq'} r dr \right) d\theta \\
&= \int_0^{2\pi} \left(\int_0^\infty \left(\int_0^{2\pi} |\mathcal{F}[\phi]_r(\theta - \omega)|^{q'} K_N(\omega) d\omega \right) \langle r \rangle^{-sq'} r dr \right) d\theta
\end{aligned}$$

$$= \int_0^{2\pi} K_N(\omega) \left(\int_0^\infty \left(\int_0^{2\pi} |\mathcal{F}[\phi]_r(\theta - \omega)|^{q'} d\theta \right) \langle r \rangle^{-sq'} r dr \right) d\omega = \|\mathcal{F}[\phi]\|_{L_{-s}^{q'}}^{q'},$$

which implies the desired result.

The proof of Proposition 12 (iii)

For the sake of simplicity, we set

$$\mathcal{F}[\phi]_r(\theta) := \mathcal{F}[\phi](r \cos \theta, r \sin \theta), \quad \sigma_N[\phi]_r(\theta) := \sigma_N[\phi](r \cos \theta, r \sin \theta).$$

Let $R > 0$ and $\varepsilon > 0$. Since $\mathcal{F}[\phi] \in C(\mathbf{R}^2)$, $\mathcal{F}[\phi]$ is uniformly continuous on any compact set on $\mathbf{R}^2$. Thus there exists $\delta > 0$ such that for any $0 \leq r \leq R$, $0 \leq \theta \leq 2\pi$ and $|\eta| < \delta$, we have

$$|\mathcal{F}[\phi]_r(\theta - \eta) - \mathcal{F}[\phi]_r(\theta)| < \varepsilon. \quad (17)$$

Now suppose that $0 \leq r \leq R$ and $0 \leq \theta \leq 2\pi$. Since $\int_0^{2\pi} K_N(\theta) d\theta = 1$, we have

$$\begin{aligned} |\sigma_N[\phi]_r(\theta) - \mathcal{F}[\phi]_r(\theta)| &= |(\mathcal{F}[\phi]_r \star K_N)(\theta) - \mathcal{F}[\phi]_r(\theta)| \\ &= \left| \int_0^{2\pi} (\mathcal{F}[\phi]_r(\theta - \eta) - \mathcal{F}[\phi]_r(\theta)) K_N(\eta) d\eta \right| \\ &\leq \int_{|\eta| < \delta} |\mathcal{F}[\phi]_r(\theta - \eta) - \mathcal{F}[\phi]_r(\theta)| K_N(\eta) d\eta \\ &\quad + \int_{|\eta| \geq \delta} |\mathcal{F}[\phi]_r(\theta - \eta) - \mathcal{F}[\phi]_r(\theta)| K_N(\eta) d\eta \\ &\leq \varepsilon + \int_{|\eta| \geq \delta} |\mathcal{F}[\phi]_r(\theta - \eta) - \mathcal{F}[\phi]_r(\theta)| K_N(\eta) d\eta. \end{aligned}$$

Moreover, since $K_N(\eta) = \frac{1}{2\pi(N+1)} \frac{\sin^2 \frac{(N+1)\eta}{2}}{\sin^2 \frac{\eta}{2}}$, there exists $C_\delta > 0$ such that $K_N(\eta) \leq \frac{C_\delta}{(N+1)}$ holds for all $\delta \leq |\eta| \leq 2\pi$. Therefore, we obtain

$$|\sigma_N[\phi]_r(\theta) - \mathcal{F}[\phi]_r(\theta)| < \varepsilon + \frac{C_\delta}{(N+1)} \sup_{|\xi| \leq R} |\mathcal{F}[\phi](\xi)|.$$

If N is sufficiently large, we obtain

$$|\sigma_N[\phi]_r(\theta) - \mathcal{F}[\phi]_r(\theta)| < 2\varepsilon$$

for all $0 \leq r \leq R$ and $0 \leq \theta \leq 2\pi$, which implies that $\{\sigma_N[\phi]\}_{N=1}^\infty$ converges uniformly to $\mathcal{F}[\phi]$ on any compact set.

5 The proof of Theorem 3

Let $\sigma(t)$ be the Radon measure in $\mathbf{R}^n$ corresponding to a mass distributed homogeneously over S^{n-1} (cf. Section 3.1), and $h_0(\xi) := \int_{\mathbf{R}^n} e^{i\xi \cdot t} d\sigma(t)$. Then, it is known that $h_0(\xi) = O(|\xi|^{-(n-1)/2})$ ($|\xi| \rightarrow \infty$) (cf. [28, p.154]). By this estimate, we can show the statement in the same way to the proof of Theorem 1. Then, we omit the details.

6 Appendix I : Application to PDEs

Let $1 < p < \infty$, $s \in \mathbf{R}$ and $\psi \in L_{-s}^{p'}(\mathbf{R}^2)$, and define the bounded linear functional ℓ_ψ on $\mathcal{F}L_s^p(\mathbf{R}^2)$ by

$$\ell_\psi(f) := \int_{\mathbf{R}^2} \mathcal{F}[f](\xi) \psi(\xi) d\xi \quad (f \in \mathcal{F}L_s^p(\mathbf{R}^2)),$$

and the support of ℓ_ψ by

$$\bigcap \{F \mid F \text{ is a closed subset of } \mathbf{R}^2 \text{ and } \ell_\psi|_{\mathbf{R}^2 \setminus F} = 0\},$$

where $\ell_\psi|_{\mathbf{R}^2 \setminus F} = 0$ means that $\ell_\psi(f) = 0$ for all $f \in \mathcal{F}L_s^{p'}(\mathbf{R}^2)$ with $\text{supp } f \subset \mathbf{R}^2 \setminus F$. Guo [16, Lemma 1 and Theorem 2'] showed the following : Let $2 \leq p < 4$ and $u \in L^p(\mathbf{R}^2)$. If $\text{supp } \ell_u \subset S^1$ (i.e., $\text{supp } \mathcal{F}[u] \subset S^1$) and u satisfies the Helmholtz equation $\Delta u + u = 0$, then $u = 0$. This result is related to Hörmander [18]. On the other hand, as a by-product of the proof of Theorem 2, we obtain the following:

Corollary 16 Let $2 < p < \infty$ and $2/p < s < 2/p + 1/2$. Then there exists non-zero $u \in L_{-s}^p(\mathbf{R}^2) \cap C(\mathbf{R}^2)$ such that $\text{supp } \ell_u \subset S^1$ and $u + \Delta u = 0$.

Proof Define $q \in (1, 2)$ by $q = p'$, and let J be the closed ideal defined in Section 4. Set

$$A := \{\phi \in \mathcal{F}L_{-s}^{q'}(\mathbf{R}^2) \mid \mathcal{F}[\phi] \in C(\mathbf{R}^2) \text{ and } \langle f, \phi \rangle = 0 \text{ for all } f \in J\}.$$

If we can prove $A \neq \{0\}$, then putting $u := \mathcal{F}[\phi]$ for some non-zero $\phi \in A$ will give us the desired result (cf. Lemma 15). Now, contrary to our claim, we assume $A = \{0\}$. Then we have $\mathcal{F}L_s^q(\mathbf{R}^2) \subset J$. In fact, let $f_0 \in \mathcal{F}L_s^q(\mathbf{R}^2)$. Then, obviously, $\langle f_0, \phi \rangle = 0$ for all $\phi \in A$. Therefore, by Proposition 11, we have $f_0 \in J$. But this is impossible since $J \neq \mathcal{F}L_s^q(\mathbf{R}^2)$. $\square$

7 Appendix II : Closed ideals in $\mathcal{F}L_s^q(\mathbf{R}^2)$

In this section, we shall prove the following theorem.

Theorem 17 Let $1 \leq q \leq 2$, $2/q' < s < 2/q' + 1/2$ and J be the closed ideal in $\mathcal{F}L_s^q(\mathbf{R}^2)$ defined in Section 4. Then, J is the smallest closed ideal I' in $\mathcal{F}L_s^q(\mathbf{R}^2)$ with $Z(I') = S^1$.

7.1 Technical Lemmas

To prove Theorem 17, we recall the following lemmas.

Lemma 18 ([19, Theorem 1.2]) Let $4/3 \leq q \leq \infty$. If $F \in C^\infty(\mathbf{R})$ and $F(0) = 0$, then $F(f) \in \mathcal{FL}_s^q(\mathbf{R}^n)$ for all real-valued $f \in \mathcal{FL}_s^q(\mathbf{R}^n)$.

Lemma 19 ([19, Lemma A.1]) Let $1 \leq p, q \leq \infty$ and $s \in \mathbf{R}$. Suppose that $\varphi \in C_c^\infty(\mathbf{R}^n) \setminus \{0\}$ and $\text{supp } \varphi \subset B_n(x_0, R)$. Then we have

$$\|\varphi \cdot f\|_{M_s^{p,q}} \approx \|\varphi \cdot f\|_{\mathcal{FL}_s^q}.$$

Here, $M_s^{p,q}(\mathbf{R}^n) = M_s^{p,q}$ denotes the modulation space defined by the norm

$$\|f\|_{M_s^{p,q}} = \left\| \langle \xi \rangle^s \left\| \mathcal{F}_{y \rightarrow \xi} [f(y) e^{-|x-y|^2}] (\xi) \right\|_{L^p(\mathbf{R}_x^n)} \right\|_{L^q(\mathbf{R}_\xi^n)}.$$

We refer to [1], [4], [9], [14] for more details on the modulation spaces.

Lemma 20 ([20, Theorem 4.13]) Let $1 < p < \infty$, $1 \leq q \leq \infty$ and $s > n/q'$. Let μ be a complex measure on $\mathbf{R}$ such that

$$\int_{\mathbf{R}} (1 + |\xi|)^{1+(s+\frac{n}{q})(1+\frac{1}{s-\frac{n}{q'}})} d|\mu|(\xi) < \infty$$

and $\mu(\mathbf{R}) = 0$. Furthermore, assume that the function F is the Fourier transform of μ . Then $F(f) \in M_s^{p,q}(\mathbf{R}^n)$ for all real-valued $f \in M_s^{p,q}(\mathbf{R}^n)$.

Making use of the fact that $\mathcal{FL}_s^q(\mathbf{R}^n)$ is a Wiener algebra in the sense of Reiter ([22], [23]), we give a lemma which corresponds to an ideal version of Bhimani-Ratnakumar's [2, Lemma 3.13].

Lemma 21 Let I be a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^n)$, and $f \in \mathcal{FL}_s^q(\mathbf{R}^n)$. Suppose that

- (i) For every $x \in \mathbf{R}^n$, there exists $g_x \in I$ such that $f = g_x$ in a neighborhood of x (i.e., $f \in I$ locally at every $x \in \mathbf{R}^n$).
- (ii) There exists $\psi \in C_c^\infty(\mathbf{R}^n)$ such that $(1 - \psi)f \in I$ (i.e., $f \in I$ locally at ∞).

Then $f \in I$.

Proof Let $\psi \in C_c^\infty(\mathbf{R}^n)$ be a function that satisfies (ii) and divide f as $f = (1 - \psi)f + \psi f$. Since $(1 - \psi)f \in I$ and I is a linear subspace of $\mathcal{FL}_s^q(\mathbf{R}^n)$, to prove $f \in I$, it suffices to show $\psi f \in I$. By virtue of (i), for any $x \in \text{supp } \psi$, there exists $g_x \in \mathcal{FL}_s^q(\mathbf{R}^n)$ such that $f = g_x$ in an open ball $U(x, r_x) := \{y \in \mathbf{R}^n \mid |x - y| < r_x\}$. Since $\text{supp } \psi$ is compact, we can find $\{x_j\}_{j=1}^N \subset \text{supp } \psi$ such that $\text{supp } \psi \subset \cup_{j=1}^N U(x_j, r_{x_j})$. Also, let $\{\phi_{x_j}\}_{j=1}^N \subset C_c^\infty(\mathbf{R}^n)$ be a

partition of unity subordinate to this cover, and let $\{g_{x_j}\}_{j=1}^N \subset I$ be such that $f = g_{x_j}$ on $U(x_j, r_{x_j})$. Since $\text{supp } \phi_{x_j} \subset U(x_j, r_{x_j})$, we have $\phi_{x_j} f = \phi_{x_j} g_{x_j}$. Hence we get

$$\psi f = \sum_{j=1}^N \psi \phi_{x_j} f = \sum_{j=1}^N \psi \phi_{x_j} g_{x_j} \in I.$$

□

Lemma 22 Let $x_0 \in \mathbf{R}^2$ and f be a real-valued function in $\mathcal{FL}_s^q(\mathbf{R}^2)$ such that $f(x_0) = 1$ and $\text{supp } f$ is in a compact neighborhood of x_0 . Then $1/f$ belongs to $\mathcal{FL}_s^q(\mathbf{R}^2)$ locally at x_0 .

Proof Choose a non-negative $v \in C^\infty(\mathbf{R})$ such that $v(x) = 1$ on $B_1(1, 1/10)$ and $\text{supp } v \subset B_1(1, 1/3)$. Define F by $F(x) = v(x)/x$ ($x \in \text{supp } v$), and $= 0$ ($x \notin \text{supp } v$). We first consider the case $q \geq 4/3$. Since $F \in \mathcal{S}(\mathbf{R})$ and $F(0) = 0$, Lemma 18 shows that $F(f) \in \mathcal{FL}_s^q(\mathbf{R}^2)$. Moreover, since $f \in \mathcal{FL}_s^q(\mathbf{R}^2) \subset C(\mathbf{R}^2)$ and $v(x) = 1$ on $B_1(1, 1/10)$, we see $F(f) = 1/f$ in some neighborhood of x_0 , and thus $1/f$ belongs to $\mathcal{FL}_s^q(\mathbf{R}^2)$ locally at x_0 . Next, we consider the case $1 < q < 4/3$. Since $F \in \mathcal{FL}_\alpha^1(\mathbf{R})$ for all $\alpha \in \mathbf{R}$, Lemma 20 shows that $F(f) \in M_{s,q}^{p,q}(\mathbf{R}^2)$. On the other hand, by the definitions of v and f , we can easily see $\text{supp } F(f)$ is compact. Thus Lemma 19 yields $F(f) \in \mathcal{FL}_s^q(\mathbf{R}^2)$. Moreover, since $F(f) = 1/f$ in some neighborhood of x_0 , $1/f$ belongs to $\mathcal{FL}_s^q(\mathbf{R}^2)$ locally at x_0 .

□

7.2 The proof of Theorem 17

Let I' be a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^2)$ with $Z(I') = S^1$. Then, it suffices to prove $J \subset I'$. Before we prove this, we remark that there exist $x_0 \in \mathbf{R}^2 \setminus S^1$ and $h \in I'$ such that $h(x) = 1$ on a neighborhood of x_0 . In fact, we first note that there exist $x_0 \in \mathbf{R}^2 \setminus S^1$ and $f \in I'$ such that $f(x_0) = 1$ by $Z(I') = S^1$. Additionally, since I' is a closed ideal in $\mathcal{FL}_s^q(\mathbf{R}^2)$, by taking $\phi \in C_c^\infty(\mathbf{R}^2)$ with $\phi(x_0) = 1$ and considering $f\phi$ instead of f , we may assume $\text{supp } f$ is compact. Moreover, since $\bar{f} \in \mathcal{FL}_s^q(\mathbf{R}^2)$, we see $|f|^2 = f\bar{f} \in I'$ and $|f(x_0)|^2 = 1$. Therefore, Lemma 22 shows that there exists $k \in \mathcal{FL}_s^q(\mathbf{R}^2)$ such that $k(x) = 1/|f(x)|^2$ in some neighborhood of x_0 . Hence, $h := k|f|^2 \in I'$ and $h(x) = 1$ in some neighborhood of x_0 . Now we prove $J \subset I'$ by Lemma 21. By Lemma 7 and the definition of J (see also (15)), it suffices to show $f \in I'$ for all $f \in J^0$ with compact support. Let $f \in J^0$ and $\text{supp } f$ is compact. Since $0 \in I'$ and $f = 0$ in a neighborhood of S^1 , f belongs to I' locally near S^1 . Moreover, if $x_0 \in \mathbf{R}^2 \setminus S^1$, then there exists $k \in I'$ such that $k(x) = 1$ near x_0 . Hence, we have $f = k \cdot f$ near x_0 and $k \cdot f \in I'$, which shows f belongs to I' locally near $\mathbf{R}^2 \setminus S^1$. We also note $f \in I'$ locally at ∞ , since $\text{supp } f$ is compact. Hence, f is in I' at all point which contains within $\mathbf{R}^2$ and ∞ , which implies $f \in I'$.

Acknowledgment

We would like to thank Professor Hans Georg Feichtinger for several valuable comments and suggestions which improve our paper. We are also grateful to the referee for valuable advice and suggestions. Remark 1 is due to the referee. This work was supported by JSPS KAKENHI Grant Numbers 22K03328, 22K03331, 25K07021.

References

- [1] Bényi, Á.; Okoudjou, K.A.: Modulation Spaces. With Applications to Pseudodifferential Operators and Nonlinear Schrödinger Equations. Applied and Numerical Harmonic Analysis. New York, NY: Birkhäuser. xvi, 169 p. (2020).
- [2] Bhimani, D.G.: Ratnakumar, P.K.: Functions operating on modulation spaces and nonlinear dispersive equations. J. Funct. Anal. 270, No. 2, 621-648 (2016).
- [3] Brandenburg, L. H.: On identifying the maximal ideals in Banach algebras. J. Math. Anal. Appl. 50, 489-510 (1975).
- [4] Cordero, E.; Rodino, L.: Time-Frequency Analysis of Operators. De Gruyter Studies in Mathematics 75. Berlin: De Gruyter. xiv, 442 p. (2020).
- [5] Domar, Y.: Harmonic analysis based on certain commutative Banach algebras. Acta Math. 96, 1-66 (1956).
- [6] Domar, Y.: On the spectral synthesis problem for $(n - 1)$ -dimensional subsets of $\mathbf{R}^n$, $n \geq 2$. Ark. Mat. 9, 23-37 (1971).
- [7] Domar, Y.: On spectral synthesis in $\mathbb{R}^n$, $n \geq 2$. Euclidean harmonic analysis, Proc. Semin., Univ. Maryland 1979, Lect. Notes Math. 779, 46-72 (1980).
- [8] Feichtinger, H. G.: Gewichtsfunktionen auf lokalkompakten Gruppen. Sitzungsber., Abt. II, Österr. Akad. Wiss., Math.-Naturwiss. Kl. 188, 451-471 (1979).
- [9] Feichtinger, H.G.: Modulation spaces on locally compact abelian groups, in: M. Krishna, R. Radha and S. Thangavelu (Eds.), Wavelets and their Applications, Chennai, India, Allied Publishers, New Delhi, 2003, 99-140, Updated version of a technical report, University of Vienna, 1983.
- [10] Feichtinger, H.G.: Personal communication (2024)
- [11] Feichtinger, H.G., Kobayashi, M., Sato, E.: On some properties of modulation spaces as Banach algebras. Stud. Math. 280, No. 1, 55-86 (2025).
- [12] Feichtinger, H.G., Kobayashi, M., Sato, E.: Further study of modulation spaces as Banach algebras. Ann. Univ. Sci. Budap. Rolando Eötvös, Sect. Comput. 56, 151-166 (2024).
- [13] Grafakos, L.: Classical Fourier Analysis. 3rd ed. Graduate Texts in Mathematics 249. New York, Springer. xvii, 638 p. (2014).
- [14] Gröchenig, K.: Foundations of Time-Frequency Analysis. Applied and Numerical Harmonic Analysis. Birkhäuser Boston, Inc., Boston, MA, (2001)

- [15] Gröchenig, K.: Weight functions in time-frequency analysis. *Pseudo-Differential Operators: Partial Differential Equations and Time-Frequency Analysis*, 343–366. Fields Inst. Commun., 52 American Mathematical Society, Providence, RI, (2007).
- [16] Guo, K.: On the spectral synthesis property and its application to partial differential equations. *Ark. Mat.* 30, No. 1, 93-103 (1992).
- [17] Herz, C. S.: Spectral synthesis for the circle. *Ann. Math.* (2) 68, 709-712 (1958).
- [18] Hörmander, L.: Lower bounds at infinity for solutions of differentiable equations with constant coefficients, *Israel J. Math.* 16, 103-116 (1973).
- [19] Kato, T., Sugimoto, M., Tomita, N.: Nonlinear operations on a class of modulation spaces. *J. Funct. Anal.* 278, No. 9, Article ID 108447, 26 p. (2020).
- [20] Reich, M., Sickel, W.: Multiplication and composition in weighted modulation spaces. Qian, Tao (ed.) et al., *Mathematical Analysis, Probability and Applications – Plenary Lectures. Springer Proceedings in Mathematics & Statistics* 177, 103-149 (2016).
- [21] Reiter, H.: Contributions to harmonic analysis. IV. *Math. Ann.* 135, 467-476 (1958).
- [22] Reiter, H.: *Classical Harmonic Analysis and Locally Compact Groups*. Oxford: At the Clarendon Press. XI, 200 p. (1968).
- [23] Reiter, H., Stegeman, J. D.: *Classical Harmonic Analysis and Locally Compact Groups*. 2nd ed. London Mathematical Society Monographs. New Series. 22. Oxford: Clarendon Press. xiii, 327 p. (2000).
- [24] Rudin, W.: *Fourier Analysis on Groups*. Interscience Tracts in Pure and Applied Mathematics. 12. New York and London: Interscience Publishers, division of John Wiley and Sons. IX, 285 p. (1962).
- [25] Schwartz, L.: Sur une propriété de synthese spectrale dans les groupes non compacts. *C. R. Acad. Sci., Paris* 227, 424-426 (1948).
- [26] Stegeman, J. D.: Wiener - Ditkin sets for certain Beurling algebras. *Monatsh. Math.* 82, 337-340 (1976).
- [27] Stegeman, J. D.: Analysis of P. Malliavin's proof of non spectral synthesis. *New Integrals, Proc. Henstock Conf., Coleraine/Ireland 1988, Lect. Notes Math.* 1419, 180-200 (1990).
- [28] Stein, E. M., Weiss, G.: *Introduction to Fourier Analysis on Euclidean Apaces*. Princeton Mathematical Series 32. Princeton, N. J.: Princeton University Press. X, 297 p. (1971).

- [29] Varopoluos, N. T.: Spectral synthesis on spheres. Proc. Camb. Philos. Soc. 62, 379-387 (1966).
- [30] Watson, G. N.: A Treatise on the Theory of Bessel Functions. 2nd ed. London: Cambridge University Press. vii, 804 p. (1966).
- [31] Wermer, J.: On a class of normed rings. Ark. Mat. 2, 537-551 (1954).