HOKKAIDO UNIVERSITY

Title	Modeling alternative strategies of male emergence timing
Author(s)	Kubo, Hidaka; Nakaoka, Shinji; Yamaguchi, Ryo
Citation	Journal of theoretical biology, 612, 112214 https://doi.org/10.1016/j.jtbi.2025.112214
Issue Date	2025-09-07
Doc URL	https://hdl.handle.net/2115/98058
Rights	© 2025. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	Caption_Fig_1.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Fig1 Box plots showing forewing length among early male, late male, and female individuals of *Fabriciana nerippe*. The horizontal axis indicates the categories of data. The vertical axis displays forewing length. The bold horizontal line in each box is the median, and the top and bottom of the box are the first and third quartiles, respectively. The whiskers extend to the maximum and minimum values observed in the data. "n" represents the number of samples in each category. The bars and asterisks at the top of the box represent the result of the Tukey-Kramer test. (*** ; $p<0.001$). The photographs of the sample individuals shown under each category were taken at the same scale.